

PAX6 (+5a) Expression in Adipose Tissue-Derived Mesenchymal Stem Cells Induces Differentiation to Retinal Ganglion Cells

Habib Rezanejad¹, Farhang Haddad^{1*}, Zahra-Soheila Soheili^{2**}, Maryam M. Matin^{1,3}, Shahram Samiei⁴, and Sepideh Zununi Vahed⁵

1. Department of Biology, Faculty of Science, Ferdowsi University of Mashhad, Mashhad, Iran
2. National Institute of Genetic Engineering and Biotechnology, Tehran, Iran
3. Cell and Molecular Biotechnology Research Group, Institute of Biotechnology, Ferdowsi University of Mashhad, Mashhad, Iran
4. Iranian Blood Transfusion Organization and Research Center, Tehran, Iran
5. School of Advanced Biomedical Sciences, Tabriz University of Medical Sciences, Tabriz, Iran

Received 26 December 2013

Accepted 17 January 2014

Abstract

Glaucoma remains one of the major causes of blindness in today's world. The progressive field of stem cells proposes an exciting potential for discovering novel therapies. Here, we report the development of an easy and high throughput method for differentiation of retinal ganglion cells (RGCs) and bipolar cells from human adipose tissue-derived mesenchymal stem cells (hADSCs) using *PAX6* (+5a) gene expression, a master gene in development of the vertebrate visual system. HADSCs were isolated from fat tissues and confirmed by their cell surface markers and differentiation potential into adipocytes and osteocytes lineages. Then, the coding region of human *PAX6* (+5a) gene was cloned and lentiviral particles were produced. HADSCs differentiation was characterized by morphological characteristics, qRT-PCR and immunocytochemistry (ICC). The hADSCs were isolated successfully with high yield and purity (99%). After 30 hours post transduction by pLEX-*PAX6* - pur lentiviral vectors in fibronectin supplemented medium, cells gradually showed the characteristic morphology of neuronal cells. QRT-PCR and ICC confirmed deriving of mainly RGCs and marginally bipolar cells. The current investigation demonstrates the feasibility of differentiation of RGCs and bipolar cells from hADSCs using expression of *PAX6* (+5a) in the medium supplemented by fibronectin.

Keywords: *PAX6* gene; adipose tissue-derived mesenchymal stem cells; retinal ganglion cell; transdifferentiation

Introduction

Glaucoma, a chronic retinal neurodegenerative disease, is the second cause of worldwide blindness in developed countries (Quigley and Broman, 2006). It has been estimated that 80 million people worldwide would have been affected by this disease by 2020. Glaucoma is characterized by the degeneration of axons in the optic nerve and apoptosis in retinal ganglion cells (RGCs) (Kuehn et al., 2005). RGCs are the first differentiating retinal cells in all vertebrates that are induced from retinal progenitor cells in embryonic development (Marquardt and Gruss, 2002). The inability of the central nervous system to regenerate new cellular components in response to damage leads to a limited capacity of structural and functional repair in the retina (Cao et al., 2002).

One of the strategies to restore vision in glaucoma patients is the functional replacement of

RGCs using autologous or heterologous transplantation (Baker and Brown, 2009; MacLaren et al., 2006; Moshiri et al., 2004; Wallace, 2007; Wong et al., 2011). For the generation and transplantation of RGCs and their precursors, the potential of different sources of stem cells, including human embryonic stem cells (hESC), bone marrow-derived stem cells, umbilical cord-derived cells, induced pluripotent stem cells (iPS), adult human Müller stem cells and fetal stem cells, have been reported (Baker and Brown, 2009; Buchholz et al., 2013; Jayaram et al., 2011; John et al., 2013; Ramsden et al., 2013; Wallace, 2007).

Adipose tissue represents an abundant and accessible source of adult stem cells (Jurgens et al., 2008; Kokai et al., 2005). A growing body of experimental evidence, from *in vitro* and *in vivo* studies, demonstrates the multipotentiality of adipose-tissue derived stem cells (ADSCs) (Gimble et al., 2007).

Corresponding authors E-mail:

*hadad@um.ac.ir

**soheili@nigeb.ac.ir

While ADSCs are similar to bone marrow stem cells in differentiation and therapeutic potential, they are much easier and safer to obtain in large quantities which make them an ideal and reliable source for regenerative medicine (Rasmussen et al., 2012). The Paired box (PAX) genes are members of the family of tissue specific transcription factors. This family plays a critical role in the formation of tissues and organs during embryonic development and it has vital functions in certain tissues in adults (Hever et al., 2006; Pichaud and Desplan, 2002). *PAX6* along with other genes, such as *SOX2* and *OTX2*, control each stage of eye development and it has been called the master gene for eye development (Hever et al., 2006). *PAX6* mutations in *Drosophila*, mouse, rat, and human demonstrate its requirement for the development of eye (Philips et al., 2005; Yasuda et al., 2002). *PAX6* (-5a) and *PAX6* (+5a) are two isoforms of the conserved *PAX6* gene that have different DNA binding specificities and functions (Azuma et al., 2005). It has been reported that *PAX6* (-5a) plays important roles in both embryogenesis and adult body homeostasis and *PAX6* (+5a) is the most critical isoform that promotes the neuronal differentiation of murine embryonic stem cells (Shimizu et al., 2009).

Based on these findings, it was considered that *PAX6* (+5a) transfection into hADSCs may induce retinal neurons, including RGCs and/or their precursors. In so doing, the expression of human *PAX6* (+5a) by lentiviral expression vectors was employed in hADSCs under a culture medium supplemented with fibronectin. This study offers an effective method for *in vitro* induction of retinal ganglion like cells that can be used in stem cell based therapy.

Materials and Methods

Adipose tissue sampling

Human adipose tissues were obtained from abdominal subcutaneous tissues of patients undergoing abdominoplasty procedures, in Tehran Medical University (N=6, age: 20-35 years). Before the surgical procedure, informed consents were obtained from the patients. About 150 ml of lipoaspirate was gathered in a sterile bottle, filled with 0.1 M phosphate-buffered saline (PBS) or DMEM(Dulbecco's modified Eagle's medium)-F12 (Sigma, Germany), in order to achieve enough number of cells.

Isolation and cell culture of hADSCs

HADSCs isolation was done according to

previously published methods (Estes et al., 2010) with some modifications. Up to 2×10^7 adipose stromal stem cells with more than 98% purity were isolated from 150 ml of lipoaspirates; however, yields varied among patient's samples. Briefly, in order to remove the majority of erythrocytes and leukocytes, the lipoaspirate was washed five times with sterile PBS containing 120 µg/ml penicillin (Fluka, China) and 220 µg/ml streptomycin (Fluka, China). Then, 0.1% (wt/vol) collagenase type I (Invitrogen, USA) was used to digest the extracellular matrix. Enzyme activity was neutralized with DMEM supplemented fetal bovine serum (FBS) 10% (Gibco, Germany) and centrifuged at 1500 rpm for 10 min to obtain a high density pellet. The cell pellet was re-suspended and extensively washed with PBS. Finally, remaining cells were cultured in 25 cm² flasks (Nunc, Denmark). DMEM-F12 medium supplemented with 10% FBS, 5 ng/ml human epidermal growth factor (Roche, Germany), 1 ng/ml human fibroblastic growth factor (Roche, Germany), 100 U/ml penicillin and 100 mg/L streptomycin was used as culturing and expansion medium. The flasks were then incubated at 37 °C with 5 % CO₂ in a humidified atmosphere. The medium was changed after 16 hours, and then twice a week. Adherent cells were harvested with 0.25% trypsin-0.02% EDTA, and re-plated at a dilution of 1:3 when the confluency was more than 80%.

Flowcytometric cell surface marker expression analysis

To characterize the phenotype of the cultured cells with flow cytometry, fluorescein isothiocyanate (FITC)-conjugated primary antibodies for human CD44, CD45, CD73, CD90 and CD105 (BD Biosciences, USA) were used. The hADSCs were trypsinized and washed three times with cooled PBS containing 2% FBS and sodium azide. Cells (5×10^4) were incubated with aforementioned antibodies. All antibodies were diluted 1:1000 and incubated with cells for 45 min at 4 °C. Then the cells were washed with PBS containing 2% FBS. After two washing steps, cells were re-suspended in 500 µl paraformaldehyde 2% containing 1% FBS for profile characterization and analyzing by fluorescence-activated cell sorting (FACS) system (Partec II, Germany).

Analysis of multipotent differentiation capacity

When over 80% confluency was reached, cells were incubated in the osteogenic and adipogenic differentiation media for four and three weeks, respectively. Osteogenic differentiation medium consisted of DMEM-F12, 20% FBS with

osteogenic supplement: 100 nM dexamethasone (Sigma-Aldrich, Germany), 50 µg/ml ascorbate-2-phosphate (Sigma-Aldrich, Germany) and 10 mM β-glycerolphosphate (Sigma-Aldrich, Germany). Adipogenic differentiation medium consisted of DMEM-F12 supplemented with 10% FBS, 100 nM dexamethasone (Sigma-Aldrich, Germany), 50 µg/ml indomethacin (Sigma-Aldrich, Germany) and 50 µg/ml ascorbate- 2 phosphate (Sigma-Aldrich, Germany). In general, the culture media were changed every 4 days. After induction, alkaline phosphatase assay (Sigma-Aldrich, Germany) and Oil Red O (Sigma-Aldrich, Germany) staining were performed to confirm the differentiation of hADSCs toward osteogenic and adipogenic lineages, respectively.

Construction of vectors

The coding sequence of human *PAX6* (+5a) gene was synthesized and cloned into pUC57 cloning vector. Vector was digested by *Bam*HI and *Xho*I restriction enzymes and subcloned into the *Bam*HI/*Xho*I site of pLEX-MCS-Pur lentiviral expression vector and was designated as pLEX-*PAX6* -Pur construct. The recombinant construct was confirmed by PCR amplification, digestion and finally DNA sequencing.

Virus particles production

Lentiviral particles were produced by calcium-phosphate transient transfection using a vector expression system in HEK 293T cells. Briefly, HEK 293T cells were plated in 6-cm plates with 7×10^5 cells in 4 ml of DMEM-HG (high-glucose) supplemented with 10% FBS. Then pLEX-*PAX6* -Pur vector, as transfer vector (11 µg), envelope encoding plasmid (4 µg) and packaging lentivirus vector (7 µg) were added and transfection was carried out using CaCl₂. Fourteen hours after transfection, medium was replaced, and the supernatant was collected 24 and 48 hours post transfection. Finally, assembled lentiviral particles were filtered, purified and concentrated by PEG 6000 (Sigma, Germany).

Transduction of hADSCs

HADSCs were transduced by 8 hour exposure to the viral supernatant in the presence of 8 µg/ml polybrene at 37 °C and 5% CO₂ without FBS. 72 hours post transduction, selection medium containing DMEM-F12 with 20% FBS and 1 µg/ml puromycin was added to the transduced-hADSCs. Every 3 days the selective medium was changed with the same medium. Transduced-hADSCs from passages 2 post transduction were analyzed for eGFP expression, cell proliferation and cell death

using ELISA kits (Roche, Germany) according to the manufacturer's instructions.

Real-time RT-PCR assay

Total RNA extraction and cDNA synthesis were performed using RNeasy kit (Qiagen, Germany), and Quantiscript® reverse transcriptase (Qiagen, Germany), respectively. Quantitative real-time PCR was performed with the Corbet Real-Time PCR system (Applied Biosystems, USA). Specific primers were used from the Quantitect primer assay (Qiagen, Germany) (Table 1). Data were normalized to the expression of *GAPDH*, a housekeeping gene, which has shown to have stable expression under different experimental conditions in similar studies. Each reaction contained 5 µl of Quantifast syber green master mix, 1 µl of forward and reverse mix primers (10 pM), 3 µl of RNase free water and 1 µl of cDNA. The reactions were conducted with initial enzyme activation at 95 °C for 5 min, followed by 45 cycles of denaturation at 95°C for 10 seconds and anneal at 60 °C for 30s. Relative gene expression was calculated using Bio-Rad software (RelQuant UpDate- for relative quantification) according to the 2-ΔΔCt method based on the threshold cycle (Ct) values (Schmittgen and Livak, 2008). All experiments were performed in duplicate and their values were presented as mean±SD. Student's t-test was used to evaluate the statistical significance of the data; $p < 0.05$ was considered statistically significant.

Table. 1 primers list used for qPCR from quantitect primer assay (Qiagen, Germany)

	Official symbol	Amplicon length (bp)	Cat. No.
1	<i>PAX6</i>	113	QT00071169
2	<i>FGF2</i>	109	QT00047579
3	<i>DVL3</i>	69	QT00999810
4	<i>SHH</i>	136	QT00205625
5	<i>SOX1</i>	96	QT00215299

Immunocytochemistry

Immunocytochemical analysis was carried out for the detection of cell specific markers according to the Santa Cruz protocol with some modifications. Briefly, transduced hADSCs were cultured on FBS pre-coated glass cover slips in a 24-well microplate at a density of 6×10^4 cells per well and washed with PBS. Paraformaldehyde fixed cells were permeabilized with chilled methanol (Merck, Germany) and then blocked in 1% BSA (Merck) in PBST (1% Triton X-100 in PBS) (Sigma) for 45 min at room temperature, followed by 1 h incubation in primary antibodies at room temperature (All antibodies were obtained from

Santa Cruz, USA). Antibodies for retinal progenitor and retinal ganglion and bipolar cell markers included the goat polyclonal anti-human *PAX6*, *Thy1* and *PKC*. A negative secondary antibody-only control was also included. Nuclei were counter-stained with DAPI (1 mg/ml, Santa Cruz, USA) to assess the total number of cells in each field. Cover slips were then mounted onto slides using an anti-fading mounting medium (90% glycerol, 10% PBS and 10% (w/v) phenylene-diamine). Samples were observed under the Axiophot Zeiss fluorescence microscope (Germany) with a 460 nm filter for DAPI and a 520 nm filter for FITC-conjugated antibodies, and digital pictures were taken.

Results

C hADSCs

The hADSCs were isolated successfully from human adipose tissues. Approximately 5×10^6 hADSCs were obtained from one gram of adipose tissue. Cells were cultured in expansion medium and passaged every 3-4 days. Human ADSCs were large, spindle-shaped cells with fibroblastic features (Figure 1A, B). Thin cell body of cells contained a large and round nucleus (Figure 1C, D). In early passages, cells displayed clonogenic properties, the ability of a single cell to proliferate independently to form a colony (Figure 1E, F). This indicated the renewing capacity of isolated stem cells. Cells kept their morphological features, without major alteration, for a maximum of 11 passages.

Analysis of multipotent differentiation capacity

Osteogenic and adipogenic differentiation capacity of hADSCs were examined. Osteogenic differentiation was confirmed by mineralization of cells in osteogenic medium at week 4, which could be observed by alizarin red staining (Figure 1 G, H). Long spindle-shape morphology of the hADSCs changed into a polygonal shape 4 days after incubation in adipogenic induction medium. By day 9, small droplets of oil lipid appeared in some of the cells. After three weeks, most of the differentiated cells showed red lipid droplets throughout the cytoplasm, which was confirmed by oil red O staining (Figure 1 I, J).

Analysis of cell surface marker

To characterize cell surface markers of isolated cells, flow cytometry was performed. The flow cytometric analysis demonstrated that approximately 99% of hADSCs expressed the surface markers CD44, CD73 and CD105 (Figure 2A-C, G). The hADSCs lacked the expression of the hematopoietic markers CD34 and CD45 (Figure 2E, F). Results verified the mesenchymal origin of the hADSCs and the lack of hematopoietic markers. Each value represents the mean of two independent experiments in at least duplicate.

Successfully gene transduction of hADSCs by lentiviral vectors

Two lentiviral vectors pLEX-eGFP-pur and




Figure 1. Characteristics of isolated hADSCs. Morphological features of cells (A-D); the hADSCs were typical fibroblast-like cells with fusiform shape from the 2nd passage and preserved their shape after expansion in vitro (A, B). Nuclei were stained with DAPI (blue), the cell body contained a large and round nucleus (C, D). Clonogenic capacity of hADSCs (E, F). Multipotential differentiation assays of hADSCs (G-J); differentiation potential of hADSCs towards osteogenic lineages was assessed through alkaline phosphatase activity assay (G, H), hADSCs differentiated towards the adipogenic lineage and formed lipid vesicles, which were stained using oil red-O (I, J). (C 100X – all other 200 X)

pLEX-PAX6 -pur, were constructed and transferred into HEK293T and hADSCs. Transduction efficiency of hADSCs was examined by expression of GFP. Lentiviral vector, pLEX-eGFP-pur, was used to transduce proliferating hADSCs at a MOI of 100. One day after transduction, 90% and 75% of HEK293T and hADSCs expressed GFP, respectively. Although, the expression of GFP was observed only 24 hour post transduction in HEK293 cells, it took about 3 days to observe GFP in hADSC cells (Figure 3 A, B).

Cell morphology characteristics after transduction

Three days post transduction, cells showed the characteristic morphology of neuronal cells and little axon-like processes emerged gradually. Four days post transduction, they gradually extended axon-like processes that finally led to the formation of neural-network-like structures. Cells had multiple dendrites with a long axon and a fat cell body resembling reliable RGCs (Figure 3 C-F).




Figure 2. Flow cytometry analysis of hADSCs isolated from fat tissue and cultured *in vitro*. Immunophenotypic characterization of hADSCs (A-F), with cells positively expressing the antigens CD44 (A), CD73 (B) and CD105 (C), while negatively expressing the antigens CD34 (E) and CD45 (F), 99% of isolated hADSCs expressed the surface markers: CD44, CD73 and CD105 (G).




Figure 3. Transduction of hADSCs with lentiviral vectors. Efficient eGFP transduction of cells (A, B); HEK293T (A) and hADSCs (B) transduced with pLEX-eGFP-pur vector were expressing GFP. Cell morphology of transduced cell at day 4 under inverted microscopic view (C-F) (A 100X, all others 300X).

Real time RT-PCR analysis

To test the expression of retinal neuron and RGC-associated mRNAs, including *SOX1*, *PAX6*, *Thy1* and *PKC*, qRT-PCR was performed. According to the qRT-PCR data, *PAX6* and *SOX1* expression levels increased substantially after transduction compared to cells transduced by empty vector or without transduction. *Thy1*, a marker of RGCs, plays an important role in the formation of the visual system. Expression of *FGF2* and *sonic hedgehog (SHH)* as molecules involved in different signaling pathways in differentiation of retinal neural cells was also examined. Results showed the increased expression of *FGF2* and *SHH* (Figure 4). At the mRNA level, expression of *PKC*, a bipolar cell marker, was also increased. Taken together, these results indicated that *PAX6* (+5a) induction of hADSCs can cause differentiation into mostly RGCs and marginally bipolar cells.




Figure 4: Relative retinal gene expression in *PAX6* (+5a)-transduced hADSCs after 6 days in comparison to the cells transduced by pLEX-MCS-pur (empty vector). The results represent the mean of 3 independent experiments in at least triplicate.

ICC confirmed the expression of some key markers of RGCs.

To determine whether *PAX6* (+5a) transcription factor and fibronectin were able to induce hADSCs into retinal cells, we examined the expression of *PAX6*, *Thy1* and *PKC* markers in *PAX6* (+5a)-transduced cells. Using ICC method, cells were positive for *PAX6* ($64 \pm 1.3\%$), *Thy1* ($32 \pm 1.1\%$) and *PKC* ($29 \pm 1.4\%$) (Figure 5 A-F). Cells transfected with empty vector did not show any of these markers.

Discussion

Glaucoma is the most common cause of irreversible blindness in the world. Despite the advances in the currently available treatments, many patients experience significant visual loss due to degeneration of RGC (Kerrigan–Baumrind et al., 2000; Kuehn et al., 2005; Quigley and Broman, 2006). Right now, there is no therapeutic strategy for functional recovery of these cells. Cell therapy offers an alternative treatment for restoring the damaged cells in neurodegenerative diseases (Buchholz et al., 2013; Cao et al., 2002; Haddad-Mashadrizeh et al., 2013; Huang et al., 2013; Huang et al., 2011; John et al., 2013). It is acknowledged that damage to the neural retina during glaucoma is restricted to the degeneration of RGCs (Kerrigan–Baumrind et al., 2000); therefore, replacement of these cells might be possible and, if so, might restore the optic nerve.

Several groups have focused on the differentiation of stem cells from different sources with various methods (Fraichard et al., 1995;




Figure 5. Fluorescence microscopy of *Thy1*, *PKC* and *PAX6* in *Pax6* (+5a)-transduced cells after 6 days. These *Pax6* (+5a)-transduced cells were analyzed by in situ immunostaining with antibodies raised against different RGC markers. A,B) ICC for *Thy1* C,D) Antibody against *PKC* E,F) antibody against *Pax6*. Each value represents the mean of 3 independent experiments in at least triplicate.

Jagatha et al., 2009; Jin et al., 2009; Osakada et al., 2009; Singhal et al., 2012; Wong et al., 2011). Most of these sources have limitations for human RGC replacement in clinics (Jayaram et al., 2011). Consequently, identifying alternative sources of cells that replace these cells in the glaucomatous eye without ethical and practical restrictions is necessary. With this target in mind, this study aimed to address some of these limitations.

This study established an effective and easy way to obtain high-yield hADSCs with high purity, 99% positive staining rate observed from results of multiple cell surface markers, using collagenase digestion and adherence screening. No significant difference was seen among different passages in the phenotypes of hADSCs, indicating that the cells can be stably amplified *in vitro* for several passages. High proliferation (data not shown) and differentiation (Figure 1G-J) capacity of isolated hADSCs is consistent with stem cell characteristics. It is reported that fibronectin may be important for differentiating ESCs into retinal neuron precursors, including RGC-like cells (Kayama et al., 2010). Based on these findings, culture medium was supplemented with fibronectin for neural cell induction. Results showed that *PAX6* (+5a) expression and fibronectin supplemented medium are sufficient to induce the differentiation of retinal precursor cells from hADSCs.

Inverted microscopic examination disclosed the appearance of generally long cells that exhibited multidendrites and one axon per cell, suggesting their neuronal differentiation (Figure 3C-F). Several genes expressed in differentiating RGCs such as *SOX1*, *PAX6*, and *Thy1* were tested by qPCR. The results revealed that upon differentiation, the cells had up-regulated expression of early neural markers, *SOX1* and *PAX6*. *SOX1* is one of the earliest transcription factors that is expressed in cells committed to the neural fate (Pevny et al., 1998). *PAX6* is a neural/retinal progenitor marker and acts as a master switch for activation of RGC regulator, thereby supposed to initiate the RGC differentiation cascade (Jagatha et al., 2009). *Thy1*, a surface glycoprotein, is uniquely expressed in RGCs in retina (Huang et al., 2006).

Different signaling molecules such as *FGF2* and *SHH* have been shown to be involved in differentiation of RGCs. It has been shown that *FGF2* is a potent stimulator of axon growth during RGC development (Sapieha et al., 2003). Sonic hedgehog (*Shh*) has been shown to play an

important role in the development of the retina in a number of different model organisms (Spence et al., 2004). Recent studies have demonstrated that the signaling molecule *Shh* secreted by differentiated RGCs is required to promote the progression of ganglion cell differentiation. *Shh* plays dual roles to orchestrate the progression of retinal neurogenic wave (Zhang and Yang, 2001) and also plays a major role in RGC axon projection inside the retina (Kolpak et al., 2005). *FGF2* and *SHh* are known to activate *PAX6* (Jagatha et al., 2009). Increased levels of *FGF2* and *SHh* and *PAX6* in this study were in line with these findings.

Since RGCs collect the messages from bipolar cells and represent the ultimate signals to the vision center in the brain, mRNA expression of *PKC* (protein kinase C), a bipolar cell marker, was also examined. Expression of *SOX1*, *PAX6*, *FGF2*, *Thy1*, *SHH* along with ICC results of *PAX6*, *Thy1* and *PKC* confirmed that the differentiated cells belonged to RGCs and bipolar cells (Figure 4 and 5). Major disadvantages associated with the use of different sources of stem cells are their ethical concerns, shortage of donor cells, limited availability, inflammation, immunoreaction as well as their safety issues regarding teratoma formation; therefore, their potential in cell therapy may be problematic (Wong et al., 2011). Human ADSCs can overcome some of these problems.

Aside from developing a reliable source of retinal cells for transplantation, there are several additional obstacles that have to be considered. For instance, limitation of integration of graft cells into host tissue (Wong et al., 2011), ongoing disease in the host environment that may present a problem for cell transplantation (Wallace, 2007) and tumorigenicity, specially, when cell cultivation period has prolonged (Wong et al., 2011). Moreover, after the transplantation of RGCs, it would need the additional challenge of regrowth of axons through the optic nerve to targets in the brain (Wallace, 2007).

Taken together, for successful retinal regeneration, improved methods for purifying donor retinal cells, optimizing host conditions, as well as using animal models of human diseases, to determine the efficacy and safety of treatments, will be crucial. Furthermore, in order to optimize the best minimal cocktail requested to achieve more authentic-differentiated neurons, more growth factors, cytokines, mRNA, microRNA and small molecules deserve to be investigated.

Future studies to determine the markers of differentiated cells and also time course expression of aforementioned genes are under investigation. Using current method, bipolar cells and RGCs beside different retinal cells including photoreceptors (our previous work, in press) were successfully differentiated from hADSCs. For an efficient differentiation of hADSCs to disease-relevant cell types, novel strategies need to be developed. The current investigation demonstrates the feasibility of the differentiation of RGCs and bipolar cells from hADSCs using expression of *PAX6* (+5a) in the medium supplemented by fibronectin that can be used in stem cell therapy.

Acknowledgement

We would like to thank Ferdowsi University of Mashhad, Mashhad, Iran for their Grant # 30368 to perform this work.

References

- 1- Azuma N., Tadokoro K., Asaka A., Yamada M., Yamaguchi Y., Handa H., Matsushima S., Watanabe T., Kohsaka S., Kida Y., Shiraishi T., Ogura T., Shimamura K. and Nakafuku M. (2005) The *PAX6* isoform bearing an alternative spliced exon promotes the development of the neural retinal structure. *Human molecular genetics* 14:735-745.
- 2- Baker P. S. and Brown G. C. (2009) Stem-cell therapy in retinal disease. *Curr Opin Ophthalmol* 20:175-181.
- 3- Buchholz D. E., Pennington B. O., Croze R. H., Hinman C. R., Coffey P. J. and Clegg D. O. (2013) Rapid and efficient directed differentiation of human pluripotent stem cells into retinal pigmented epithelium. *Stem Cells Translational Medicine* 2:384-393.
- 4- Cao Q., Benton R. L. and Whittemore S. R. (2002) Stem cell repair of central nervous system injury. *Journal of neuroscience research* 68:501-510.
- 5- Estes B. T., Diekman B. O., Gimble J. M. and Guilak F. (2010) Isolation of adipose-derived stem cells and their induction to a chondrogenic phenotype. *Nat Protoc* 5:1294-1311.
- 6- Fraichard A., Chassande O., Bilbaut G., Dehay C., Savatier P. and Samarut J. (1995) *In vitro* differentiation of embryonic stem cells into glial cells and functional neurons. *J Cell Sci* 108 (Pt 10):3181-3188.
- 7- Gimble J. M., Katz A. J. and Bunnell B. A. (2007) Adipose-derived stem cells for regenerative

medicine. *Circulation research* 100:1249-1260.

8- Haddad-Mashadrizheh A., Bahrami A. R., Matin M. M., Edalatmanesh M. A., Zomorodipour A., Gardaneh M., Farshchian M. and Momeni-Moghaddam M. (2013) Human adipose-derived mesenchymal stem cells can survive and integrate into the adult rat eye following xenotransplantation. *Xenotransplantation* 20:165-176.

9- Hever A., Williamson K. and Van Heyningen V. (2006) Developmental malformations of the eye: the role of *PAX6*, *SOX2* and *OTX2*. *Clinical genetics* 69:459-470.

10- Huang L., Liang J., Geng Y., Tsang W. M., Yao X., Jhanji V., Zhang M., Cheung H. S., Pang C. P. and Yam G. H. (2013) Directing adult human periodontal ligament-derived stem cells to retinal fate. *Investigative Ophthalmology and Visual Science* 54:3965-3974.

11- Huang W., Fileta J., Guo Y. and Grosskreutz C. L. (2006) Downregulation of *Thy1* in retinal ganglion cells in experimental glaucoma. *Current eye research* 31:265-271.

12- Huang Y., Enzmann V. and Ildstad S. T. (2011) Stem cell-based therapeutic applications in retinal degenerative diseases. *Stem Cell Rev* 7:434-445.

13- Jagatha B., Divya M. S., Sanalkumar R., Indulekha C. L., Vidyanand S., Divya T. S., Das A. V. and James J. (2009) *In vitro* differentiation of retinal ganglion-like cells from embryonic stem cell derived neural progenitors. *Biochemical and biophysical research communications* 380:230-235.

14- Jayaram H., Becker S. and Limb G. A. (2011) *Stem Cell Based Therapies for Glaucoma*.

15- Jin Z. B., Okamoto S., Mandai M. and Takahashi M. (2009) Induced pluripotent stem cells for retinal degenerative diseases: a new perspective on the challenges. *J Genet* 88:417-424.

16- John S., Natarajan S., Parikumar P., Shanmugam P. M., Senthilkumar R., Green D. W. and Abraham S. J. (2013) Choice of Cell Source in Cell-Based Therapies for Retinal Damage due to Age-Related Macular Degeneration: A Review. *J Ophthalmol* 2013:465169.

17- Jurgens W. J., Oedayrajsingh-Varma M. J., Helder M. N., Zandiehoulabi B., Schouten T. E., Kuik D. J., Ritt M. J. and van Milligen F. J. (2008) Effect of tissue-harvesting site on yield of stem cells derived from adipose tissue: implications for cell-based therapies. *Cell and tissue research* 332:415-426.

18- Kayama M., Kurokawa M. S., Ueda Y., Ueno H., Kumagai Y., Chiba S., Takada E., Ueno S., Tadokoro M. and Suzuki N. (2010) Transfection with *PAX6* gene of mouse embryonic stem cells and subsequent cell cloning induced retinal neuron

progenitors, including retinal ganglion cell-like cells, *in vitro*. Ophthalmic research 43:79-91.

19- Kerrigan-Baumrind L. A., Quigley H. A., Pease M. E., Kerrigan D. F. and Mitchell R. S. (2000) Number of ganglion cells in glaucoma eyes compared with threshold visual field tests in the same persons. Investigative Ophthalmology & Visual Science 41:741-748.

20- Kokai L. E., Rubin J. P. and Marra K. G. (2005) The potential of adipose-derived adult stem cells as a source of neuronal progenitor cells. Plast Reconstr Surg 116:1453-1460.

21- Kolpak A., Zhang J. and Bao Z. Z. (2005) Sonic hedgehog has a dual effect on the growth of retinal ganglion axons depending on its concentration. The Journal of neuroscience 25:3432-3441.

22- Kuehn M. H., Fingert J. H. and Kwon Y. H. (2005) Retinal ganglion cell death in glaucoma: mechanisms and neuroprotective strategies. development 1:3.

23- MacLaren R. E., Pearson R. A., MacNeil A., Douglas R. H., Salt T. E., Akimoto M., Swaroop A., Sowden J. C. and Ali R. R. (2006) Retinal repair by transplantation of photoreceptor precursors. Nature 444:203-207.

24- Marquardt T. and Gruss P. (2002) Generating neuronal diversity in the retina: one for nearly all. Trends in neurosciences 25:32-38.

25- Moshiri A., Close J. and Reh T. A. (2004) Retinal stem cells and regeneration. Int J Dev Biol 48:1003-1014.

26- Osakada F., Jin Z. B., Hirami Y., Ikeda H., Danjyo T., Watanabe K., Sasai Y. and Takahashi M. (2009) *In vitro* differentiation of retinal cells from human pluripotent stem cells by small-molecule induction. Journal of cell science 122:3169-3179.

27- Pevny L. H., Sockanathan S., Placzek M. and Lovell-Badge R. (1998) A role for SOX1 in neural determination. Development 125:1967-1978.

28- Philips G. T., Stair C. N., Young Lee H., Wroblewski E., Berberoglu M. A., Brown N. L. and Mastick G. S. (2005) Precocious retinal neurons: *PAX6* controls timing of differentiation and determination of cell type. Developmental biology 279:308-321.

29- Pichaud F. and Desplan C. (2002) Pax genes and eye organogenesis. Curr Opin Genet Dev 12:430-434.

30- Quigley H. A. and Broman A. T. (2006) The number of people with glaucoma worldwide in 2010 and 2020. British Journal of Ophthalmology 90:262-267.

31- Ramsden C. M., Powner M. B., Carr A. J., Smart M. J., da Cruz L. and Coffey P. J. (2013)

Stem cells in retinal regeneration: past, present and future. Development 140:2576-2585.

32- Rasmussen J. G., Frobert O., Holst-Hansen C., Kastrup J., Baandrup U., Zachar V., Fink T. and Simonsen U. (2012) Comparison of human adipose-derived stem cells and bone marrow-derived stem cells in a myocardial infarction model. Cell Transplant.

33- Sapieha P. S., Peltier M., Rendahl K. G., Manning W. C. and Di Polo A. (2003) Fibroblast growth factor-2 gene delivery stimulates axon growth by adult retinal ganglion cells after acute optic nerve injury. Molecular and Cellular Neuroscience 24:656-672.

34- Schmittgen T. D. and Livak K. J. (2008) Analyzing real-time PCR data by the comparative C(T) method. Nat Protoc 3:1101-1108.

35- Shimizu N., Watanabe H., Kubota J., Wu J., Saito R., Yokoi T., Era T., Iwatsubo T., Watanabe T., Nishina S., Azuma N., Katada T. and Nishina H. (2009) *PAX6* -5a promotes neuronal differentiation of murine embryonic stem cells. Biol Pharm Bull 32:999-1003.

36- Singhal S., Bhatia B., Jayaram H., Becker S., Jones M. F., Cottrill P. B., Khaw P. T., Salt T. E. and Limb G. A. (2012) Human Müller Glia with Stem Cell Characteristics Differentiate into Retinal Ganglion Cell (RGC) Precursors *In vitro* and Partially Restore RGC Function *In vivo* Following Transplantation. Stem cells translational medicine 1:188-199.

37- Spence J. R., Madhavan M., Ewing J. D., Jones D. K., Lehman B. M. and Del Rio-Tsonis K. (2004) The hedgehog pathway is a modulator of retina regeneration. Development 131:4607-4621.

38- Wallace V. A. (2007) Stem cells: a source for neuron repair in retinal disease. Canadian Journal of Ophthalmology/Journal Canadien d'Ophthalmologie 42:442-446.

39- Wong I. Y. H., Poon M. W., Pang R. T. W., Lian Q. and Wong D. (2011) Promises of stem cell therapy for retinal degenerative diseases. Graefes Archive for Clinical and Experimental Ophthalmology 249:1439-1448.

40- Yasuda T., Kajimoto Y., Fujitani Y., Watada H., Yamamoto S., Watarai T., Umayahara Y., Matsuhisa M., Gorogawa S., Kuwayama Y., Tano Y., Yamasaki Y. and Hori M. (2002) *PAX6* mutation as a genetic factor common to aniridia and glucose intolerance. Diabetes 51:224-230.

41- Zhang X. M. and Yang X.-J. (2001) Regulation of retinal ganglion cell production by Sonic hedgehog. Development 128:943-957.