

Impact Factor:

ISRA (India) = 1.344	SIS (USA) = 0.912	ICV (Poland) = 6.630
ISI (Dubai, UAE) = 0.829	PIHHI (Russia) = 0.179	
GIF (Australia) = 0.356	ESJI (KZ) = 1.042	
JIF = 1.500	SJIF (Morocco) = 2.031	

SOI: [1.1/TAS](#) DOI: [10.15863/TAS](#)

International Scientific Journal Theoretical & Applied Science

p-ISSN: 2308-4944 (print) e-ISSN: 2409-0085 (online)

Year: 2015 Issue: 10 Volume: 30

Published: 30.10.2015 <http://T-Science.org>

Ravshon Khudaybergan-ugli Khudayberganov
PhD, Head of the «Social sciences» Department
The Tashkent Automobile and Road Institute,
Uzbekistan
paradigma777@bk.ru

SECTION 30. Philosophy.

SOCIAL-PHILOSOPHICAL ANALYSIS OF SUICIDAL DEATH AS ONE OF THE ASPECT OF THANATOLOGY

Abstract: This article is devoted to one of the most complex psychological issues associated with the philosophical teaching of thanatology. Here is socio-philosophical analysis of this problem.

Key words: suicide, death, suicidal tendencies, euthanasia, personality, statistics.

Language: English

Citation: Khudayberganov RK (2015) SOCIAL-PHILOSOPHICAL ANALYSIS OF SUICIDAL DEATH AS ONE OF THE ASPECT OF THANATOLOGY. ISJ Theoretical & Applied Science 10 (30): 124-126.

Soi: <http://s-o-i.org/1.1/TAS-10-30-26> **Doi:**  <http://dx.doi.org/10.15863/TAS.2015.10.30.26>

Deaths related to suicide, is one of the socio-economic and health problems are now very relevant in forensic science and practice. The study of suicides based on forensics in Uzbekistan only in episodic form of special interest to the results of which are specified in the present study, as well as the role and significance of the hazards that create a propensity for such actions.

In cases of suicide significance of the results of forensic examination of the corpse, with the successful conduct of the investigation, the nature of the existing injury resolved the problem of proportionality of their mechanism and details of the subject. A study of the materials forensics indicates the presence of a number of problems in the stages of use. Organization and conduct of examinations. In particular, the position of embracing the control of the forensic examination of all of these cases are in place, above all, makes it impossible for a legal assessment of the situation, and secondly, have a negative impact on the development of preventive measures aimed at obtaining objective information about them, planned the study, and prevention of suicide accidents. In is possible to emphasize the abstractness of tasks before the examination to the discrepancy in certain cases the details of the case. At the same time based on the position of the two cases must indicate the usefulness of the issues of under examination.

Suicidal state is an attack on his life, antihuman attitude. Therefore, Abramov A.G. confirms that suicidal case is arbitrary and mental action mentally healthy person to an attempt on his life. It can be

finished (suicide) and not finished (parasuicidal) [1], suicidal action, is attempt on his life made arbitrarily and deliberately. Comprehensive full inspection suicide cases, the study of the hazards that affect the formation of these cases are important in reducing the number of these cases, the development of effective preventive measures to prevent them.

The most alarming part related to increased destructiveness of man: the cases as unjustified aggression and necrophilia, sadism (violence), the mass of young people attempt on his life, terrorism, become common in developed countries. For example, as the R.L.Harpe "In the U.S. suicide cases take place in the fourth cause of death" [3]. Consideration of all these problems in an industrial civilization context helps to analyze important issues in modern society, such as terrorism, euthanasia, demonstrative death, collective ritual attempt on his life.

In order to study in our country reason of suicide attempts on his life, and death has been active National Center for Pathological Anatomy at the Ministry of Health, as well as centers of forensic examination. They also include the definition of the epidemiology of suicide cases and pending in Uzbekistan, hazards that are important in their genesis, territorial characteristics, organization of services in these cases, forensic, identification of gaps and development activities due to these for their further improvement. In this case, it would be useful definition of assassination attempts on his life, as well as suicides and many similar cases that are contrary to human life and in Uzbekistan and doing



Impact Factor:

ISRA (India) = 1.344	SIS (USA) = 0.912	ICV (Poland) = 6.630
ISI (Dubai, UAE) = 0.829	PIHHI (Russia) = 0.179	
GIF (Australia) = 0.356	ESJI (KZ) = 1.042	
JIF = 1.500	SJIF (Morocco) = 2.031	

various measures against them. Below a classification of death by suicide: 1-mechanical aspics, resulting of hanging, 2-Acute poisoning, 3-immolation cases, 4-samosbrasyvaniya with height, 5-rail injury bloodletting, 6-bow [5]. Emphasizes the importance of forensic medical examination to determine the circumstances of the

reasons of murder and suicides committed intentionally and suddenly, on the assumption that other types of homicide are not included in the materials forensics.

It should be emphasized reduction of suicidal death in Uzbekistan from year to year (Table 1).

Table 1

The results of cases of suicidal deaths in 2000-2008 in the Republic of Uzbekistan.

Place	2000	2001	2002	2003	2004	2005	2006	2007	2008
Andijan	134	141	126	93	80	92	120	129	116
Bukhara	52	27	19	21	24	22	15	26	34
Jizzah	45	63	46	26	34	22	24	32	36
Navoi	89	94	65	49	45	65	47	43	50
Namangan	125	121	148	174	100	116	141	154	115
Samarkand	172	163	138	116	113	113	107	91	96
Sirdarya	68	89	60	49	48	50	48	45	59
Surkhandarya	75	40	31	30	36	21	38	28	30
Tashkent	313	308	300	248	222	229	231	188	184
Tashkent region	540	527	420	365	379	377	379	361	361
Fergana	243	235	231	197	225	199	205	192	218
Khorezm	80	91	75	57	44	53	59	49	31
Kashkadarya	73	58	63	48	68	27	43	45	36
Karakalpakstan	71	41	23	20	29	24	25	37	31
Total	2080	1998	1745	1493	1447	1410	1482	1420	1397

Examination analysis conducted on suicide cases of suicide in the office OPLU forensics in 2000-2008 (Figure 1).

For example, consider Andijan region: in 2000 the total mortality rate reached 6.4% -134, in 2008,

this case fell to 5.5% 116. Nablyudalis suicide suicides in Samarkand in 2000, 8.2% -172, in 2008 4.6% 96, in the Tashkent region in 2000, 26% -540, in 2008, 17.3% -361, in the Republic of Karakalpakstan in 2000, 9,46-71, in 2008, 1,4% -31.




Figure 1 -The analysis of examination spent in suicidal cases on sex in a forensic medical examination bureau in 2000-2008.

There are at least two forms of suicidal attempts on his life. First of them suicide-sensitive, meaning an irrational suicide person in all forms. It should be

stressed that the views of society, "Hemlock" about this form of suicide reflect the views of the American Association of Suicidology, as well as all other

Impact Factor:

ISRA (India) = 1.344	SIS (USA) = 0.912	ICV (Poland) = 6.630
ISI (Dubai, UAE) = 0.829	PIHHI (Russia) = 0.179	
GIF (Australia) = 0.356	ESJI (KZ) = 1.042	
JIF = 1.500	SJIF (Morocco) = 2.031	

communities and it makes clear the need to prevent all cases of this suicide. One not endorse any form of attempts on his life, saying psychological illness or the difficulties of life.

However, in our opinion, the second rate of suicide can be justified to call suicide. On implies deliverance through rationally pre-planned attempt on his life. In other words, it is euthanasia using suicide as a means.

In the analysis of suicide in the spiritual and moral sense to take into account the following reasons—firstly, an incurable disease, intolerable suffering which allows this man. Every man is freed from all suffering through the attempt on his life. Secondly, the state of serious physical injury that staff, after some thought could not take his using. This attempt on his life occurs seldom. In history that always can be an example of a lawsuit brought by Mrs. Elizabeth Bovina. By this regard, it is worth noting and following. Firstly, to obtain the first report of the disease, which causes a danger of human life should not try to get rid of the disease by the attempt on his life. It is necessary to provide special medical helpness.

Receiving notification of a serious illness with a fatal outcome can not approve the decision of the attempt on his life. Secondly, this report should be your doctor, as well as to take into account its opinion. Doctor's opinion will depend on cases. No despite this, the physician should information of this decision. Because in a state of severe emotional and physical pain he could not hear or understand the diagnosis. Thirdly, a man encroaching on their lives, care should be taken so as not to attract other criminal answer. Haw mentioned above, assistance in committing suicide a criminal answer. Fourth, the

person must leave a letter, stating openly about the attempt on his life. If implementation attempts on his life is in a hotel, hospital or other public institution, we must not forget to leave a letter requesting apology from management of the institution for the inconvenience as assistance in the implementation of suicide may be the cause of criminal answer. Some people do not want to endanger the family and prefer to implement suicide elsewhere.

Should a person instead of the attempt on his life to go in hospice? Briefly, the hospice is very good at the hardest of all works. They do their job with great skill and love. Action of euthanasia support them. No one want to push all his death as helpness. Without that all of the Hospice of the forces they can not give an opportunity to stop the approaching death. In is necessary to recognize the fact that most good hospices will organize the most convenient conditions and care at a high level for each man. How recent surveys show that many hospitals use the standards for hospice itself. This proves that hospices are very well established educational works [6]. However, in our opinion wrong to think that the interests of supporters of the hospice euthanasia and intersect at one point. All of them is designed for people of different categories that have completely different values.

Thus, analyzing the materials training and medical examination, conducted in our country in 2000-2008, can clearly see the reduction of suicides by 33%. This suggests that human life is the highest value and, most importantly, people of various people have begun to understand. It means that the moral and spiritual position of our people have changed attitude to life and it is possible to reasonably get out of any problem situations.

References:

1. Ambrumova AG (1981) Principle of the organization suicidal services and prospects of scientific researches. // Works of the Moscow scientific research institute of psychiatry. Volume № 92. 1981. – pp. 13-16.
2. Ambrumova AG, Postovalova LI (1983) Analyses of axonal notes of suicide. FF Scientific and organizational problems suicidal: Research. - M: 1983. – pp. 53-74.
3. R. La Harpe (1995) Suicide in the Geneva canton (1971-1990). An analysis of the forensic medicine autopsy sample. // Arch Criminal. 1995. Mar-Apr. 195. (3-4). 65-74.
4. Giyozov ZA, Mahsumhonov KA (2006) A suicide court-tibby questions. - T: "Uzbekistan Scientifics print", 2006. – pp.25.
5. Giyozov ZA, Mahsumhonov KA (2006) A suicide court-tibby questions. - T: "Uzbekiston Scientific print", 2006. – pp.29.
6. Robert El. Kane, et al. (1984) A. Randomized Controlled Thai of Hospice (The lancer 1984. N 8382. April 21). pp. 891-893.
7. Mechnikov II (1988) Etyudy optimizma . - Izd . -ye . - M .: Nauka , 1988. - 328 s . (1 - ye izdaniye - 1907 ; 2 - ye - 1964).
8. Shor GV (2002) O smerti cheloveka (vvedeniye v tanatologiyu) . - SPb .: Izd - vo SPbGMU , 2002. - 272 p.
9. Demichev AV (1997) Diskursy smerti : Vvedeniye v filosofskuyu tanatologiyu . - Spb .: INAPRESS , 1997. - 144 p.
10. Ryazantsev S (1994) Tanatologiya (ucheniye o smerti) / Vostochno - Yevropeyskiy institut psikoanaliza .. - Spb .: VEIP , 1994. - 380 p.

