



Contents lists available at ScienceDirect

Asian Pacific Journal of Tropical Disease

journal homepage: www.elsevier.com/locate/apjtd

Document heading doi:10.1016/S2222-1808(11)60026-1

Prevalence of hypoglycemia among severe malaria children in a rural African population

Osonuga OA^{1,2}, Osonuga AA³, Osonuga IO⁴, Osonuga A^{2*}, Derkyi Kwarteng L²¹Department of Pharmacology, Olabisi Onabanjo University, Remo Campus, Nigeria²School of Medical Sciences, University of Cape-coast, Ghana³Department of Nursing, University of Cape-coast, Ghana⁴Department of Physiology, Olabisi Onabanjo University, Remo Campus, Nigeria

ARTICLE INFO

Article history:

Received 15 May 2011

Received in revised form 10 June 2011

Accepted 10 July 2011

Available online 28 September 2011

Keywords:

Hypoglycemia
Severe malaria
Prevalence
African
Glucose

ABSTRACT

Objective: To determine the prevalence of hypoglycemia in severe malaria children in a rural African community. **Methods:** Thirty two children who fulfilled the inclusion criteria were recruited for the study from two hospitals with intensive care facilities. Blood sugar levels of the patients were monitored serially at admission and then every 4 hours (after admission) for 24 hours using a glucometer; taking values <2.2 mmol/L as hypoglycemia. The children received intensive care according to WHO guidelines for severe malaria. **Results:** Fifteen out of 32 children recruited (representing 46.9%) had hypoglycemia with about 60% under 5 years of age. The mean age of the children was (4.49±2.89) years. The pre-correction and average post correction blood glucose levels were (2.04±0.13) mmol/L and (3.18±0.23) mmol/L, respectively. **Conclusions:** The result has demonstrated that about 1 in 2 children with severe malaria may suffer from hypoglycemia in an African rural environment hence there is a need for at least three-point glucose estimation (at recruitment, 4 hours following correction and the end of the 24 hours).

1. Introduction

Malaria continues to be a major global health problem, with over 40% of the world's population (more than 2 400 million people) exposed to varying degrees of malaria risk in some 100 countries in Africa, Asia, central America, Oceania and south America.

It causes more than one million deaths worldwide each year, and over 90% of them occur in Africa. Severe malaria is caused by *Plasmodium falciparum* infection and usually occurs as a result of delay in treating an uncomplicated attack of falciparum malaria. In children, it could however develop rapidly^[1–4].

Severe malaria is approximately 1%–2% of clinical attacks of malaria in African children. About 1–2 million deaths occur mostly in young children (under 5 years) with a child dying every 30 seconds. Hence it is a medical emergency to require specialist care^[4,5].

Hypoglycemia (blood glucose <2.2 mmol/L) has been

identified as an independent risk of death in children presenting with severe malaria along with coma, repeated convulsions, shock, and hyperparasitemia. In Africa, 153 000–267 000 malaria related deaths are attributed to hypoglycemia. Generally, 8% of adults, 30% of children and 50% of pregnant women in late pregnancy develop hypoglycemia. Sudden unexplained deterioration in a patient with severe falciparum malaria is commonly due to hypoglycemia. It is easily overlooked clinically because the manifestations may be similar to those of cerebral malaria^[6–8].

Since studies in African children with uncomplicated malaria cannot be extrapolated to children with severe disease hence^[9], this study seeks to find out the prevalence of hypoglycemia in severe malaria cases among Nigerian children on the day of hospital admission.

2. Materials and methods

2.1. Patients

A total of 32 patients were recruited from Overcomers Specialist Hospital, Ilishan and General Hospital Ikenne, Ikenne local government area of Ogun State, Nigeria. The

*Corresponding author: Osonuga Ayokunle, School of Medical Sciences, University of Cape coast, Ghana.

E-mail: overcomers2007@yahoo.com

hospitals have facilities for resuscitating and handling emergency.

Patients who satisfied inclusion criteria (see below) were admitted for estimation of blood glucose and treatment in the ward. On enrolment, a brief history was obtained from accompanying adult (which may be the parent or guardian) and clinical examination was performed.

Ethical and parental approval for the study was obtained from Olabisi Onabanjo University Teaching Hospital, Nigeria, Joint Ethical Review Committee and informed consent from the parents or guardians.

Inclusion criteria: (1) Children from either sex with age ranging from 1 year to 12 years; (2) Fever with temperature higher than 37.5 °C within the last 24 hours; (3) Presence of convulsion, vomiting, hypoglycemia, anemia and headache; (4) Informed consent obtained from the parents and guardians; (5) Assurance that patients will be resident within catchments of study for follow-up; (6) Absence of concomitant illness such as bronchopneumonia, typhoid, meningitis, urinary tract infection; (7) Absent history of administration of antipyrexia; (8) The Blantyre coma score of <3.

Exclusion criteria: (1) History of blood transfusion in the last two months; (2) Presence of concomitant illness; (3) History of previous allergy to quinine and artemether; (4) Lack of informed consent.

Withdrawal criteria: (1) If any concomitant illness developed during the study; (2) If informed consent is withdrawn by parents or guardian; (3) If patient (or parents/guardian) is unwilling to continue in the study; (4) Failure to comply with protocol.

2.2. Blood sugar monitoring

Blood sugar level was monitored upon arrival of the patient by using a digital glucometer. Hypoglycemia was defined as blood glucose level < 2.2 mmol/L. Prompt correction of hypoglycemia (in children who were hypoglycemic) was done by infusion of 50% dextrose at a dose of 1.0 mL/kg b.w. to the child, diluted in an approximately equal volume of normal saline infusion fluid over a period of about 5 minutes. This was followed with a continuous intravenous infusion of 5% glucose fluid. Serial monitoring was done every 4 hours on the day of admission. The blood glucose level upon

admission and the average values (post correction) over a period of 24 hours were recorded.

Standard treatment protocols for treating severe malaria children were employed for the treatment of other symptoms.

2.3. Statistical analysis

All calculations were done using the SPSS-V15 statistical software package for analysis of the data. The data were presented as Mean±SD, and statistical analysis was carried out using the student's paired *t*-test and ANOVA. Differences were considered to be statistically significant at an error probability of less than 0.05 ($P \leq 0.05$).

3. Results

Fifteen out of 32 children recruited (representing 46.9%) had hypoglycemia with about 60% were under 5 years of age. Table 1 showed the pattern of their blood sugar levels at enrollment and the average values obtained for blood glucose after correction with glucose infusion. The mean age of the children was (4.49±2.89) years. The pre-correction and average post correction blood glucose levels were (2.04±0.13) mmol/L and (3.18±0.23) mmol/L, respectively. *P* value less than 0.05 was significant.

4. Discussion

Effort in this study was to evaluate the prevalence of hypoglycemia in children presenting with severe malaria on hospital admission.

The mechanism of hypoglycemia in severe malaria is unknown. Impaired hepatic gluconeogenesis (even in the presence of needed substrate *e.g.* glycerol and alanine), increasing metabolic demands of the host and the parasite are thought to be probable causes^[6,10]. In our study, 46.9% of the children were hypoglycemic. This supports work done previously by other researchers who identified hypoglycemia as one of the common features of severe malaria in children (independent of quinine induced hypoglycemia)^[11,12]. However, an Indian study concluded that hypoglycemia is observed in up to 30% of cases^[13]. This disparity may be

Table 1
Pattern of hypoglycemia among severe malaria children at recruitment and after correction.

Age (months/years)	Blood sugar level (at enrolment)*	Average blood sugar level (post-correction)*
12.0**	1.8	3.00
13.0**	2.0	3.00
15.0**	2.1	2.80
2.0	1.8	3.00
2.5	2.0	3.20
3.0	2.0	3.00
3.5	2.0	3.10
4.0	2.1	3.30
4.5	2.2	3.10
5.0	2.0	3.20
6.0	2.2	3.06
6.5	2.2	3.50
8.0	2.0	3.60
9.0	2.2	3.50
10.0	2.0	3.40

*: Blood glucose < 2.2 mmol/L was considered as hypoglycemia. **: Age in months.

attributed to late presentation of patients to the hospital in most rural Nigerian communities as well as few referral hospitals in the rural areas that can handle such pediatric emergency. Also, there is usually a history of prolonged vomiting, anorexia and forceful feeding with native medication (rampant among rural dwellers). Other newer studies have also reported associated risk of hypoglycemia in severe malaria to starvation^[14,15].

Hypoglycemia being associated with higher risk of death as well as being an inclusive criteria for severe malaria, deserves close monitoring unlike other parameters commonly monitored in severe malaria treatment such as parasitemia clearance, temperature and PCV which are usually done on day 0, 3 and 14^[16].

Age also had an impact on hypoglycemia presentation in the children with severe malaria studied. 60% of the children with hypoglycemia were under 5 years. This is in agreement with the findings of other researchers that indicated that hypoglycemia is more common in younger children than older ones^[15,17–21].

The burden of hypoglycemia is multifactorial as it can worsen convulsion and may deepen coma in children with cerebral malaria making recovery and prognosis very poor^[22–27]. The hypoglycemic children were offered prompt glucose correction. No patient had hypoglycemia after 24 hours of correction.

The result has demonstrated that about 1 in 2 children with severe malaria may suffer from hypoglycemia in an African rural environment where there is dearth of health care practitioners with specialist training, paucity of investigation and facilities, poor environmental sanitation, lack of door and window netting. All these factors contribute to high severe malaria burden. Since hypoglycemia has been identified as a common feature of severe malaria, there is a special need for serial monitoring and management of blood glucose. The study has demonstrated the need for at least three-point glucose estimation (at recruitment, 4 hours following correction and the end of the 24 hours) in such communities.

Conflict of interest statement

We declare that we have no conflict of interest.

References

- [1] WHO. *Malaria case management operations manual*. Geneva: WHO; 2009. [Online] Available from: whqlibdoc.who.int/publications/2009/9789241598088_eng.pdf. [Accessed on 20 July, 2010]
- [2] WHO. *Roll back malaria partners set ambitious financial targets for 19 African countries fighting malaria*. Geneva: WHO; 2007. [Online] Available from: http://www.rollbackmalaria.org/amd2007/pr/pr_rbmAMD2007-e.pdf. [Accessed on 17 Feb, 2010]
- [3] WHO. *Guidelines for the treatment of malaria*. 2nd ed. Geneva: WHO; 2010. [Online] Available from: <http://www.who.int/malaria/docs/TreatmentGuidelines2010.pdf>. [Accessed on 7 Aug, 2010]
- [4] WHO. *Management of severe malaria: a practical handbook*. 2nd ed. Geneva: WHO; 2000.
- [5] Osonuga O, Osonuga AA, Osonuga IO, Osonuga A. Temperature changes in the course of treatment of severe malaria patients with quinine and artemether. *Asian J Med Sci* 2011; **2**: 41–45.
- [6] Ashley EA, Nicholas J. White: malaria diagnosis and treatment. In: Jong EC, Sanford C. (eds.) *The travel and tropical medicine manual*. 4th ed. Philadelphia: WB Saunders/Elsevier; 2008, p. 308–326.
- [7] WHO. *Malaria case management operations manual*. Geneva: WHO; 2009. [Online] Available from: whqlibdoc.who.int/publications/2009/9789241598088_eng.pdf. [Accessed on 12 Dec, 2010]
- [8] Breman JG, Mills A, Snow RW, Mulligan JA, Lengeler C, Mendis K, et al. *Disease control priorities in developing countries: conquering malaria*. 2nd ed. Washington DC: World Bank; 2006, p. 413–431.
- [9] Agbenyega T, Angus BJ, Bedu-Addo G, Baffoe-Bonnie B, Guyton T, Stacpoole PW, et al. Glucose and lactate kinetics in children with severe malaria. *J Clin Endocrinol Metab* 2000; **85**: 1569–1576.
- [10] English M, Wale S, Binns G, Mwangi I, Sauerwein H, Marsh K. Hypoglycemia on and after admission in Kenyan children with severe malaria. *QJM* 1998; **91**: 191–197.
- [11] Severe falciparum malaria. World Health Organization, Communicable Disease Cluster. *Trans R Soc Trop Med Hyg* 2000; **94**(Suppl 1): s1–s90.
- [12] Maitland K, Nadel S, Pollard AJ, Williams TN, Newton CR, Levin M. Management of severe malaria in children: proposed guidelines for the United Kingdom. *BMJ* 2005; **331**(7512): 337–343.
- [13] Gupta D, Chugh K, Sachdev A, Soni A. ICU management of severe malaria. *Indian J Pediatr* 2001; **68**(11): 1057–1061.
- [14] Thien HV, Kager PA. Hypoglycemia in *Falciparum malaria*: is fasting an unrecognized and insufficiency emphasized risk factor? *Trends Parasitol* 2006; **22**(9): 410–415.
- [15] Zijlmans W, van Kempen A, Ackermans M, de Metz J, Kager P, Sauerwein H. Glucose kinetics during fasting in young children with severe and non-severe malaria in Suriname. *Am J Trop Med Hyg* 2008; **79**: 605–612.
- [16] Aguwa CN, Ukwe CV, Adibe MO. A comparative study of quinine and artemether in the treatment of severe malaria in Nigerian children. *Trop J Pharm Res* 2010; **9**(1): 11–17.
- [17] Mockenhaupt FP, Ehrhardt S, Burkhardt J, Bosomtwe SY, Laryea S, Anemana SD, et al. Manifestation and outcome of severe malaria in children in northern Ghana. *Am J Trop Med Hyg* 2004; **71**(2): 167–172.
- [18] Zijlmans WC, van Kempen AA, Serlie MJ, Sauerwein HP. Glucose metabolism in children: influence of age fasting and infectious diseases. *Metabolism* 2009; **58**(9): 1356–1365.
- [19] Khan Haris M, Fatima Shujatullah, Mohammad Ashfaq, Adil Raza. Changing trends in prevalence of different *Plasmodium* species with dominance of *Plasmodium falciparum* malaria infection in Aligarh (India). *Asian Pac J Trop Med* 2011; **4**(1): 64–66.
- [20] Ekanem EI, Agan TU, Efiok EE, Ekott MI, Okodi E. A study of anemia in women with asymptomatic malaria parasitaemia at their first antenatal care visit at the General Hospital, Ikot Ekpene, Akwa Ibom State, Nigeria. *Asian Pac J Trop Med* 2010; **3**(7): 567–570.
- [21] Afonso de Almeida, Virgílio E do Rosário, Ana Paula Arez, Pedro Cravo. Malaria epidemiology in the Democratic Republic of East Timor. *Asian Pac J Trop Med* 2011; **3**(4): 283–287.
- [22] Ogetii GN, Akech S, Jemutai J, Boga M, Kivaya E, Fegan G, et al. Hypoglycaemia in severe malaria, clinical associations and relationship to quinine dosage. *BMC Infect Dis* 2010; **10**: 334.
- [23] Lorenz V, Karanis P. Malaria vaccines: looking back and lessons learnt. *Asian Pac J Trop Biomed* 2011; **1**(1): 74–78.
- [24] Krungkrai SR, Krungkrai J. Malaria parasite carbonic anhydrase: inhibition of aromatic/heterocyclic sulfonamides and its therapeutic potential. *Asian Pac J Trop Biomed* 2011; **1**(3): 233–242.
- [25] Zerihun T, Degarege A, Erko B. Association of ABO blood group and *Plasmodium falciparum* malaria in Dore Bafeno Area, Southern Ethiopia. *Asian Pac J Trop Biomed* 2011; **1**(4): 289–294.
- [26] Wiwanitkit V. Concurrent malaria and dengue infection: a brief summary and comment. *Asian Pac J Trop Biomed* 2011; **1**(4): 326–327.
- [27] Dondorp AM, Lee SJ, Faiz MA, Mishra S, Price R, Tjitra E, et al. The relationship between age and the manifestations of and mortality associated with severe malaria. *Clin Infect Dis* 2008; **47**: 151–157.