



Contents lists available at ScienceDirect

Asian Pacific Journal of Tropical Disease

journal homepage: www.elsevier.com/locate/apjtd



Floral research

doi: 10.1016/S2222-1808(15)61051-9

©2016 by the Asian Pacific Journal of Tropical Disease. All rights reserved.

Determination of the total phenols, flavonoids and antimicrobial activity of the crude extracts from locally grown neem stems

Hoda Salim Khamis Al-Jadidi, Mohammad Amzad Hossain*

School of Pharmacy, College of Pharmacy and Nursing, University of Nizwa, P. O. Box 33, Postal Code 616, Nizwa, Sultanate of Oman

ARTICLE INFO

Article history:

Received 8 Oct 2014

Received in revised form 21 Oct 2014,

2nd revised form 27 Feb 2016

Accepted 29 Apr 2016

Available online 3 May 2016

Keywords:

Azadirachta indica

Neem

Total phenols

Total flavonoids

Antimicrobial activity

UV-visible spectroscopy

ABSTRACT

Objective: To prepare the crude extracts from the stems of neem with maceration extractor and determine their total phenolics, flavonoids contents and antimicrobial activity by established methods.

Methods: The dry stem samples were used to prepare different crude extracts using maceration method. The total phenolics and total flavonoids contents were determined through Folin-Ciocalteu's reagent and $AlCl_3$ methods. The antimicrobial activity of the crude extracts from the local neem was determined by disc diffusion method against one Gram-positive *Staphylococcus aureus* and three Grams-negative *Escherichia coli*, *Pseudomonas aeruginosa* and *vulgaris* pathogenic bacterial strains. The standard amoxicillin was used as a positive control for the antimicrobial study.

Results: The results of the total phenolics content of the crude extracts from different stems ranged from 20.80 mg/100 g to 107.29 mg/100 g of the powder crude extracts. The total flavonoids content of different crude extracts from powder stems of neem ranged from 136.50 mg/100 g to 484.50 mg/100 g of the powder samples. The antimicrobial activity of the crude extracts from different stems of neem at different concentrations against four bacterial strains didn't give any activity.

Conclusions: Thanks to the contents of the total phenolics and flavonoids, the crude extracts could be used as medicines for different ailments.

1. Introduction

Azadirachta indica A. Juss (*A. indica*) is a kind of tree belonging to Meliaceae family. It comes from Pakistan, India and Thailand[1]. Its English name is neem. Locally it is called "alshurisha". Neem tree grows very fast. The branches are wide and spreading. Leaves are mixed of both young and mature. The mature leaves are bright green and the young leaves are reddish to purplish color. Typical leaves consist of three main parts: leaf base, petiole and lamina. Leaves are attached to the stem by their leaf bases and may bear two lateral small leaves like structures called stipules[2].

Neem is used as traditional medicine in Indian culture for the treatment of different ailments. Neem is one of the main sources of many therapeutic agents[3]. In Thai traditional medicine system,

its leaves and flowers are used as a tonic for the treatment of fever[4]. The leaf and seed oils are very important medicines used for the treatment of symptoms of psoriasis and relieves itching[5]. Similarly, the fruits are used as an anthelmintic for the treatment of hemorrhoids and abnormal urination[6]. Traditionally, this tree was preferred over others because it produces fewer allergies and side effects[7].

Medicinally it is used widely as alternative therapeutic tools for the prevention or treatment of many diseases[8]. Each part of neem tree has several medicinal values to treat a wide range of human disorders such as antiseptic, diuretic, cough, nausea, vomiting, fever and peptic ulcer[9]. The most common use of this plant's juice is for gastrointestinal diseases[9]. Traditionally, Indian people used the leaves for the treatment of chicken pox sleep[10].

There is much extensive work by scientists on the analysis of chemical constitutions of the crude extracts neem. Many bioactive components have been isolated and identified from neem crude extracts such as azadirachtin, salannin, meliantriol and nimbin[11]. Among them the most active ingredient is reported as azadirachtin[11]. These chemical constitutions belong to the classes: beta-sitosterol, stigmasterol and limonoids[11,12]. Also, the other tricyclic and tetracyclic compounds have been isolated from this plant such as sulphides, flavonol glycosides, nimatol, quercetin,

*Corresponding author: Mohammad Amzad Hossain, School of Pharmacy, College of Pharmacy and Nursing, University of Nizwa, P. O. Box 33, Postal Code 616, Nizwa, Sultanate of Oman.

Tel: +968 99708496

Fax: +968 25446236

E-mail: hossainabi@gmail.com

Foundation Project: Supported by University of Nizwa, Nizwa, Sultanate of Oman (Grant No. SOP/ADO/72/2014).

The journal implements double-blind peer review practiced by specially invited international editorial board members.

myricetin and kaempferol[12].

Traditionally, Omanis use the whole plant for the treatment of fever, to kill worm and so on[12]. The literature search reveals that no scientific data on the total phenolics, flavonoids and antimicrobial activity are available in Oman. Therefore, the main objectives of this present study is to prepare crude extracts using different polarities of solvents by maceration method and determine their total phenolics, flavonoids and antimicrobial activity from the crude extracts of neem stems collected from Muscat.

2. Materials and methods

2.1. Materials and chemicals

The solvents acetone, butanol, chloroform, ethyl acetate, methanol, ethanol, hexane dichloromethane and dimethyl sulphoxide (DMSO) were collected from Sigma-Aldrich Company, Germany. The standard gallic acid and quercetin for the determination of the total phenolics and flavonoids were collected from Sigma Company Ltd, Germany. Sodium hydroxide and aluminum chloride were obtained from Philip Harris, England. Folin-Ciocalteu's reagent (FCR) and sodium nitrate for the total phenolics were obtained from Scharlau, Spain. UV-visible spectrophotometer (UV-visible) was from Jasco Company, Japan.

2.2. Bacterial strains

The employed bacterial strains against different crude extracts of neem such as *Staphylococcus aureus* (*S. aureus*), *Escherichia coli* (*E. coli*), *Pseudomonas aeruginosa* (*P. aeruginosa*) and *Proteus vulgaris* (*P. vulgaris*) were collected from the Department of Biological Sciences, College of Arts and Sciences, University of Nizwa, Sultanate of Oman.

2.3. Plant sample

The stem samples were collected from Al-Amerat, Muscat, Sultanate of Oman. The samples were harvested on February 8, 2014 at 3 pm. The collected samples were transported to the Natural Product Lab, School of Pharmacy, College of Pharmacy and Nursing, University of Nizwa, Sultanate of Oman. The plant was identified by botanists and the specimen number (006) was deposited in the lab.

2.4. Preparations of samples

The stem samples were washed to remove the unwanted substances. Then, the samples were cut into small pieces and dried at room temperature for three days. The dried stem samples were powdered by a grinding machine.

2.5. Extraction procedure

The stem powder (150 g) were taken in a beaker and 250 mL of methanol was added. The mixture was kept for 3 days. The sample was filtered by using a Bruckner funnel. After filtration, the filtrate was taken in a round bottom flask and evaporated by using a rotary evaporator. After evaporation, the crude extract was dissolved in water (150 mL) and shaken until it was dissolved. The dissolved mixture was transferred into a separatory funnel and solvents with

different polarities were added twice with increasing polarities such as hexane, chloroform, ethyl acetate and butanol (30 mL, and 20 mL). After the extraction procedure was finished, the solvent was evaporated to get crude extracts with different polarities.

2.6. Determination of the total phenolics content

2.6.1. Preparation of reagents

FCR (10 mL) was taken in a 100-milliliter volumetric flask with 90 mL of water. Three grams of sodium carbonate was taken in a 50-milliliter volumetric flask with 50 mL of water.

2.6.2. Procedure

The prepared crude extracts from the stems of neem (4 mg) were taken in separate test tubes and 4 mL of methanol was added. A total of 200 μ L of solutions from each crude extract was taken in a test tube and 1.5 mL of FCR was added and then kept in dark place for 5 min. After dark, 1.5 mL of sodium carbonate was added to the same solution and mixed by hand. All the tubes were kept at dark place for 2 h. Finally, the absorbance was recorded by using UV-visible at the wave length of 760 nm[13].

2.7. Determination of the total flavonoids content

The prepared crude extracts (4 mg) were taken in separate test tubes and 4 mL of methanol was added. Then, 0.25 mL of each crude solution was taken in a test tube and 1.25 mL of water and 75 μ L of sodium nitrate were added. The whole mixture was kept for 6 min and 150 μ L of aluminum chloride was added to react for 5 min in the dark place. Finally, 0.5 mL of sodium hydroxide and 0.275 mL of water were added to it. The absorbance was measured by UV-visible at the wave length of 450 nm[14].

2.8. Antimicrobial activity test

Each crude extracts from the stems of neem (4 mg) was taken in a volumetric flask and 4 mL of DMSO was added[15]. Different concentrations such as 2.00 mg/mL, 1.00 mg/mL, 0.50 mg/mL and 0.25 mg/mL were prepared by serial dilution technique. Amoxicillin (3 mg) was used as a standard by adding 3 mL of DMSO. Each prepared concentrations of each crude extracts was tested for its antimicrobial activity against one Gram-positive bacteria *S. aureus* and three Gram-negative bacteria *E. coli*, *P. aeruginosa* and *P. vulgaris* on MacConkey agar plate using disc diffusion method. Whatmann filter paper No. 41 sterile filter paper discs with a diameter of 6 mm were impregnated with each crude extract and placed on the inoculated agar. The concentration of amoxicillin standard used for this study was 1 mg/mL. All the plates were incubated at 37 °C for 24 h. The antibacterial activity was evaluated by measuring the diameter of the inhibition zones against the tested bacteria. Each method in this experiment was replicated three times.

3. Results

The powdered stem samples were used for the extraction with methanol by maceration method for 3 days. After completing extraction, methanol was evaporated by a rotary evaporator and defatted with water. Then, different solvents such as hexane, chloroform, ethyl acetate and butanol were used for extraction with increasing polarities.

3.1. Total phenolics content

All crude extracts such as hexane, ethyl acetate, chloroform, butanol, methanol and water crude extracts from the stems of neem were used for the determination of the total phenol contents by using FCR[13]. The result showed that the highest amount of the total phenols in the stem crude extracts was in ethyl acetate and the lowest was in hexane. The order was ethyl acetate > chloroform > butanol > methanol > hexane. The gallic acid was used as a standard calibration curve (Figure 1).




Figure 1. Gallic acid standard calibration curve.

3.2. Total flavonoids contents

The total flavonoid content in all neem crude extracts of stems was determined through aluminum chloride method based on the quercetin standard curve (Figure 2)[14]. The highest amount of the total flavonoids was in hexane (484.50 mg/100 g dry powdered samples) and the lowest was in butanol (136.50 mg/100 g dry powdered samples) and followed by ethyl acetate (339.50 mg/100 g), methanol (314.00 mg/100 g), chloroform (151.00 mg/100 g) and butanol (136.50 mg/100 g) among the five stems crude extracts.




Figure 2. Quercetin standard calibration curve.

3.3. Antimicrobial activity

The antimicrobial activity was determined by the disc diffusion method against four different bacterial strains[15]. Six crude extracts of stems at different concentrations were used to determine the antimicrobial activity through disc diffusion method. A standard amoxicillin antibiotic was used in this study. The result obtained from this study was presented in Table 1. Almost all the crude extracts from the stems of neem did not show any potential activity against the employed Gram-positive and Gram-negative bacterial strains. All the crude extracts did not show activity against *E. coli* except for ethyl acetate. The ethyl acetate crude extract at concentrations of 2.00 and 1.00 mg/mL showed small activity

against *E. coli* bacterial strain. On the other hand, almost all crude extracts did not show any activity against *S. aureus* at all concentrations except ethyl acetate (2.00 mg/mL) and water (1.00 mg/mL). The small activity was obtained from both crude extracts of ethyl acetate and water at the concentrations of 2.00 mg/mL and 1.00 mg/mL with the values of 7 and 4 mm. Similarly, *P. aeruginosa* bacterial strain showed some activity at different concentrations of the employed crude extracts from the stems of neem. Both hexane and chloroform crude extracts at concentration of 0.25 mg/mL gave the activity with the value of 6 mm. But butanol, methanol and water crude extracts of stems at concentration of 2.00 mg/mL gave the range of activity with the value of 7 mm (Table 1). However, all crude extracts at all concentrations did not give any activity against the *P. vulgaris* bacterial strain.

Table 1

Antimicrobial activities of different crude extracts form stems of neem. mm.

| Extracts | Concentrations (mg/mL) | <i>E. coli</i> | <i>S. aureus</i> | <i>P. aeruginosa</i> | <i>P. vulgaris</i> |
|---------------|------------------------|----------------|------------------|----------------------|--------------------|
| Hexane | 2.00 | nd | nd | nd | nd |
| | 1.00 | nd | nd | nd | nd |
| | 0.50 | nd | nd | nd | nd |
| | 0.25 | nd | nd | 6.00 ± 0.55 | nd |
| | Standard | 9.00 ± 0.13 | 9.0 ± 0.29 | nd | nd |
| Ethyl acetate | 2.00 | 7.00 ± 0.21 | 7.00 ± 0.37 | nd | nd |
| | 1.00 | 6.00 ± 0.43 | nd | nd | nd |
| | 0.50 | nd | nd | nd | nd |
| | 0.25 | nd | nd | nd | nd |
| | Standard | nd | 10.00 ± 0.22 | 7.00 ± 0.11 | nd |
| Chloroform | 2.00 | nd | nd | nd | nd |
| | 1.00 | nd | nd | nd | nd |
| | 0.50 | nd | nd | nd | nd |
| | 0.25 | nd | nd | 6.00 ± 0.39 | nd |
| | Standard | nd | 9.00 ± 0.25 | nd | nd |
| Butanol | 2.00 | nd | nd | 7.00 ± 0.18 | nd |
| | 1.00 | nd | nd | nd | nd |
| | 0.50 | nd | nd | nd | nd |
| | 0.25 | nd | nd | nd | nd |
| | Standard | 9.00 ± 0.12 | 8.00 ± 0.56 | 6.00 ± 0.72 | nd |
| Methanol | 2.00 | nd | nd | 7.00 ± 0.17 | nd |
| | 1.00 | nd | nd | nd | nd |
| | 0.50 | nd | nd | nd | nd |
| | 0.25 | nd | nd | nd | nd |
| | Standard | nd | 9.00 ± 0.30 | nd | nd |
| Water extract | 2.00 | nd | nd | 6.00 ± 0.15 | nd |
| | 1.00 | nd | 4.00 ± 0.10 | nd | nd |
| | 0.50 | nd | nd | nd | nd |
| | 0.25 | nd | nd | nd | nd |
| | Standard | nd | 8.00 ± 0.23 | nd | nd |

nd: Not detected.

4. Discussion

Neem is an indigenous herb which was chosen for this study. It belongs to the family of Meliaceae. The lack of information and medicinal importance on this plant facilitate this kind of study. *In vitro* total phenolics and flavonoids contents were determined for the crude extracts from the stems of neem. The total phenolics content of the crude extracts from the stems was expressed as gallic acid equivalents (GA) per gram of dry extract ranged between 12.18 to 128.5 mg GA/g. The results showed that the highest amount of the total phenolics was in ethyl acetate and the lowest was in hexane. The order was ethyl acetate > chloroform > butanol > methanol > hexane. The gallic acid was used as a standard calibration curve.

The total amount of flavonoids detected in the tested five crude extracts of stems was described. The total flavonoids content of different crude extracts from stems of neem was determined using UV-visible spectrophotometric method with aluminum chloride method[14]. Among the five crude extracts of neem stems, the highest amount of the total flavonoids was in hexane (484.50 mg/100 g dry powdered samples) and the lowest was in butanol (136.50 mg/100 g dry powdered samples) and followed the sequence ethyl acetate (339.50 mg/100 g), methanol (314.00 mg/100 g), chloroform (151.00 mg/100 g) and butanol (136.50 mg/100 g). The entire plant kingdom containing flavonoids are important components of human and animal diet. The different biological activities depend on the plant secondary metabolites and the regular consumption of them by human being may result in serious consequences for health, both positive and negative[16].

The antimicrobial result obtained from this study was presented. Almost all the crude extracts from stems of neem did not show any potential activity against the employed Gram-positive and Gram-negative bacterial strains. All the crude extracts did not show activity against *E. coli* except ethyl acetate. The ethyl acetate crude extract at concentrations 2.00 and 1.00 mg/mL showed small activity against *E. coli* bacterial strain. On the other hand, almost all crude extracts at all concentrations did not show any activity against *S. aureus* except ethyl acetate (2.00 mg/mL) and water (1.00 mg/mL). The small activity was obtained from all crude extracts of ethyl acetate and water at the concentrations of 2.00 mg/mL and 1.00 mg/mL with the values of 7 and 4 mm. Similarly, *P. aeruginosa* bacterial strain showed some activity at different concentrations of the employed crude extracts from the stems of neem. Both hexane and chloroform crude extracts at concentration of 0.25 mg/mL showed the activity with a value of 6 mm. But butanol, methanol and water crude extracts of stems at concentration of 2.00 mg/mL gave the range of activity with the value of 7 mm. However, *P. vulgaris* bacterial strain did not give any activity against all crude extracts at all concentrations. The total phenol, flavonoids and antimicrobial activities of different crude extracts from stems of neem were demonstrated from many previous studies[16-20].

From this study, it may be concluded that crude extracts from stems of neem are a good source of phytochemicals. Different extractives showed significant total phenolics and flavonoids contents. Hence, further studies are needed to be undertaken to isolate pure compounds from the crude extract and to better understand the mechanism of such actions scientifically.

Conflict of interest statement

We declare that we have no conflict of interest.

Acknowledgments

The author is grateful to Prof. Dr. Nafsiah Binti Shamsudin, Dean, College of Pharmacy and Nursing, University of Nizwa, Sultanate of Oman, for her continuous encouragement and laboratory facilities. The authors are also grateful to University of Nizwa, Nizwa, Sultanate of Oman for providing all chemicals and other expenses from their internal fund to carry out this graduation project. The authors wish to express sincere gratitude to the Central Instrument Laboratory, College of Agriculture and Marine Sciences, Sultan Qaboos University, Sultanate of Oman where some of the test were confirmed (Grant No. SOP/ADO/72/2014).

References

- [1] Sithisarn P, Gritsanapan W. Free radical scavenging activity and total flavonoid content of Siamese neem tree leaf aqueous extract from different locations. *Mahidol Univ J Pharm Sci* 2005; **32**(1-2): 31-5.
- [2] Norton E. *Neem: India's miraculous healing plant*. 1st ed. Rochester: Healing Arts Press; 1996, p. 15-21.
- [3] Maragathavalli S, Brindha S, Kaviyarasi NS, Annadurai BB, Gangwar SK. Antimicrobial activity in leaf extract of neem (*Azadirachta indica* Linn.). *Int J Sci Nat* 2012; **3**(1): 110-3.
- [4] Bhuiyan MM, Nishimura M, Matsumura S, Shimono T. Antibacterial effect of the crude *Azadirachta indica* neem barks extract on *Streptococcus sobrinus*. *Pediatr Dent J* 1997; **7**(1): 61-4.
- [5] Conrick J. *Neem: the ultimate herb*. 1st ed. Wisconsin: Louts Press; 2001, p. 1-6.
- [6] Chaturvedi P, Bag A, Rawat V, Jyala NS, Satyavali V, Jha PK. Antibacterial effects of *Azadirachta indica* leaf and bark extracts in clinical isolates of diabetic patients. *Natl J Integr Res Med* 2011; **2**(1): 5-9.
- [7] Siswomihardjo W, Badawi SS, Nishimura M, Hamada T. The difference of antibacterial effect of neem leaves and stick extracts. *Int Chin J Dent* 2007; **7**: 27-9.
- [8] Kiranmai M, Kumar CBM, Ibrahim M. Comparison of total flavonoid content of *Azadirachta indica* root bark extracts prepared by different methods of extraction. *Res J Pharm Biol Chem Sci* 2011; **2**(3): 254-5.
- [9] Kranmai M, Sri BU, Reddy DS, Kumar CBM, Ibrahim M. Evaluation of total phenolic contents and antiulcerogenic activity of root bark of *Azadirachta indica*. *Glob J Med Res* 2012; **12**(6): 23-9.
- [10] Asif M. Antimicrobial potential of *Azadirachta indica* against pathogenic bacteria and fungi. *J Pharmacogn Phytochem* 2012; **1**: 78-83.
- [11] Kokate CK, Purohit AP, Gokhale SB. *Pharmacognosy*. 45th ed. Pune: Nirali Prakashan; 2010.
- [12] Hismath I, Wan Aida WM, Ho CW. Optimization of extraction conditions for phenolic compounds from neem (*Azadirachta indica*) leaves. *Int Food Res J* 2011; **18**(3): 931-9.
- [13] Hossain MA, Al Kalbani MSA, Al Farsi SAJ, Weli AM, Al-Riyami Q. Comparative study of total phenolics, flavonoids contents and evaluation of antioxidant and antimicrobial activities of different polarities fruits crude extracts of *Datura metel* L. *Asian Pac J Trop Dis* 2014; **4**(5): 378-83.
- [14] Hossain MA, AL-Raqmi KA, AL-Mijizy ZH, Weli AM, Al-Riyami Q. Study of total phenol, flavonoids contents and phytochemical screening of various leaves crude extracts of locally grown *Thymus vulgaris*. *Asian Pac J Trop Biomed* 2013; **3**(9): 705-10.
- [15] Hossain MA, AL Harbi SR, Weli AM, Al-Riyami Q, Al-Sabahi JN. Comparison of chemical constituents and antimicrobial activities of three essential oils from three different brands' clove samples collected from Gulf region. *Asian Pac J Trop Dis* 2014; **4**(4): 262-8.
- [16] Mossini SA, Arrotía CC, Kimmelmeier C. Effect of neem leaf extract and neem oil on *Penicillium* growth, sporulation, morphology and ochratoxin A production. *Toxins (Basel)* 2009; **1**(1): 3-13.
- [17] Chaisawangwong W, Gritsanapan W. Total phenolic and flavonoid contents and free radical scavenging capacity of Siamese neem flower extracts. *Philipp Agric Sci* 2009; **92**(4): 431-6.
- [18] Girish K, Shankara Bhat S. Neem - a green treasure. *Electron J Biol* 2013; **4**(3): 102-11.
- [19] Stobiecki M, Kachlicki P. Isolation and identification of flavonoids. In: Grotewold E, editor. *The science of flavonoids*. New York: Springer; 2006, p. 47-70.
- [20] Ibrahim H, Aziz AN, Syamsir DR, Ali NAM, Mohtar M, Ali RM, et al. Essential oils of *Alpinia conchigera* Griff. and their antimicrobial activities. *Food Chem* 2009; **113**(2): 575-7.