



Contents lists available at [ScienceDirect](http://www.sciencedirect.com)

Asian Pacific Journal of Reproduction

journal homepage: www.apjr.net

Original research <http://dx.doi.org/10.1016/j.apjr.2016.11.001>

Effect of selenium on human sperm parameters after freezing and thawing procedures

Zahra Rezaeian³, Hossein Yazdekhesti¹, Sima Nasri², Zahra Rajabi¹, Parvin Fallahi³, Fardin Amidi^{1*}

¹Department of Anatomy, Tehran University of Medical Sciences, Tehran, Iran

²Payame-Noor University, Tehran, Iran

³Department of Infertility, Shariati Hospital, Tehran University of Medical Sciences, Tehran, Iran

ARTICLE INFO

Article history:

Received 3 Jul 2016

Received in revised form 31 Oct 2016

Accepted 1 Nov 2016

Available online 9 Nov 2016

Keywords:

Sperm parameters

Selenium

Oxidative stress

Freezing

Thawing

Infertility

ABSTRACT

Objective: To evaluate the effects of pre-freezing treatment of human semen samples with selenium on semen parameters after thawing procedure.

Methods: Each sample was divided into four groups: two groups were washed and other two remained unwashed. 5 µg/mL of selenium was added to one of the washed and unwashed group.

Results: The results showed that sperm motility in unwashed selenium-treated samples was higher compared to washed untreated samples ($P < 0.05$). DNA damage percentage in unwashed treated samples compared to both untreated samples showed no difference ($P > 0.05$) but in washed sperm samples, DNA damaged percentage decreased in treated samples compared to untreated ones ($P < 0.05$). Moreover, the effect of selenium on morphology of washed treated sperms was higher than unwashed untreated samples after thawing procedure. Normal morphology percentage in unwashed treated samples was significantly higher than both unwashed samples after thawing ($P < 0.001$). In the washed treated samples a higher number of sperms with normal morphology compared to washed untreated samples ($P < 0.001$) were observed.

Conclusions: Based on the results, we recommend 5 µg/mL of selenium should be used in infertility clinics to increase sperm quality after freezing and thawing procedures.

1. Introduction

Causes of infertility may include male, female or both and other unexplained factors. About 50% of infertility is believed that result from male factors associated with abnormal semen parameters. Some forms of abnormalities that may be observed in these men include oligospermia, asthenospermia, teratospermia (oligoasthenozoospermia) and azoospermia. Human sperm freezing is used extensively in assisted reproductive technology especially before chemotherapy, radiotherapy and some surgical procedures [1]. Oxidative stress and reactive oxygen radicals (ROS) can be produced in semen, and it has been shown to have some harmful effects on spermatozoa [2]. In semen abnormal spermatozoa, germinal cell precursors and leukocytes have potential to produce ROS [3]. On the other

hand, ROS generated by spermatozoa plays an important role in normal physiological processes such as sperm capacitation, acrosome reaction and maintenance of fertilization ability [4]. Due to high susceptibility of spermatozoa to oxidative stress and ROS, several damages on spermatozoa such as lipid peroxidation, DNA damage, apoptosis and sperm motility reduction can be induced [5].

Antioxidants are molecules that have ability to inhibit or reduce oxidative process in other molecules by scavenging released free radicals [6]. Antioxidants are divided into two groups, enzymatic and non-enzymatic. Enzymatic antioxidants include superoxide dismutase, glutathione peroxidase, catalase and glutathione reductase, whereas non-enzymatic antioxidants include vitamins A, C, E, pyruvate, glutathione, and coenzyme Q. Antioxidant function of selenium (Se) is mediated through glutathione peroxidase enzyme activity. There are several antioxidants in semen that are known to improve sperm quality such as vitamin E and C, as well as Se and Zn which are components of antioxidant systems. However, presumably these antioxidant agents are insufficient in preventing lipid peroxidation and

*Corresponding author: Fardin Amidi, Department of Anatomy, Tehran University of Medical Sciences, Tehran, Iran.

Fax: +98 21 66419072

E-mail: famidi@sina.tums.ac.ir

Peer review under responsibility of Hainan Medical College.

sperm plasma membrane injury during the freezing-thawing process because the protective antioxidant systems in sperms originate from cytoplasm and sperms discard their huge part of cytoplasm as residual body during final stages of differentiation. Freezing and thawing procedures cause significant reduction in motility and metabolic activity in sperms, however, vitamin E and Se have been found to improve freezing induced damages and reduced sperm motility [17]. During cryopreservation, semen is exposed to cold shock and atmospheric oxygen which in turn increases the susceptibility to lipid peroxidation due to higher production of ROS. Sperm plasma membrane is one of the important structures affected by cryopreservation which then can affect the viability and fertility of the sperm cells [8].

Se is an essential metal-like trace element in mammalian diet and its importance has been well established in humans [9]. Foods such as sea foods, liver, grain, egg yolk, milk, water and soil are major natural sources of Se with its levels generally depending on soil [10]. Typical dietary intake of Se in the US is 80–120 µg/d, and the recommended daily allowance is 70 µg in men and 50 µg in women [11]. Se is a component of the glutathione peroxidase enzyme which is known as an important antioxidant and a marker of oxidative stress [12]. Experimental evidences confirmed the positive effects of Se on human health [13]: reduction of disease symptoms and improvement of some disorders such as infertility, viral infections, cancers, cardiovascular disease and so on [14]. Also Se is an essential element in spermatogenesis and male fertility as well [15]. In a study that was performed by Moslemi *et al.* [16]. Effect of Se–vitamin E supplementation on different parameters of 690 infertile men and pregnancy rate was evaluated. Their results showed that supplemental Se and vitamin E may improve semen quality and pregnancy rate in idiopathic male infertility diagnosed with asthenoteratospermia or asthenospermia in semen analysis, and also have beneficial and protective effects especially on sperm motility [16]. Other studies have also shown that administration of Se to subfertile patients induced a statistically significant rise in sperm motility [17]. Various other studies have revealed the effects of cryopreservation on human sperms function and fertility whereas those conducted on animal model using Se as the most potent component of antioxidant system improved sperm quality and quantity [18]. For example, Sermak *et al.* demonstrated that addition of Se in the quantity of 1 µg/mL to cryomedium of ram semen improved survival rate following freezing-thawing process [19]. Unfortunately, there are few or no literature publications on the effects of post treatment of human semen with Se on semen parameters. As a first study, this research aimed to evaluate the effect of pre-freezing treatment of human semen with Se on different parameters of human sperm after thawing process.

2. Materials and methods

This study was approved by the Ethical Committee of the Faculty of Basic Sciences of Payame Noor University (Tehran, Iran). This experimental study was performed on 42 semen samples referred to Shariati infertility center in Tehran during February and March 2013, because of infertility problems of their wives but did not show any indications of infertility. All the 42 donors who participated had an informed written consent and the mean age of these individuals was (32.1 ± 3.9) years. In all participants, after 3 d of sexual abstinence, semen samples were collected by

masturbation in sterile containers and were transferred to laboratory immediately after ejaculation. The semen samples were allowed to complete liquefaction at 37 °C for 30 min. After liquefaction, a small aliquot was removed from each specimen and the sperm parameters were determined according to World Health Organization (WHO) guidelines [20]. Each sample was then divided into two groups: one group unwashed and the other washed. Washed samples were processed by swim-up technique. In swim-up technique, semen be overlaid directly with culture medium and the sperm allowed to swim from the seminal plasma into the culture medium. Washed and unwashed samples were also divided into two subgroups: one without Se treatment as control and the other with 5 µg/mL of sodium selenite (Na_2SeO_3). Control and treated specimens were then cryopreserved by liquid nitrogen vapor method [21].

An aliquot of the freezing medium equaling 25% of the original specimen volume was added to each of the samples. This process was repeated until the ratio of freezing medium to ejaculum (semen) became 1:1. Cryovials were loaded with 1 mL freezing medium/semen mixture and kept at –20 °C for 8 min. The samples were then exposed to liquid nitrogen vapor at –79 °C for 2 h before plunging into liquid nitrogen for storage at –196 °C. After 2 wk interval, the samples were thawed at room temperature for 5 min and then transferred to 37 °C water bath for 20 min. To remove the freezing medium, the thawed samples were diluted with Hams F₁₀(Biochrom.K.G) medium at a ratio of 1 volume of freezing medium/semen to 3 volumes of freezing medium and then washed by centrifugation at 300g for 7 min. The supernatant was removed and the pellet resuspended in 0.7 mL Hams F₁₀ medium. A small aliquot was removed from each sample for assessment of different parameters.

Light microscope (Nikon, Tokyo, Japan) with 40× magnification was used for motility evaluation. Sperms morphology was evaluated using Papanicolaou staining method [20]. In this method the staining was as follows; anterior area of acrosome stained light blue, posterior area dark blue, sperm neck slightly red, tail blue and excess residual cytoplasm pink.

Eosin-nigrosin staining was used for vitality evaluation. Stained sperms indicated permeability of sperm to dye and membrane damage whereas unstained sperms had healthy membrane which did not permit the dye to enter into the sperm cells. If sperm head stains red or dark pink it was considered dead whilst those whose head stained light pink or white was considered alive.

In order to evaluate sperm motility, we used light microscope with 40× magnification and based on WHO protocol, sperm motility was classified into three classes: progressive, non-progressive and immotile.

Ultimately, DNA damage was analyzed by toluidine blue staining based on WHO protocol. Toluidine blue staining is a metachromatic staining procedure used for chromatin structure evaluation. The basis of this procedure is the highly incorporation of Toluidine blue dye into damaged and compacted chromatin. When this dye is incorporated into histone and arginine-rich chromatin, the nucleus stains dark purple whereas its incorporation into protamine-rich chromatin stains the nucleus light blue [22]. To determine the significant differences between the studied groups, statistical analysis including mean \pm SD, correlation coefficients and two way Anova tests were performed in three independent experiments using Prism (version 5) software. All tests were performed at a confidence level of 95%.

3. Results

Results of effects of Se on different sperm parameters are shown in Figure 1.

3.1. Percentage of sperm cells with normal morphology in treated and control samples after freezing-thawing procedure

In this respect, our results showed that the percentage of sperms with normal morphology in washed treated samples was higher than both unwashed samples after thawing ($P < 0.001$). In washed treated sperms, a higher number of sperms with normal morphology compared to untreated samples ($P < 0.001$) were also observed. The effect of Se on morphology of washed treated sperms was higher than unwashed treated samples after thawing procedure, however, washed or unwashed status of sample did not affect morphology of sperms after thawing process ($P > 0.05$) (Figure 1a).

3.2. Percentage of dead sperm cells in treated and control samples after freezing-thawing procedure

In this research, Eosin-nigrosin staining procedure was employed for the evaluation of sperm cells viability. The results of this staining method was as follows: the head of dead sperm

cells stained red or dark pink and fresh sperms head showed white or light pink color. The percentage of dead sperm cells in both treated samples was lower compared to both untreated samples ($P < 0.001$). Our results also showed that the percentage of dead sperm cells in both washed samples was lower than both unwashed ($P < 0.05$) (Figure 1b).

3.3. Percentage of sperm cells with DNA damage in treated and control samples after freezing-thawing procedure

Percentage frequency of sperm cells with DNA damage in treated samples compared to both untreated samples showed a lower value ($P < 0.01$) but this difference was higher in washed sperm samples ($P < 0.05$) (Figure 1c).

3.4. Sperm cells motility alteration in treated and control samples after freezing-thawing procedure

Results of the evaluation of sperm cells motility showed that the percentage of motile sperms in unwashed treated samples was higher than washed untreated samples ($P < 0.05$). A significant difference was also found in the percentage of motile sperms between treated and untreated washed samples, however, in both treated sperms, motile sperms were higher than both untreated samples ($P < 0.001$). Our results also showed that Se




Figure 1. Effect of Se on different semen parameters, on morphology (a), on sperm vitality (b), on Percentage of sperm cells with DNA damage (c) and on sperm motility (d). Results are the mean \pm SD of three independent experiments. *: $P < 0.05$, **: $P < 0.01$, ***: $P < 0.001$, ns: not significant.

has no significant effect on sperm motility in unwashed group ($P > 0.05$). These results showed that Se is more effective on washed sperm cells than on unwashed samples ($P < 0.001$) (Figure 1d).

4. Discussion

Previous results showed that freezing procedure affects the different sperm parameters [23]. Our results showed that using Se as a component of antioxidant system optimizes sperm parameters after freezing-thawing procedure. Previous studies have also shown that freezing-thawing procedure decreases sperm motility, normal morphology, viability, and increases frequency of sperms with DNA damage [24]. In this regard, Meryman, *et al.* [25] stated that when sperm's environmental temperature reduces to zero, adenosine triphosphate production stops, sperm cells hibernate and all other activities come to a halt and/or sperm cells die afterwards. Cell damage occurs as the state of sperm cells changes from normal to freeze condition and vice versa, and the production of ROS has a very important role in this phenomena [25]. Different studies have indicated the effect of several antioxidants in the optimization of sperm quantity and quality after freezing-thawing process [26].

In this study, the effect of Se as a component of antioxidant system on different parameters of sperm after cryopreservation was studied. Kefer *et al.* [27] using x-ray fluorescence microscopy showed that Se is involved in spermatogenesis [27]. Moreover, Keskes-Ammar *et al.* [28] confirmed the protective and beneficial effects of Se on semen quantity and suggested its application in male infertility treatment [28].

In this current study, normal morphology percentage in washed treated and unwashed groups showed an increase compared to both untreated groups ($P < 0.001$). Seremak *et al.* [19] found that percentage of ram sperms with progressive movement in Se (1 μg Se/mL) treated group was identical to control group [19]. Watana be showed that in Se deficient mice, the proportion of abnormal sperm was high, ranging from 6.8% to 49.6%, and the most frequently occurring abnormalities in sperm shape were in the sperm head [33]. In another study [29] indicated that 6 and 8 ppm of dietary Se caused dose time dependent reduction in body and reproductive organs' weights and increased the number of morphologically abnormal spermatozoa [29]. Additionally, Gadea *et al.* [30] showed that addition of Glu to thawed sperm samples had no effect on optimization of standard semen parameters such as morphology [30]. In the same line, our results indicated that the addition of 5 $\mu\text{g}/\text{mL}$ of Se to human sperm before freezing-thawing procedures caused an increase in sperms with normal morphology. Sperm morphology is an important indication of seminiferous epithelial status, and degenerative changes in seminiferous epithelium have been found to have negative effects on sperm morphology [31].

On the effect of Se on sperm viability, Dorostkar *et al.* [31] showed that addition of 1 and 2 $\mu\text{g}/\text{mL}$ Se to water buffalo semen samples significantly increased sperm motility of fresh and equilibrated semen compared to the control without affecting sperms viability, in freezed-thawed semen however, extenders containing 1 and 2 $\mu\text{g}/\text{mL}$ of Se significantly improved sperm viability beside other parameters [32]. In some other studies, different doses of natural and synthetic antioxidants were shown to improve sperms viability [33].

Our investigation showed that 5 $\mu\text{g}/\text{mL}$ Se has a potential effect on the reduction of DNA damage. Eghbali *et al.* [33] performed an investigation on buffalo bulls and they found that 1 and 2 mg/mL of Se can increase semen antioxidants activity and reduce DNA damage after freezing and thawing procedure [34]. In terms of protective effect of Se against sperm cells' DNA damage, Xu *et al.* [34] determined the association between concentrations of cadmium and Se in seminal plasma and the degree of oxidative DNA damage in human spermatozoa. Their results showed that cadmium in seminal plasma can affect semen quality and oxidative DNA damage in human spermatozoa, and Se can protect against oxidative DNA damage in human sperm cells [35]. Similar to the study above, our results confirmed that more than semen Se concentration and treatment of sperm cells with Se before cryopreservation can have a protective effect against DNA damage. In different section of this study, we compared the effects of Se on different sperm parameters between washed and unwashed samples. Our results indicated that Se is more effective on different parameters of washed samples. Overall, semen has controlled generation of ROS and also involved in normal function and physiology of spermatozoa. From our results, it can be inferred that washing sperm samples can eliminate ROS and improve the effectiveness of antioxidants.

In this study, percentage of motile sperm in Se treatment groups (washed and unwashed groups) was significantly higher than control group. After thawing, class A (fast progressive) and B (progressive) sperm motility decreased to about 0%. However, class C (Non-progressive) and D (Non-motile) sperm motility did not decrease significantly. Freezing effect was so high that it could extremely increase class D sperm motility. In this study, motility mean was calculated from the summation of class C and D sperm motility after freezing and thawing procedure. Dorostkar *et al.* [31] evaluated different doses of Se (1, 2, 4 and 8 $\mu\text{g}/\text{mL}$) in buffaloes and they concluded that 1 and 2 $\mu\text{g}/\text{mL}$ Se dosages significantly improved sperm motility in treatment groups compared to control group and 4 and 8 $\mu\text{g}/\text{mL}$ Se dosages showed a significant decrease in sperm motility compared to control group. Florin *et al.* (2012) freezed Mangalita boar semen with the addition of vitamin E and C supplements and they observed that these supplements improved sperm motility after thawing [36]. Vitamin E can dissolve covalent linkages, and vitamin C can also inhibit free radicals production and thus can reduce harmful effects of freezing and thawing procedures. Ozkavukcu *et al.* [36] evaluated effects of freezing and thawing procedures on human sperms and they found that after freezing and thawing procedures, class A, B and C sperm motility decreased but class D increased [37]. These results confirmed our findings, however, Muratori, *et al.* [37] by adding glutathione peroxidase as antioxidant to domestic animal sperm samples found that sperm motility increased after freezing and thawing procedures compared to control group [38]. Effects of antioxidants on sperm quality could be as a result of very low usage of antioxidant doses or short period of treatment because supplements should reach a threshold concentration to have sufficient effects on oxidative stress [39]. In our investigation, we observed significant increase in sperm motility in Se treated groups, as well, that could have resulted from the effects of Se on sperm mitochondrial oxidative phosphorylation.

Based on our results, we suggest that 5 $\mu\text{g}/\text{mL}$ Se can improve sperm parameters and sperm vitality after freezing and

thawing procedures and this antioxidant could be utilized in clinic for treatment of infertility of sub fertile men. Moreover, our results showed that washed sperms (with or without Se) have better quality than unwashed sperms.

Conflict of interest statement

The authors declare that they have no conflict of interest.

Acknowledgments

We are grateful to all couples who participated in the study for their kind cooperation.

References

- Anger JT, Gilbert BR, Goldstein M. Cryopreservation of sperm: indications, methods and results. *J Urol* 2003; **170**(4): 1079-1084.
- Serviddio G, Bellanti F, Vendemiale G. Free radical biology for medicine: learning from nonalcoholic fatty liver disease. *Free Radic Bio Med* 2013; **65**: 952-968.
- Mahfouz R, Sharma R, Thiagarajan A, Kale V, Gupta S, Sabanegh E, et al. Semen characteristics and sperm DNA fragmentation in infertile men with low and high levels of seminal reactive oxygen species. *Fertil Steril* 2010; **94**(6): 2141-2146.
- Makker K, Agarwal A, Sharma R. Oxidative stress & male infertility. *Indian J Med Res* 2009; **129**(4): 357-367.
- Hess RA, Renato de Franca L. Spermatogenesis and cycle of the seminiferous epithelium. *Adv Exp Med Biol* 2008; **636**: 1-15.
- Kalthur G, Adiga SK, Upadhy D, Rao S, Kumar P. Effect of cryopreservation on sperm DNA integrity in patients with teratospermia. *Fertil Steril* 2008; **89**(6): 1723-1727.
- Paradiso Galatioto G, Gravina GL, Angelozzi G, Sacchetti A, Innominato PF, Pace G, et al. May antioxidant therapy improve sperm parameters of men with persistent oligospermia after retrograde embolization for varicocele? *World J Urol* 2008; **26**(1): 97-102.
- Watson P. The causes of reduced fertility with cryopreserved semen. *Anim Reprod Sci* 2000; **60**: 481-492.
- Akinloye O, Arowojolu AO, Shittu OB, Adejuwon CA, Osotimehin B. Selenium status of idiopathic infertile Nigerian males. *Biol Trace Elem Res* 2005; **104**(1): 9-18.
- Brown DG, Burk RF. Selenium retention in tissues and sperm of rats fed α torula yeast diet. *J Nutr* 1973; **103**(1): 102-108.
- Allowances NRCCoD, Food NRC, Board N. . *Recommended dietary allowances*, vol. 2941. National Academies; 1980.
- Mizuno K, Hirata S, Hoshi K, Shinohara A, Chiba M. Analysis of the phospholipid hydroperoxide glutathione peroxidase mRNA in the rat spermatozoon and effect of selenium deficiency on the mRNA. *Biol Trace Elem Res* 2000; **74**(1): 71-79.
- Rayman MP. The importance of selenium to human health. *Lancet* 2000; **356**(9225): 233-241.
- Beck MA, Levander OA, Handy J. Selenium deficiency and viral infection. *J Nutr* 2003; **133**(5): 1463S-1467S.
- Chavarro JE, Toth TL, Wright DL, Meeker JD, Hauser R. Body mass index in relation to semen quality, sperm DNA integrity, and serum reproductive hormone levels among men attending an infertility clinic. *Fertil Steril* 2010; **93**(7): 2222-2231.
- Moslemi MK, Tavanbakhsh S. Selenium-vitamin E supplementation in infertile men: effects on semen parameters and pregnancy rate. *Int J Gen Med* 2011; **4**: 99.
- MacPherson A, Scott R, Yates R. The effect of selenium supplementation in subfertile males. *Trace Elem Man Anim* 1993; **19**(6): 566-570.
- Zubair M, Ali M, Ahmad M, Sajid SM, Ahmad I, Gul ST. Effect of selenium and vitamin E on cryopreservation of semen and reproductive performance of animals (a review). *J Entomol Zool* 2015; **3**(1): 82-86.
- Seremak B, Udala J, Lasota B. Influence of selenium additive on ram semen freezing quality. *Electron J Pol Agric Univ* 1999; **2**(1). Series Animal Husbandry.
- Cooper TG, Noonan E, von Eckardstein S, Auger J, Baker HW, Behre HM, et al. World Health Organization reference values for human semen characteristics. *Hum Reprod Update* 2009; **dmp048**.
- Esteves SC, Sharma RK, Thomas A, Agarwal A. Cryopreservation of human spermatozoa with pentoxifylline improves the post-thaw agonist-induced acrosome reaction rate. *Hum Reprod* 1998; **13**(12): 3384-3389.
- Krzanowska H. Toluidine blue staining reveals changes in chromatin stabilization of mouse spermatozoa during epididymal maturation and penetration of ova. *J Reprod Fertil* 1982; **64**(1): 97-101.
- O'connell M, McClure N, Lewis S. The effects of cryopreservation on sperm morphology, motility and mitochondrial function. *Hum Reprod* 2002; **17**(3): 704-709.
- Asturiano JF, Marco-Jiménez F, Peñaranda DS, Garzón DL, Pérez L, Vicente JS, et al. Effect of sperm cryopreservation on the European eel sperm viability and spermatozoa morphology. *Reprod Domest Anim* 2007; **42**(2): 162-166.
- Meryman HT. Cryopreservation of living cells: principles and practice. *Transfusion* 2007; **47**(5): 935-945.
- Topragaleh TR, Shahverdi A, Rastegarnia A, Ebrahimi B, Shafiepour V, Sharbatoghli M, et al. Effect of cysteine and glutamine added to extender on post-thaw sperm functional parameters of buffalo bull. *Andrologia* 2014; **46**(7): 777-783.
- Kefer JC, Agarwal A, Sabanegh E. Role of antioxidants in the treatment of male infertility. *Int J Urol* 2009; **16**(5): 449-457.
- Keskes-Ammar L, Feki-Chakroun N, Rebai T, Sahnoun Z, Ghazzi H, Hammami S, et al. Sperm oxidative stress and the effect of an oral vitamin E and selenium supplement on semen quality in infertile men. *Syst Biol Reprod Med* 2003; **49**(2): 83-94.
- Kaur R, Kaur K. Effects of dietary selenium (SE) on morphology of testis and cauda epididymis in rats. *Indian J Physiol Pharmacol* 2000; **44**(3): 265-272.
- Gadea J, García-Vazquez F, Matás C, Gardón JC, Cánovas S, Gumbao D. Cooling and freezing of boar spermatozoa: supplementation of the freezing media with reduced glutathione preserves sperm function. *J Androl* 2005; **26**(3): 396-404.
- World Health Organization. WHO laboratory manual for the examination and processing of human semen. Switzerland: WHO Press; 2010.
- Dorostkar K, Alavi-Shoushtari SM, Mokarizadeh A. Effects of in vitro selenium addition to the semen extender on the spermatozoa characteristics before and after freezing in water buffaloes (*Bubalus bubalis*). *Vet Res Forum* 2012; **3**(4): 263-268.
- Breining E, Beorlegui NB, O'Flaherty CM, Beconi MT. Alphatocopherol improves biochemical and dynamic parameters in cryopreserved boar semen. *Theriogenology* 2005; **63**(8): 2126-2135.
- Eghbali M, Alavi-Shoushtari SM, Asri-Rezaei S, Khadem Ansari M-H. Calcium, magnesium and total antioxidant capacity (TAC) in seminal plasma of water buffalo (*Bubalus bubalis*) bulls and their relationships with semen characteristics. *Vet Res Forum* 2012; **1**(1): 12-20.
- Xu DX, Shen HM, Zhu QX, Chua L, Wang QN, Chia SE, et al. The associations among semen quality, oxidative DNA damage in human spermatozoa and concentrations of cadmium, lead and selenium in seminal plasma. *Mutat Res Genetic Toxicol Environ Mutagen* 2003; **534**(1): 155-163.
- Ghiuru FV, Ioan L, Roman I, Hettig A, Marius Z, Miclea V. Antioxidant medium for Mangalita Boar Semen cryopreservation. Bulletin of university of agricultural sciences and veterinary medicine Cluj-Napoca. *Animal Science and Biotechnologies* 2010; **67**(1-2).
- Ozkavukcu S, Erdemli E, Isik A, Oztuna D, Karahuseynoglu S. Effects of cryopreservation on sperm parameters and ultrastructural morphology of human spermatozoa. *J Assist Reprod Gen* 2008; **25**(8): 403-411.
- Muratori M, Luconi M, Marchiani S, Forti G, Baldi E. Molecular markers of human sperm functions. *Int J Androl* 2009; **32**(1): 25-45.
- Agarwal A, Said TM. Oxidative stress, DNA damage and apoptosis in male infertility: a clinical approach. *BJU Int* 2005; **95**(4): 503-507.