

**GREEN CHEMISTRY — CURRENT AND FUTURE ISSUES****Prof. Sineen Taj***D.G. Tatkare Mahavidyalay, Mangaon, Raigad***Abstract**

*The beginning of green chemistry is frequently considered as a response to the need to reduce the damage of the environment by man-made materials and the processes used to produce them. A quick view of green chemistry issues in the past decade demonstrates many methodologies that protect human health and the environment in an economically beneficial manner. This article presents selected examples of the implementation of green chemistry principles in everyday life in industry, the laboratory and in education. A brief history of green chemistry and future challenges are also mentioned.*

**Keywords:** *Green chemistry, green analytical chemistry, clean chemistry, atom economy, sustainable development.*



*Scholarly Research Journal's is licensed Based on a work at [www.srjis.com](http://www.srjis.com)*

**❖ History:**

The term green chemistry was first used in 1991 by P.T. Anastas in a special program launched by the US Environmental Protection Agency (EPA) to implement sustainable development in chemistry and chemical technology by industry, academia and government. In 1995 the annual US Presidential Green Chemistry Challenge was announced. Similar awards were soon established in European countries. In 1996 the Working Party on Green Chemistry was created, acting within the framework of International Union of Applied and Pure Chemistry. One year later, the GreenChemistry Institute (GCI) was formed with chapters in 20 countries to facilitate contact between governmental agencies and industrial corporations with universities and research institutes to design and implement new technologies. The first conference highlighting green chemistry was held in Washington in 1997. Since that time other similar scientific conferences have soon held on a regular basis. The first books and journals on the subject of green chemistry were introduced in the 1990s, including the Journal of Clean Processes and Products (Springer-Verlag) and Green Chemistry, sponsored by the Royal Society of Chemistry. Other journals, such as Environmental Science and Technology and the Journal of Chemical Education, have

devoted sections to green chemistry. The actual information also may be found on the Internet.

❖ **The 12 Principles of Green Chemistry:**

**1. Prevention**

It is better to prevent waste than to treat or clean up waste after it has been created.

**2. Atom Economy**

Synthetic methods should be designed to maximize the incorporation of all materials used in the process into the final product.

**3. Less Hazardous Chemical Syntheses**

Wherever practicable, synthetic methods should be designed to use and generate substances that possess little or no toxicity to human health and the environment.

**4. Designing Safer Chemicals**

Chemical products should be designed to effect their desired function while minimizing toxicity.

**5. Safer Solvents and Auxiliaries**

The use of auxiliary substances (e.g. solvents, separation agents, etc.) should be made unnecessary wherever possible and innocuous when used.

**6. Design for Energy Efficiency**

Energy requirements of chemical processes should be recognized for their environmental and economic impacts and should be minimized. If possible, synthetic methods should be conducted at ambient temperature and pressure.

**7. Use of Renewable Feedstocks**

A raw material or feedstock should be renewable rather than depleting whenever technically and economically practicable.

**8. Reduce Derivatives**

Unnecessary derivatization (use of blocking groups, protection/deprotection, temporary modification of physical/chemical processes) should be minimized or avoided if possible, because such steps require additional reagents and can generate waste.

**9. Catalysis**

Catalytic reagents (as selective as possible) are superior to stoichiometric reagents.

**10. Design for Degradation**

Chemical products should be designed so that at the end of their function they break down into innocuous degradation products and do not persist in the environment.

## **11. Real-time analysis for Pollution Prevention**

Analytical methodologies need to be further developed to allow for real-time, in-process monitoring and control prior to the formation of hazardous substances.

## **12. Inherently Safer Chemistry for Accident Prevention**


Substances and the form of a substance used in a chemical process should be chosen to minimize the potential for chemical accidents, including releases, explosions, and fires.

### **❖ Examples of Implementation of Green Chemistry Principles Into Practice:**

In some industrial chemical processes, not only waste products but also the reagents used for the production, may cause a threat to the environment. The risk of exposure to hazardous chemical compounds is limited in daily work by protective equipment such as goggles, breathing apparatus, face-guard masks, etc. According to the principles of green chemistry, a threat can be eliminated in a simpler way, by applying safe raw materials for production process. Large amounts of adipic acid [ $\text{HOOC}(\text{CH}_2)_4\text{COOH}$ ] are used each year for the production of nylon, polyurethanes, lubricants and plasticizers. Benzene — a compound with convinced carcinogenic properties — is a standard substrate for the production of this acid. Chemists from State University of Michigan developed green synthesis of adipic acid using a less toxic substrate. Furthermore, the natural source of this raw material — glucose — is almost inexhaustible. The glucose can be converted into adipic acid by an enzyme discovered in genetically modified bacteria. Such a manner of production of this acid guards the workers and the environment from exposure to hazardous chemical compounds.

Green chemistry tries, when possible, to utilize benign, renewable feedstocks as raw materials. From the point view of green chemistry, combustion of fuels obtained from renewable feedstocks is more preferable than combustion of fossil fuels from depleting finite sources.


For example, many vehicles around the world are fueled with diesel oil, and the production of biodiesel oil is a promising possibility. As the name indicates, biodiesel oil is produced from cultivated plants oil, e.g. from soya beans. It is synthesized from fats embedded in plant oils by removing the glycerine molecule (Fig. 1) — a valuable raw material for soap production.



**Fig. 1. Reaction for biodiesel oil production**

Biodiesel oil also can be obtained from wasted plant oils, e.g. oils used in restaurants. In the technological process, a potential waste product is transformed into valuable fuel. (Combusted biodiesel oil smells like fried potatoes.) The advantages of using biodiesel oil are obvious. It's fuel from renewable resources and contrary to normal diesel oil, the combustion of biodiesel does not generate sulphur compounds and generally does not increase the amount of carbon dioxide in the atmosphere. CO<sub>2</sub> formed in the combustion of fuel was removed earlier by plant. The great threats to the environment are organic solvents applied in many syntheses. The new solutions for practical synthesis aim at complete elimination of solvents or to substitute the compounds belonging to VOCs by cheap technological media, harmless for humans and the environment.

The use of supercritical fluids (SCFs) in chemical processes is becoming more and more prevalent. The term "supercritical fluids" comprises the liquids and gases at temperatures and pressures higher than their critical temperatures and pressures (Fig. 2). Above the critical point the liquid-vapour phase boundary disappears while the formed phase exhibits properties between those of gas and liquid. High compressibility of supercritical fluids in the vicinity of the critical point makes it easy to adjust density and solution ability by a small change of temperature or pressure. Due to this, the supercritical fluids are able to dissolve many compounds with different polarity and molecular mass. Among many possible supercritical fluids, fulfilling the green chemistry demands as the reaction media are carbon dioxide (scCO<sub>2</sub>) and water(scH<sub>2</sub>O).



**Fig. 2 Phase diagram showing supercritical fluid region.**

Carbon dioxide as a supercritical fluid is most frequently used as medium for reactions. It is inflammable, easily available (from natural sources, from power engineering) and cheap. Its application gives considerable energy savings because the critical point is easy to reach due to a low evaporation heat of CO<sub>2</sub>. Carbon dioxide as a supercritical fluid dissolves non-polar compounds and some polar (e. g. methanol, acetone) like fluorocarbon solvents. The discovery of a new surfactant with high surface activity in supercritical carbon dioxide opened a way to new processes in textile and metal industries and for dry cleaning of clothes.

Micells Technologies Company offers technology for removal of stains using liquid carbon dioxide instead of the perchloroethylene more commonly applied. Most of the common liquids (e.g., water, ethanol, benzene, etc.) are molecular. That is, regardless of whether they are polar or non-polar, they are basically made up of molecules. However, since the early 1980s an exciting new class of room-temperature liquids have become available. These are the ambient-temperature ionic liquids. Unlike the molecular liquids, regardless of the degree of association, they are basically constituted of ions. This gives them the potential to behave very differently from conventional molecular liquids when they are used as solvents. Room-temperature ionic liquids are considered to be environmentally benign reaction media because they are low-viscosity liquids with no measurable vapor pressure.

However, the lack of sustainable techniques for the removal of products from the room-temperature ionic liquids has limited their application. Professors Brennecke and Beckman have shown that environmentally benign carbon dioxide, which has been used extensively, both commercially and in research for the extraction of heavy organic solutes, can be used to extract nonvolatile organic compounds from room temperature ionic liquid. They found that extraction of a material into carbon dioxide represents an attractive means for recovery of products from ionic liquids because:

(a) CO<sub>2</sub> dissolves in the ionic liquid to facilitate extraction, and  
(b) the ionic liquid does not dissolve appreciably in the CO<sub>2</sub>, so that the product can be recovered in pure form. The research groups of Professors Brennecke and Beckman have shown that ionic liquids (using 1-butyl-3-methylimidazolium hexafluorophosphate as a prototype) and CO<sub>2</sub> exhibit extremely unusual, and very attractive, phase behavior. The solubility of CO<sub>2</sub> in ionic liquids is substantial, reaching mole fractions as high as 0.6 at just 8 MPa. Yet the two phases do not become completely miscible, so CO<sub>2</sub> can be used to extract compounds from the ionic liquids. Most importantly, the composition of the CO<sub>2</sub>-rich phase is essentially pure CO<sub>2</sub>, and there is no measurable cross-contamination of the CO<sub>2</sub> by the ionic liquid. Moreover, non-volatile organic solutes (using naphthalene as a prototype) may be quantitatively extracted from the ionic liquid with CO<sub>2</sub>, demonstrating the tremendous potential of ionic liquid/CO<sub>2</sub> biphasic systems as environmentally benign solvents for combined reaction and separation schemes.

❖ **Teaching of Green Chemistry:**

The main rule: Teaching must be in harmony with practice.

The question of how to educate the future generation of chemists possessing the skill and knowledge to practice environmentally friendly chemistry lies in the center of educational materials related to green chemistry. Education is especially important in the popularization of green chemistry. It is realized both at the level of academia and on the level of pro-environmental education for broad circles of society. Young chemists are currently acquainted with new methods of organic compound syntheses instead of traditional methods and with new analytical chemistry techniques allowing them to assess the state of environmental pollution in an increasing number of high schools. Different international institutions, i.e. the American Chemical Society (ACS) and Polish Chemical Society (PTChem), are active in publishing materials that promote the rules and achievements of green chemistry. The green chemistry program should lead to sustainability by designing and using the methods in which natural raw materials will be economically processed, rational usage of energy sources, elimination of hazardous gaseous, liquid and solid wastes and by introduction of safety products for man. The popularization of green chemistry in schools, among the workers at plants of chemical industry and distributors of chemical products is also very important. The broad usage of greenGreen Chemistry ...393 chemistry achievements will enable us to balance eco-development profitable for society, economy and the environment. The numerous educational materials, available currently on

market and on the Internet, are very useful in everyday teaching of green chemistry principles.

### ❖ **Conclusions**

Green chemistry is not a new branch of science. It is a new philosophical approach that through application and extension of the principles of green chemistry can contribute to sustainable development. Presently it is easy to find in the literature many interesting examples of the use of green chemistry rules. They are applied not only in synthesis, processing and using of chemical compounds. Many new analytical methodologies are also described which are realized according to green chemistry rules. They are useful in conducting chemical processes and in evaluation of their effects on the environment. The application of proper sample preparation techniques, (e.g.SPME, SPE, ASE) allows us to obtain precise and accurate results of analysis. Great efforts are still undertaken to design an ideal process that starts from non-polluting initial materials, leads to no secondary products and requires no solvents to carry out the chemical conversion or to isolate and purify the product. However, more environmentally friendly technologies at the research stage do not guarantee that they will be implemented on an industrial scale. Adoption of environmentally benign methods may be facilitated by higher flexibility in regulations, new programs to facilitate technology transfer among academic institutions, government and industry and tax incentives for implementing cleaner technologies. Furthermore, the success of green chemistry depends

on the training and education of a new generation of chemists. Student at all levels have to be introduced to the philosophy and practice of green chemistry. Finally, regarding the role of education in green chemistry:

THE BIGGEST CHALLENGE OF GREEN CHEMISTRY IS TO USE ITS RULES IN PRACTICE.

### **References**

- ANASTAS P. T., WARNER J. C. *Green Chemistry: Theory and Practise. Oxford University Press, Oxford 1998.*
- NAMIECENIK J., WARDENCKI W. *Solventless sample preparation techniques in environmental analysis. J. High Resol. Chromatogr. 23, 297, 2000.*
- SATO K., AOKI M., NOYORI R. *A "Green" Route to Adipic Acid: Direct Oxidation of Cyclohexenes with 30 percent hydrogen peroxide. Science. 281, 1646, 1998.*
- <http://www.epa.gov/greenchemistry>
- BARDLEY D., DYSON P., WELTON T. *Room temperature ionic liquids. Chem. Rev. 9(5), 18, 2000.*

- ROMANO U., GARBASSI F. *The environmental issue. A challenge for new generation polyolefins. Pure Appl. Chem.* 72, 1383, 2000.
- NICOLAS N., BENVEGNI T., PLUSQUELLEC D. *Surfactants from renewable resources. Actualite Chimique* 11-12, 70, 2002.
- STASHENKO E. E., PUERTAS A. M., SALGAR W., DELGADO W., MARTINEZ J. R. *Solid-phase microextraction with on fibre derivatization applied to the analysis of volatile carbonyl compounds. J. Chromatogr. A.* 886, 175, 2000.
- ACARDI A., BIANCHI G., DI GIUSEPPE S., MARINEL-LI F. *Gold catalysis in the reaction of 1,3-dicarbonyls with nucleophiles. Green Chemistry.* 5(1), 64, 2003.
- SCOTT G. *Green polymers. Polym. Degrad. Stab.* 68(1),1, 2000.
- TUNDO P., SELVA M., MEMOLI S. *Dimethylcarbonate as a green reagent. ACS Symp. Ser., 767 (Green Chemical Syntheses and Processes),* 87, 2000.
- MERRILL M., PARENT K., KIRCHHOFF M. *Green Chemistry. Stopping Pollution before it starts. Chem Matters.* April, 7, 2003.