

## Screening for group B streptococci in antenatal women: Isolation and its antibiogram

Balan K.<sup>1,\*</sup>, Abirami Lakshmy<sup>2</sup>, Ashwini MA<sup>3</sup>, Vijayalakshmi TS.<sup>4</sup>, Sheila Doris Devamani<sup>5</sup>

<sup>1</sup>Associate Professor, <sup>2</sup>Assistant Professor, <sup>3</sup>2nd year Student, <sup>4</sup>Professor & HOD, <sup>5</sup>Professor, Dept. of Microbiology, Karpaga Vinayaga Institute of Medical Sciences and Research Center, Madurantagam, Tamil Nadu, India

**\*Corresponding Author:**

Email: balankbalan@gmail.com

### Abstract

Group B streptococci (GBS) is one of those pathogens causing neonatal septicaemia and neonatal meningitis. Objective of this study is to screen antenatal women for Group B Streptococcus colonization and its antibiotic sensitivity pattern by Kirby bauer disk diffusion method. This study was carried out in Department of Microbiology during a period of 4 months. A total of 100 pregnant women attending antenatal clinics in the age group 20-40 years were included in the study. Two vaginal swab were taken aseptically and immediately transported to the laboratory for processing. One swab for direct gram staining and other swab was inoculated into sheep blood agar and incubated at 37 °C for 24-48 hours. Identification of organism was based on gram staining, colony morphology, catalase reaction, Christie Atkins Munch Peterson (CAMP) and Hippurate hydrolysis test. Results of this study show that prevalence rate for GBS colonization is 8%. Pregnant women less than 20 years of age and primigravida are more commonly colonized with Group B Streptococci. All the isolates were sensitive to Penicillin, Erythromycin, Ampicillin, Gentamicin, Ceftriaxone and Chloramphenicol. Hence all pregnant women should be screened for Group B Streptococci carriage between 35 and 37 weeks of gestation and intrapartum antibiotic prophylaxis be given to colonized women at the time of labour or rupture of membranes.

**Keywords:** Group B Streptococci, Antenatal pregnant women, Prevalence, Antibiotic susceptibility pattern.

### Introduction

Group B streptococci (GBS) also referred as *Streptococcus agalactiae*, is a Gram positive opportunistic pathogen which is beta hemolytic streptococci of Lancefield group B, that colonizes the gastrointestinal and genitourinary tract upto 50 % of healthy individuals. However in neonates GBS is one of the most common organisms causing neonatal sepsis and meningitis.<sup>1</sup> The antenatal women GBS colonizes in 15-20% of pregnant women in the lower vaginal tract. During passage of neonate via the mother's birth canal newborn acquires the GBS from the lower vaginal tract since the mother's birth canal is the principal reservoir of this GBS for infants.<sup>2</sup> Very few Indian studies have been carried out, to study the prevalence of GBS colonization and its antibiogram in the vaginal flora of the pregnant women. The incidence of GBS colonization in the vaginal flora of pregnant women varied from 0.48 to 24.3%.<sup>3</sup> Hence the present study was done to know the prevalence of GBS colonization and the antibiotic susceptibility pattern in the antenatal pregnant women. GBS is mostly thought of as a commensal organism that resides in the genitourinary and lower gastrointestinal tract of healthy adults. Approximately 20% to 40% of women are colonized with GBS during pregnancy. While maternal colonization is usually asymptomatic, it places the newborn at immediate risk for vertical transmission.<sup>4</sup> Up to 50% of newborns born to colonized mothers will also become colonized with GBS. Among colonized newborns, 1-2% will develop early onset acquisition from colonized individuals. Guidelines shows all

antenatal pregnant women at 35-37 weeks of gestation are universally screened for GBS, and if colonized, intrapartum antibiotic prophylaxis to reduce the risk of transmission from mother to neonates. Antibiotic prophylaxis reduces the incidence of early onset Group B streptococcal disease by 36%, however the incidence of Late onset Group B streptococcal disease has not been affected.<sup>5</sup> Henceforth GBS remains the leading cause of bacterial disease in newborns with an overall incidence rate of ~0.3 infections per 1,000 live births. Evidence shows that GBS develop resistant to second line antibiotics in recent days and is emerging pathogen among adult and elderly populations which emphasize the need to make alternative measures for preventing disease in both newborns and the elderly.<sup>6</sup>

### Materials and Methods

A cross sectional study was conducted in the department of Microbiology for the period of 4 months. A total of 100 antenatal pregnant women attending antenatal clinics were included in our study. The inclusion criteria were the all antenatal pregnant women in the age group 20-40 years willingness to participate and provide informed consent. The exclusion criteria were those women undergone pelvic examination prior to vaginal swab, already receiving antibiotics and antepartum haemorrhage. Prior to vaginal examination, two sterile vaginal swab were taken aseptically and were immediately transported to the Microbiology department. One swab is used for direct gram staining and the Culture was done with other swab which was inoculated on to 5% sheep blood agar and it was further

incubated at optimum temperature 37°C for 24-48 hours. Preliminary identification of GBS was done by Direct Gram's staining shows gram positive cocci arranged in short chain and in culture colonies are 1 mm in diameter smooth, convex, moist, regular, surrounded by a small hazy zone of  $\beta$  hemolysis.

GBS was further tested for catalase activity where 3% hydrogen peroxide was used, presence of effervescence shows a catalase positive and further confirmed by CAMP test, hippurate hydrolysis test and Lancefield grouping. The serogrouping for GBS was done by using latex agglutination test available as commercial detection kit.

The Antibiotic sensitivity test was performed on Muller Hinton agar plates by Kirby-Bauer disk diffusion method as per Clinical and Laboratory Standards Institute (CLSI) guidelines. The following antibiotics (Hi-Media, Mumbai, India) were tested by disk diffusion method: Penicillin (10  $\mu$ g/disc), Erythromycin (15  $\mu$ g/disc), Ampicillin (10  $\mu$ g/disc), Gentamicin (10  $\mu$ g/disc), Chloramphenicol (30  $\mu$ g/disc) and Ceftriaxone (30  $\mu$ g/disc).<sup>11</sup> Data was analysed by chi-square test, paired 't' test, and a p-value of <0.05 was taken as statistically significant.

## Results and Discussion

In our study out of 100 women, 8(8%) were showed GBS colonization. Most of the isolates were from age group  $\leq 20$  years (16%) followed by 21-25 years (5.5%) Table 1. The GBS was isolated maximum from the primigravida 4(12.5%) followed by second gravida 2(8%) and 01(4.3%) were multigravida. It was observed GBS was not isolated from multigravida women above 20 years of age. Thenceforth the correlation between GBS colonization with age & gravida was statistically not significant.

All most all the strains were 100% sensitive to beta lactam antibiotic (penicillin, ampicillin, cefuroxime and ceftriaxone), Erythromycin, ampicillin, gentamicin, Chloramphenicol,

In newborn and adults Group A Streptococci causes variety of life threatening serious diseases like septicemia, meningitis and pneumonia which has given an alarm in recent years. However during the early onset of this infection lung is the major organ involved for the initiation of infection and for the clinical sequelae, but for the those infants whose death occurred within few hour of birth inflammatory response is minimal or absent. GBS shows very high morbidity and mortality and most of the clinicians are not aware about the prevalence of GBS, so an attempt was made to find out the prevalence of GBS in pregnant women.

Present study shows the prevalence rate of GBS colonization was 8% in third trimester. The incidence rate was significantly high in primigravida followed by second gravida. The incidence in our study (8%) which correlates with two Indian studies by Annie Rajaratnam *et al* and Vinay haraze *et al* were colonization rate is

8.3% and 7.5%.<sup>12,13</sup> In contrast by Motlova *et al* and Konikkara *et al* reported that the prevalence of GBS carriage is 29.3% and 12% was relatively high.<sup>14,15</sup> Many reasons are attributed for GBS colonization and the rate of colonization varies from place to place depending upon various factors like socioeconomic status, sampling techniques to collect the clinical sample. However genetic polymorphism and Ethnicity might play a very important role in variation of the rates of infection with GBS.

In the present study most of the cases belongs to age group  $\leq 20$  years (16%) followed by the age group 21-25 years (6.6%), thus most of the GBS colonized pregnant women were from teenage and the younger age group was found to be at par with that of Vinay hazare *et al* and Tsering Chomu *et al*.<sup>13,16</sup> The reason for the predisposition of women younger  $\leq 20$  years of age to vaginal colonization with GBS is less apparent. This relationship could be the results of age related development of local or humoral immunological responses that interfere with mucosal attachment and/or persistence of GBS.

In the present study 9.8% of GBS isolated from third trimester which correlates with Fatemi *et al*, however in contrast Baker *et al* found that the colonization rate almost doubles between the second trimester and delivery.<sup>17,18</sup>

All strains were 100% sensitive penicillin, Erythromycin, ampicillin, gentamicin, Chloramphenicol, Ceftriaxone were Arora *et al* reported that 60 GBS were susceptible to Penicillin, Ampicillin & erythromycin.<sup>19</sup> In 2012 Vinay *et al* showed all the 15 strains were 100% sensitive Penicillin, Ampicillin & erythromycin.<sup>13</sup>

**Table 1: Age wise distribution of study population in relation to Group B streptococci**

Age (Years)	No. of Mothers	Group B streptococci infection			
		Present		Absent	
		No	%	No	%
$\leq 20$	24	4	16	20	83
21-25	30	2	6.6	28	93
26-30	19	1	5.2	18	94.7
31-35	14	1	5.2	13	92.8
36-40	9	0	0	9	100
$\geq 40$	4	0	0	4	100
Total	100	8	8	92	92

$X^2$  p>0.05 insignificant

**Table 2: Distribution of Pregnant women according to gravida and Group B Streptococci infection**

Gravida	No. of Mothers	Group B streptococci infection			
		Present		Absent	
		No	%	No	%
Primi	32	4	12.5	28	87.5
Second	25	2	8	23	92

Third	20	1	5	19	95
Multi	23	1	4.3	22	95.6
Total	100	8	8	92	92

$X^2 p > 0.05$  insignificant

### Conclusion

Colonization of GBS in the genital tract of the parturient women is the single most important factor for GBS colonizer in the newborn babies. However various strategy are used to prevent the transmission of GBS from mother to newborn, one such method is intravenous intrapartum antibiotic prophylaxis by penicillin or ampicillin to avoid early –onset GBS disease in the infant and to treat all neonates born to such women with penicillin shortly after birth.

### Financial support and sponsorship

This work was supported by Indian Council of Medical Research (ICMR) STS 2017-03057, Ministry of Health and Family welfare, Government of India

### Conflicts of Interest: Nil

### References

- Verani JR, Schrag SJ. Group B streptococcal disease in infants: Progress in prevention and continues challenges. *Clin Perinatol* 2010;37:375-92
- Edwards MS, Baker CJ. Group B Streptococcal infections. In : Remington JA ed. *Infectious Diseases of the Fetus and Newborn Infant*. Philadelphia. WB Saunders, 2001:1091-156
- Powers RJ, Wirtschafter D. Prevention of Group B Streptococcus early onset disease: A toolkit by the California Perinatal quality care Collaborative. *J Perinatol* 2010;30:77-87
- Weston EJ, Pondo T, Lewis MM, et al. The burden of invasive early onset neonatal sepsis in the United states. 2005-2008. *Pediatric Infections Disease Journal* 2011;30:937-41.
- Stoll, Barbagera J, Schuchat, Anne MD. Maternal carriage of Group B streptococci in developing countries. *Pediatric Infections Disease Journal* 1998;17(6):499-503.
- Hoogkamp-Korstanje JA, Gerards LJ, Cats BP: Maternal carriage and neonatal acquisition of Group B Streptococci. *The Journal of Infectious Disease*. 1982;145(6):800-3.
- Tettelin H, Massignani V, Cieslewicz MJ, Donati C, Medini D, et al. Genome analysis of multiple pathogenic isolates of *Streptococcus agalactiae*: implications for the microbial "pan-genome". *Proc Natl Acad Sci U S A*. 2005;102:13950-5.
- Medini D, Donati C, Tettelin H, Massignani V, Rappuoli R. The microbial pangenome. *Curr Opin Genet Dev*. 2005;15:589-94.
- Mira A, Martin-Cuadrado AB, D'Auria G, Rodriguez-Valera F. The bacterial pan-genome: a new paradigm in microbiology. *Int J Microbiol*; 2010 13:45-57.
- Lakshmi V, Das S, Shivananda PG, Savitri P, Roa K: Incidence of group B hemolytic Streptococci in the vaginal flora of pregnant women. *Indian Journal of Pathology & Microbiology*, 1998;31(3):240-4.
- Clinical Laboratory Standards Institute (CLSI). Performance Standards for Antimicrobial Susceptibility Testing; 27<sup>th</sup> Informational Supplement. M100-S27. Wayne: The Institute, 2017.
- Annie Rajaratnam, Thomas S, Kuruvilla, Benna Antony. Prevalance of Group B streptococci colonization among pregnant women in a tertiary care hospital in coastal Karnataka. *Int J. Appl. Biol. Pharm. Technol* 2013;308
- Vinay Hajare et al. Antibioqram of group B streptococci isolated from the vagina of pregnant women in third trimester of pregnancy. *People J, Scientific Res*. 2012;5(2):22-6.
- Motlova J, Strakova L, Urbaskova P, Sak P, Sever T. Vaginal and rectal carriage of *Streptococcus agalactiae* in the Czech Republic: incidence, serotypes distribution & susceptibility to antibiotics. *Indian J. Med Res* 2004;119:84-7.
- Konikkara KP, Baliga S, Shenoy SM, Bharati B. Comparison of various culture methods for isolation of Group B streptococci from intrapartum vaginal colonization 2013. *J. Lab Physicians*; 5:42-5.
- Tsering Chomu Dechen, Kar Sumit, Pal Ranabir. Correlates of vaginal colonization with Group B streptococci among Pregnant women. *Global Infect Dis* 2010;2(3):236-41.
- Farnaz Fatemi et al. Colonization rate of Group B streptococci in pregnant women using GBS Agar Medium. *Acta Medica Iranica* 2009;47(1):25-30.
- Baker CJ et al. Vaginal colonization with Group B streptococci: A study in college women. *J Infect Dis*. 1977;135(3):392-39.
- Arora S, Jindal N, Walia G, Gulati VL. Presumptive identification and antibiotic susceptibility of Group B streptococci. *Indian Journal of Pathology & microbiology*, 1994;37(2):185-90.