

STABILITY AND OPTIMAL LIPOLYTIC ACTIVITY OF SONICATED *PRUNUS AVIUM* LIPASE AND ITS APPLICATION AS FAT DEGRADATION AND HYDROLYSIS OF POULTRY WASTEWATER

Hifza Akhtar¹, Muhammad Ilyas Tariq², Nasrgis Sultana³ & Ambreen Imtiaz⁴

^{1,2}Pakistan Council of Scientific and Industrial Research, Pakistan, South Asia

³Department of Chemistry, University of Sargodha, Sargodha, Pakistan, South Asia

⁴Department of Biological Sciences, Forman Christian College (A Chartered University), Lahore, Pakistan, South Asia

ABSTRACT

Present work is an optimized lipolytic activity of sonicated defatted *Prunus Avium* seed lipases. The characterization and the stability of *P. avium* lipase were assayed at incremental steps of pH and temperature. Comparison of fat liberation potential of the Lipases treated and untreated poultry wastewater and mutton fat showed the importance of *Avium* lipase towards fat hydrolysis. A 300W sonicated waves in 30 minutes made meal clear for enzyme extraction by removing all interference of seeds fat, evidenced by the images of scanning electron microscope. 10 % olive oil emulsion ; a substrate, one-hour incubation time and 4000rpm agitation speed were the optimal variables for extraction of enzyme amongst the 9 runs of experiments suggested by orthogonal L9 array design. Substrate concentration was found the actual contributor to facilitating the lipase for complete hydrolysis. A 43.19% contribution (ANOVA result) of olive oil: substrate was found responsible for maximum lipolytic activity (1.52 μ U) of *Avium* lipase at pH 6 and 40^oC. Best stability of lipase was observed in n-heptane. A release of 6 times more poultry wastewater fat as compared to acid hydrolysis was the successful application of *P. avium* after 180minutes of the incubation period. Degradation of lipids as increase the %age of monounsaturated acid as compared to polyunsaturated acid (GLC analysis) speculating that *P. avium* enzyme can be utilized in oil/ fat, detergent and wastewater industry due to fat degradation property.

KEYWORDS: Hydrolysis, Lipolytic Activity, Orthogonal Array, *Prunus Avium*, Scanning Electron Microscope, N-Heptane, Sonic Waves

Article History

Received: 28 Sep 2018 | Revised: 07 Nov 2018 | Accepted: 16 Nov 2018

RUNNING TITLE

*Lipolytic Activity and the Important Application of *Prunus Avium* Lipase*

INTRODUCTION

Lipase is the enzyme known as triglycerol hydrolases. The subsequent ability to release the glycerols and fatty acids make lipase participation vital in worldwide enzyme industry market. Moreover, being regioselective, chemoselective or enantioselective, the lipases are becoming more and more important for special catalytic reactions in bio-technological controlled manufacturing processes [1]. Versatility for reactions like hydrolysis, esterification, interesterification is another reason for lipase acceptability in food, detergent, cosmetic, pharmaceutical, leather, lubricant,

polyester industries [2]. Most recent participation of lipases in the production of bio-modified fats is the unique characteristic of this enzyme which opens the new era of value addition in already existed fatty acids based products to meet the nutritional and safety requirements of the current global dietary rules [3].

Lipases fate for its suitability and acceptability in different processes is very much dependent on their sources. Presently, available lipases main sources are gastric, hepatic and pancreatic parts of the animal, bacterial and fungal microorganisms and plants seeds [4]. The selection of the source dependent fully on their catalytic properties towards different reactions like esterification, interesterification, and transesterification including acidolysis and alcoholysis in industrial application [5, 6]. Amongst all plant seed lipases source has been focusing the attention due to its low cost, huge and diversity of sources (plants), easy handling and simple purification methods along with the specificity for the targeted reactions [7]. Nevertheless, the heat independency characteristic of lipase activity makes it more important for oil chemical industries to cut down their energy expenses and the degradation of compounds as compared to chemical processes. Literature evidenced the advantages of seed lipases over the animal and microbial lipases for hydrolysis of oils and fats and its application in food, oil chemicals, detergent, and fine chemical industries [3]. Recently, increasing interest has been observed of these enzymes in wastewater treatment to reduce sludge flotation after reducing fat and fatty matter concentration which enable smooth aerobic and anaerobic biological treatments [8]. The plant source lipase would make the wastewater treatment feasible as compared to high-cost lipases preparation of microbial and animal source.

Oil-bearing seeds worldwide used as the source of oil as well as for the production of biodiesel with the representation of serious industrial waste environmental concern of the defatted seeds (meal). Previous researchers suggested the lipases preparation from different nonconventional plants but no such study was reported on the *Prunus* species cultivated tremendously almost all the region of the world. As *Prunus* species demonstrated good fruit nutritive attributes as well as other values: physicochemical, fatty acid profile, antioxidant and antimicrobial so their seeds were chosen for the preparation of lipase enzymes. Ongoing study based on the extraction of enzymes by means of low heat extraction technique –ultrasonic assisted extraction for enzymatic activity (lipolytic) of *Prunus Avium* indigenous to Pakistan. Species under investigation has grown in the northern area of Gilgit and Baltistan. As the total production consumed in the home market, so ample raw material available for the industries for the extraction of the enzymes. Multivariate parameters used to achieve the optimal conditions for ideal lipolytic action with the tolerance power of this enzymes in different solvents.

Characterization of *P. avium* enzyme, its ability to hydrolyze olive oil and the application of extracted enzyme as fat degradation of meat and poultry wastewater is the part of the study. The findings will help to produce a vital industrial product – enzymes with optimal conditions which can be important for making the industrial procedures feasible. The other fruitful outcome will be the utilization of agro-waste which is the main issue of food processing units.

MATERIALS AND METHODS

Collection and Defattification of Seeds: The fruits of *Prunus Avium* were collected from the Karim Abad region of Gilgit-Baltistan- Pakistan. Seeds were collected manually from the fruits grown in garden and orchids. Decortication was done with a 20% salt solution [9]. Collected seed kernel was kept in the dry forced oven (Memmert Germany) at 50⁰ C for six hours. They were ground into fine powder. Ultrasonic assisted extraction employed for complete defattification of seeds. Hielscher ultrasonic UP 400 S used for sonication. A lipid was extracted by opting conditions after optimization of different parameters like solvent and seed ratio (S/S ml/mg), sonic power and solvents used. The conditions used for seed

defatting for the present study was 300W sonic power, 10ml/gm s/s ratio and a mixture of chloroform and methanol (2:1v/v) were used as the solvent. Extraction time was 30 minutes and the temperature was 50°C [10]. The images of the *Prunus avium* seeds after oil extraction were scanned with Scanning electron microscopic at 25.0kVx1.00k magnification factor.

Optimal Extraction of Enzymes by Taguchi Method/Orthogonal Array (Design of Experiment): Nine experiments were designed by using the advanced statistical technique (L9 orthogonal array of Taguchi method) for optimal extraction of lipase from *Prunus Avium*. The experimental runs based on three variable parameters with three levels namely substrate concentration (10, 20 and 30 %), incubation time (30, 40 and 60 minutes) and agitation speed (2000, 3000 and 4000 rpm) as mentioned in table-1. For each experiment 20gm, defatted seeds of *Prunus Avium* were suspended separately in 0.1M citrate buffer at pH 6.0. The mixtures were shaken in an orbital shaker for an hour at 45°C. The extracts centrifuged with different agitation speed and time based on the statistical model. The supernatant was diluted further with 1:1 v/v ratio of distilled water and 0.1M citrate buffer. The extracts containing lipases were stored in a refrigerator for enzymes assays [11].

Enzymes Assays: The spectrophotometric method was used for the determination of lipase activity. Olive oil emulsion of different concentration (10-30%) was used as the substrate for lipase activity. Assay mixture containing 5ml substrates emulsions, 5ml chloroform, hexane (1:1v/v) mixture, 2.5ml Cu-TEA (Copper-Triethanolamine) reagent. Mixtures were centrifuged for 30-60 minutes at 2000-4000 rpm according to experimental conditions (optimization process). An absorbance of supernatants of all the individual experiments was measured at 440nm after adding 0.5ml of 0.15% sodium diethyldithiocarbamate. Stearic acid fractions used as the reference material for enzyme assay.

LIPOLYTIC ACTIVITY EVALUATION

Effect of pH and Temperature: Optimum pH for the lipolytic activities of lipases was determined by using phosphate and tris buffer covering the range of pH between 3.0-9.0. For optimum temperature absorbance were measured after every 5° rise of temperature from 30 to 70°C while adopting the same method mentioned above in enzyme assay [1].

Enzyme Stability in the Presence of Different Solvents: The stability of lipase was studied in the presence of di-isopropyl ether, n-heptane, and cyclohexane by incorporated each solvent into the assay mixture and tested the stability using 10% olive oil emulsion. The pH of the assay was maintained at 6.0 and 40°C was the temperature [12].

Hydrolysis of Poultry Waste Water: Poultry wastewater was collected from a local poultry processing industry (K&N Manga Road, Pakistan). Five samples were treated with different concentrations of enzyme extracts (5-15mg/mL) added to 0.5M citrate buffer for 180 minutes at pH 6.0 and 40°C. The comparison was made with acid hydrolysis of the same waste -water.

Degradation of Mutton Fat: Mutton pieces were collected contained fat particles from the local slaughterhouse (Baker mandi, Pakistan). Dissolved the 5gm, piece in enzyme extract (40mg/mL) and incubated at 40°C for 72 hours. The pH of the mixture was maintained at 6.0. The change in the particle size measured as the decrease of particle mass. The methyl ester of the enzyme-treated as well as control (without treatment) was prepared by using BF₃ – methanol as mentioned in the previous work [13].

Statistical Analysis: All the experiments were conducted in triplicate. Signal to noise ratio (SNR) was used to determine the maximum extraction of enzymes from the *Prunus Avium* by considering larger the better (LTB) formula for

each experiment (Table2)

Larger the better– $SNR_i = -10 \log_{10} \frac{1}{n} \sum_{j=1}^n \frac{1}{y_j^2}$ (1) Whereas, analysis of variance (ANOVA) used to determine the %age parameter contribution towards the optimal extraction of enzymes (Table3) with the help of below mentioned formula

$$\% \text{age contribution of a factor} = \frac{SSf}{SST} \times 100 \quad (2)$$

RESULTS AND DISCUSSIONS

Extraction of Oils: Plants seeds are not only the main source of oils, protein, carbohydrates, and minerals but also produce enzymes utilized by the developing embryo during the storage time in the form of fatty acid [14]. Enzymes activity indicated by the hydrolysis of glycerides and experimentally evidenced by the liberation of the free fatty acids. The concentration of these free fatty acids attributed the lipolytic activities of lipase in the seeds bearing polar and neutral lipids. The amount of extracted oil by UAE from seeds of *Prunus Avium* were 42.75 %. The reason for UAE selection was the previous research work which supported that ultrasonic-assisted extraction (UAE) was a highly efficient method for seeds oil extraction [15]. Secondly, a technique was ideal for isolation the seeds compounds present in minute quantities and sensitive to heat [16]. The present study focused on the seeds which itself contain rich amount of oil and free fatty acid so, defatification was done prior to the enzymatic assay. Scanning electron microscope (SEM) images at magnification power 25.0kV x 1.00k ensured the complete removal of lipid and free fatty acids of ultrasonically treated *P. avium* seeds (figure-1).



Figure 1: Scanning Electron Microscopic Image of Sonicated Defatted Prunus Avium Seeds

Extraction of Enzymes: Incubation time, temperature, pH, nature of buffer, agitation speed (rpm) were found the influential parameters of enzymes assays from previously reported research findings. As the amount, conditions and index of pH vary from seeds to seeds so, optimization of the process was essential to achieve the target of the present research work. Nine designed experiments were run with incubation time, the concentration of the substrate and agitation speed (variable parameters) along with their levels (low to high values). The concentration of free fatty acids showed the action the performance of the enzymes as these enzymes have been recognized as the hydrolysis promoters [17]. A concentration of liberated fatty acid (μL) as a result of hydrolysis of fats of each experiment (in triplicate). The selection of the Taguchi method for the present study was on the basis of this model version for investigating the production problems and the loss function [18]. Loss of function calculated the differences between experimental and predicted values and this was only possible to calculate the signal to noise ratio (SNR). The SNR values were the probable outcome served as the objective of

¹l, j and n stands for experiment no, trial no and no of experiments

this optimization study. The variable parameters, the concentration of liberated fatty acids, their mean values, SNR and SNRT were mentioned in table-1 by successfully adopting L9 orthogonal array with 9 runs (Taguchi model).

Table 1: L9 Orthogonal Array Design (Taguchi Method) for Concentration of Free Fatty Acids (µl) and SNRs due to Variable Parameters

Experiments	Concentration of Substrate(A)	Incubation Temperature (B)	Agitation Speed (C)	Concentration of Free Fatty Acid(µl)			Mean Values of Fatty Acids (µl)	SNR	SN RT
				1	2	3			
1	10	30	2000	110	108	112	110	40.87	44.3
2	10	45	3000	195	203	202	200	46.0	
3	10	60	4000	277	268	265	270	48.62	
4	20	30	2000	171	178	191	180	45.08	
5	20	45	3000	203	202	200	202	46.09	
6	20	60	4000	238	234	231	234	47.4	
7	30	30	2000	106	108	110	108	40.66	
8	30	60	3000	121	117	123	120	41.60	
9	30	60	4000	132	125	138	132	42.36	

To achieve an optimal combination of the variable parameter with an effective level, signal to noise level-based values (SNRL) calculated. SNRL values of each experimental variable for each of its specified level separately mentioned in table-2.

Table 2: Level Mean SNRs of Variable Parameters for Specific Level

Parameter ID	Parameter Description	Levels		
		1	2	3
A	Concentration of substrate	45.16	46.19	41.54
B	Incubation time	42.20	44.56	46.13
C	Agitation speed	42.17	44.56	46.13

The SNRL values of each independent variable (A, B, C) at their levels (1, 2, 3) showed the direct influence on the concentration of fatty acids (CoFA). Bigvalue of SNRL greater the influence of that variable of that level to perform the hydrolysis activity of enzymes on the substrate. Analysis of variance (ANOVA) applied to determine the significant contribution towards the maximum CoFAat optimal levels represented in table-3. The values obtained after taking the sum of the square of the fifth factor (SSf) and the total sum of square (SST) as calculated by earlier researchers [18].

Table 3: Parameters Contribution towards Optimal Cofa

Variable Parameters	Significant Contribution in terms of Percentage
Concentration of substrate	43.19
Incubation time	28.16
Agitation speed	28.65

The concentration of the substrate found the more significant parameter in the present study as compared to incubation time and agitation speed. The optimal concentration of the substrates was found in most of the enzymatic studies [19].

Lipase Characterization: The lipase characterization based on its action and the lipolytic activity is the attributor of lipases activity, so evaluation of the activity was made by monitoring the enzymes activities and stabilities at various temperature, pH, and solvents. Prior to taking the absorbance of liberated fatty acids of *Prunus Avium* a standard curve of different concentrations of the stearic acid plot was drawn (50-500 µL). The selection of stearic acid was made due to

stability and easily availability according to local climate and environment. Previous work reported the successful use of stearic acid [20] along with other fatty acids (oleic acid) as the control reagent for enzymatic studies [21]. Literature revealed that characterization of enzymes influenced by the external factors like pH and temperature moreover; the ripening stage, soil nature, extraction procedure, and seed origin also contributor of enzymes properties. These internal and external parameters make the extracted enzymes viable for different industries, to control the rancidity and increase the shelf life of the product [22]. Variation in pH plays the vital role in the efficiency of lipolytic activity of enzymes. The optimization of the pH for lipases is mandatory step to make the enzymes specific and viable for reaction either for catalyzed synthesis or for the formation of products through reactions (hydrolysis, acylation, esterification, and cleaving).

Different experiments were carried out with the variation of pH covering the range of acidic to alkaline 3.0-9.0. Extracted lipase hydrolyzed the substrate triglycerides (10% emulsion fraction of olive oil) measured as the concentration of liberated fatty acids (LFA) in the form of absorbance. lipase showed single peak point when measured absorbance at different pH which clearly indicated the presence of single nature enzyme, the present finding was in accordance with the previous study of lipolytic activities of extracted lipases of peanut, walnut, castor, lipin, pinus and rape seeds which showed single optimal point of pH [22]. Similarly, Variation in the temperature not only influenced the rate of lipolytic activities of lipases but can also be leading to the total denaturing of the enzymes as well. Grosh and coworkers reported the complete loss of enzymatic activities at elevated temperature [23]. The absorbance of fatty acid corresponding to the different temperature in the present study was measured at 440.0nm. The noted absorbance values at different pH and temperature are the indicators of the carboxyl ester bond cleavage of olive oil with the removal of fatty acids and their comparison with standard curve values of stearic acid showed their concentration. The lipolytic activity of lipases at different temperature and pH profiled in figure-2 (b&a) after incorporating the LFA concentration in the Guven's equation[24]




Figure 2: Effect of Ph (A) and Temperature (B) on P. Avium Lipase Activity Lipolytic Activity was Calculated by Measuring the Absorbance of Liberated Fatty Acids using 10% Emulsion of Olive Oil

The result of present work showed the optimal activity of extracted lipase of *Prunus Avium* was observed at 40^o C when assayed at pH 6.0, likewise the stability of lipolytic activities regarding pH was around 4.0-7.0 for olive oil substrate, whereas at 35-55^o C the *P. avium* lipase exhibited thermal stability which was in accordance to the previous work [25] on different vegetable oil-bearing seeds. The thermal and alkaline stability range of same origin enzymes can be varying due to the change of the substrate, incubation time and agitation speed.

The stability of enzymes is very much dependent on the solvents because the solvents can alter the pH memory, thermal stability and even substrate selectivity of the enzymes. Gupta and coworker reported inhibition of enzymes activity in different concentration of organic solvents [26]. Di-isopropyl ether, n –heptane, and cyclohexane were used to study the

stability of the *Prunus Avium* extracted enzymes on already observed optimal points of temperature and pH. All the solvents showed stability against these optimal points with a variation of their activities in term of the concentration of fatty acid liberation and their absorbance. The figure-5 depicted the stability trend of extracted lipases due to the combination of hydrophilic solvent together with a non-aqueous solvent which is in accordance with the Tsuzuki and Kitamura studies [27]




Figure 3: Surface plot of *P. Avium* Lipase Activity in Different Solvents at 40⁰ C and 6.0 Ph.

Hydrolysis of Poultry Waste Water: The wastewater of poultry industries contained organic and fatty based suspended solid fractions. The sources might be the washing steps of slaughter birds, equipment and processing plant washing [28]. The free fatty acids concentration in poultry field waste-water was analyzed before and after the lipolytic activity with different concentration of lipase extract. The maximum concentration of fatty acids was observed with 40mM lipase extract after 180 minutes of incubation time which was six times higher than the control wastewater sample (figure 4).




Figure 4: Effect of Lipase on the Liberation of Fatty Acid of Poultry Wastewater at Different Incubation Times

Polzellireported the liberation of fatty acids 7.4 times higher with enzymatic treatment than the normal acid hydrolysis [5]. The variation in the activity might be due to the variation of wastewater characteristics of different processing plants, consumption of water for each slaughtered bird and the source of enzymes. The results are important for the poultry industrial wastewater process for the conversion of triacylglycerols into fatty acids and will step forward for industry development as few studies reported to use lipase in fat wastewater.

Fat Degradation: Degradation of the fat particle in flash of animals was reported in different studies with a different concentration of enzymes and at different incubation time, temperature and pH. Using 25mL, *P. avium* enzyme for 5gm, mutton piece, 6 % mass reduction was observed by calculating the weight of the particle. The result did not match with the findings of Masses studies of 40 % reduction of beef particle size with pancreatic lipase and others with no claim of particle size change [29]. The apparent reason of drastic variations of results in case of a mass reduction up to 40% is the source change; pancreatic lipase (animal source of enzyme) which has higher activity as compared to plant source lipase. The fatty acid profile of the lipase treated and untreated mutton revealed the reason for mass reduction. Change in the composition of higher molecular weight fatty acids (linoleic and linolenic acid) might be the reason for loss of particle

size. Previous work mentioned the activity of lipase on oleic acid as well as linoleic and linolenic acids [30]. The change in the concentration of oleic acids and other fatty acids of lipase treated mutton fat and control of the present study was mentioned in table-4.

Table 4: Fatty Acid Composition of Lipase Treated and Un-Treated Mutton Fat

Fatty Acids	Fat Particle %Age (Untreated)	Fat Particle % (Lipase Treated)
Myristic acid	4.1	5.0
Palmitic acid	18.9	20.7
Palmitoleic acid	1.6	0.7
Stearic acid	30.8	31.3
Oleic acid	39.05	39.87
Linoleic acid	3.57	1.7
Linolenic acid	1.1	0.3
TSFAs	53.8	57.0
TUFAs	45.32	42.57
PUFAs	4.67	2.0
MUFAs	40.65	40.57

All the results of studied parameters showed the extracted lipase effective activity at optimal pH, temperature and its successful application in fat degradation and wastewater hydrolysis. The difference in the ranges of the lipolytic activity can be due to some disadvantages like selection of the substrate for application as shown lipase extraction, an interaction of the existed fat of plant seed, origin of lipase, hydrolysis time and the concentration of lipase.

CONCLUSIONS

In this study, the organic solvent stable lipases from *Prunus Avium* were successfully studied regarding optimal activity at different temperature and pH, along with the specificity and concentration of the substrates. Taguchi method made possible to extract the enzymes after optimizing the parameters like substrate concentration, incubation time and agitation speed. The single neutral nature enzymes with single optimal pH and temperature generated the possibility of the usefulness of this enzyme in fat particle degradation and poultry wastewater treatment.

REFERENCES

1. Gadge, P.P., Madhikar.S.D, Yewle. J.N, Jadhav. U.U, Chougale. A.D, Zambare.V.P, Padul. M.V. *Biochemical studies of lipase from germinating oil seeds (Glycine max). American Journal of Biochemistry and Biotechnology.*2011,7(3):141-5.
2. Cancino, M., Bauchar.t P, Sandoval. G, Nicaud.J.M, André.I, Dossat. V, Marty.A. A. variant of *Yarrowialipolytica* lipase with improved activity and enantioselectivity for resolution of 2-bromo-arylacetic acid esters. *Tetrahedron: Asymmetry.* 2008,19(13):1608-12.
3. Neklyudo.v, A.D, Berdutin.a. A.V, Ivankin. A.N, Karpo.B.S. Kinetic characterization of enzymatic hydrolysis of complex protein substrates for producing nutrient media. *Applied Biochemistry and Microbiology.* 2002,38(4):328-34.
4. Barros, M., Fleuri.L.F, Macedo. G.A. Seed lipases: sources, applications and properties-a review. *Brazilian Journal of Chemical Engineering.* 2010,27(1):15-29.

5. Zhong, Q., Glatz, C.E. Enzymatic assay method for evaluating the lipase activity in complex extracts from transgenic corn seed. *Journal of agricultural and food chemistry*. **2006**, 54(9):3181-5.
6. Darvishi, F., Nahvi, I, Zarkesh-Esfahani, H, Momenbeik, F. Effect of plant oils upon lipase and citric acid production in *Yarrowialipolytica* yeast. *BioMed Research International*. **2009**, 8;2009.
7. Polizelli, P.P., Facchini, F.D, Bonilla-Rodriguez, G.O. Stability of a Lipase extracted from seeds of *Pachira aquatica* in commercial detergents and application tests in poultry wastewater pretreatment and fat particle hydrolysis. *Enzyme research*. **2013**;2013..
8. Cammarota, M.C., Freire, D.M. A review on hydrolytic enzymes in the treatment of wastewater with high oil and grease content. *Bioresource technology*. **2006**, 97(17):2195-210.
9. Gupta, A., Sharma, P.C, Tilakratne, B.M, Verma, A.K. Studies on physico-chemical characteristics and fatty acid composition of wild apricot (*Prunus armeniaca* Linn.) kernel oil.
10. Tian, Y., Xu, Z, Zheng, B, Lo, Y.M. Optimization of ultrasonic-assisted extraction of pomegranate (*Punicagranatum* L.) seed oil. *Ultrasonics Sonochemistry*. **2013**, 20(1):202-8.
11. Hifza, A., Tariq, M.I, Sultana, N, Akhtar, T, Nawaz, S. Effects of different parameters on the lipolytic activities of lipase and phospholipase extracted from *Prunus arminica* meal of hunza cultivar. *Pakistan Journal of Science*. **2017**, 69(2):168-172.
12. Akhtar, H., Hamid, S, Khan, J.I. Activity of extracted lipase and phospholipase of (Perlette) Grape cultivar seed meal. *Journal Chemical Society of Pakistan*. **2007**, 29(2):167-170
13. Hifza, A., Tariq, M.I, Sultana, N, Nosheen, S, Habib, F, Shafiq, M. extraction and bioanalysis of ultrasonic assisted pakistani cultivar *Prunus dulcis* seed: an optimization study. *Pakistan Journal of Science*. **2018**, 70(1):1-7.
14. Abigor, R.D., Uadia, P.O, Foglia, T.A, Haas, M.J, Scott, K, Savary, B.J. Partial purification and properties of lipase from germinating seeds of *Jatropha curcas* L. *Journal of the American Oil Chemists' Society*. **2002**, 79(11):1123-6.
15. Cravotto, G., Bicchi, C, Mantegna, S, Binello, A, Tomao, V, Chemat, F. Extraction of kiwi seed oil: Soxhlet versus four different non-conventional techniques. *Natural product research*. **2011**, 25(10):974-81.
16. Jerman, Klen, T., Mozetič, Vodopivec, B. Ultrasonic extraction of phenols from olive mill wastewater: comparison with conventional methods. *Journal of agricultural and food chemistry*. **2011**, (24):12725-31.
17. Nelofe, R., Ramanan, R.N, Rahman, R.N, Basri, M, Ariff, A.B. Sequential optimization of production of a thermostable and organic solvent tolerant lipase by recombinant *Escherichia coli*. *Annals of microbiology*. **2011**, 61(3):535-44.
18. Kumar, R.S., Sureshkumar, K, Velraj, R. Optimization of biodiesel production from *Manilkara zapota* (L.) seed oil using Taguchi method. *Fuel*. **2015**, 15;140:90-6.
19. Jamie, A., Alshami, A.S, Maliabari, Z.O, Ateih, M.A. Development and validation of a kinetic model for enzymatic hydrolysis using *Candida rugosa* lipase. *J. Biotechnol. Bioprocess*. **2017**, 7:297.

20. Waheed, A., Mahmood.S and Saleem.M. Activity of extracted lipase and phospholipase of tobaccum seed meal. *Natural products science.***2001**,7:17.
21. Eze,S.O.O.andChilaka.F.C. (2). Lipolytic activity of some species of germinating cucubitaceae world. *Journal Agricultur Science.***2012**,6: 700-706.
22. Seyhan,.F., Tijskens.L.M.M and Evranuz.O. Modelling temperature and pH dependence of lipase and peroxidase activity in turkishhazelnuts.*Journal of food engineering.***2002**,52:387-395.
23. Laskawy, G., Senser. F, Grosch. W. Storage stability of roasted hazelnuts. CCB, Review for Chocolate, Confectionery and Bakery. **1983**.
24. Güven, K.C., Berğişadi. N, Güler. E. A Modification of Duncombe's Method and its Application to the Lipolytic Activity Assay of Heparin. *Fette, Seifen, Anstrichmittel.* **1979**;81(4):152-4.
25. Altaf,A., Ankers.T.V, Kaderbhai. N, Mercer. E.I, Kaderbhai. M.A. Acid lipase of castor bean lipid bodies: Isolation and characterisation. *Journal of plant Biochemistry and Biotechnology.* **1997**,6(1):13-8.
26. Gupta, M.,Batra.R.,Tyagi.R.,Sharma. A..Polarity index: the guiding solvent parameter for enzyme stability in aqueous organic cosolvent mixture.*Biotechnol.Prog.***1997**,13:284-288.
27. Tsuzuki, W., Ue.A, Kitamura.Y. Effect of dimethylsulfoxide on hydrolysis of lipase. *Bioscience, biotechnology and biochemistry.***2001**,65(9):2078-82.
28. Valladão,A.B., Torres. A.G, Freire. D.M, Cammarota.M.C. Profiles of fatty acids and triacylglycerols and their influence on the anaerobic biodegradability of effluents from poultry slaughterhouse. *BioresourceTechnology.* **2011**, 102(14):7043-50.
29. Masse, L., Massé. D.I, Kennedy. K.J. Effect of hydrolysis pretreatment on fat degradation during anaerobic digestion of slaughterhouse wastewater. *Process Biochemistry.* **2003**,38(9):1365-72.
30. Polizelli, P.P., Facchini. F.D, Cabral.H, Bonilla-Rodriguez.G.O. A new lipase isolated from oleaginous seeds from *Pachiraaquatica* (Bombacaceae). *Applied biochemistry and biotechnology.* **2008**150(3):233-42.