

Centella asiatica increases B-cell lymphoma 2 expression in rat prefrontal cortex

Kuswati*, Djoko Prakosa**, Brian Wasita***, and Nanang Wiyono****

ABSTRACT

*Department of Anatomy,
Faculty of Medicine,
Islamic University of Indonesia
**Department of Anatomy,
Embryology and Anthropology,
Faculty of Medicine
Gadjah Mada University.

***Department of Pathological
Anatomy,
Faculty of Medicine,
Sebelas Maret University
****Department of Anatomy,
Faculty of Medicine,
Sebelas Maret University

Correspondence

dr. Kuswati, M.Sc
Department of Anatomy,
Faculty of Medicine,
Islamic University of Indonesia
Yogyakarta 55281
Phone: +6285228478487
Email : kuswati2005@yahoo.com

Univ Med 2015;34:10-6

DOI: 10.18051/UnivMed.2015.v34.10-16

BACKGROUND

Stress is one of the factors that cause apoptosis in neuronal cells. *Centella asiatica* has a neuroprotective effect that can inhibit apoptosis. This study aimed to examine the effect of *Centella asiatica* ethanol extract on B-cell lymphoma 2 (Bcl-2) protein expression in the prefrontal cortex of rats.

METHODS

An experimental study was conducted on 34 brain tissue samples from male *Sprague Dawley* rats exposed to chronic restraint stress for 21 days. The samples were taken from following groups: non-stress group K, negative control group P1 (stress + arabic gum powder), P2 (stress + *C.asiatica* at 150 mg/kgBW), P3 (stress + *C.asiatica* at 300 mg/kg BW), P4 (stress + *C.asiatica* at 600 mg/kg body weight) and positive control group P5 (stress + fluoxetine at 10 mg/kgBW). The samples were made into sections that were stained immunohistochemically using Bcl-2 antibody to determine the percentage of cells expressing Bcl-2. Data were analyzed using one way ANOVA test followed by a post - hoc test.

RESULTS

There were significant differences in mean Bcl-2 expression between the groups receiving *Centella asiatica* compared with the non-stress group and stress-only group (negative control group) ($p < 0.05$). The results were comparable to those of the fluoxetine treatment group.

CONCLUSION

The *Centella asiatica* ethanol extract was able to increase Bcl-2 expression in the prefrontal cortex of *Sprague Dawley* rats exposed to restraint stress. This study suggests that *Centella asiatica* may be useful in the treatment of cerebral stress.

Keywords : *Centella asiatica*, B-cell lymphoma 2, prefrontal cortex, stress, rat

***Centella asiatica* meningkatkan ekspresi protein B-cell lymphoma 2 pada korteks prefrontalis tikus**

ABSTRAK

LATAR BELAKANG

Stres merupakan salah satu faktor yang dapat menyebabkan apoptosis pada sel saraf. *Centella asiatica* memiliki efek neuroprotektif yang dapat menghambat apoptosis. Penelitian ini bertujuan untuk menilai peningkatan ekspresi protein B-cell lymphoma 2 (Bcl-2) setelah pemberian ekstrak etanol *Centella asiatica* pada tikus yang diberi perlakuan stres restraint kronik.

METODE

Sebuah penelitian eksperimental dilakukan pada 34 jaringan otak tikus Sprague Dawley yang diberi perlakuan stres restraint kronik selama 21 hari. Jaringan otak tikus berasal dari 6 kelompok berikut: kelompok non stres (K), kelompok kontrol negatif stres + pulvis gummy arabicum (P1), kelompok stres + *C. asiatica* dosis 150 mg/kgBB (P2), kelompok stres + *C. asiatica* dosis 300 mg/kgBB (P3), kelompok stres + *C. asiatica* dosis 600 mg/kgBB (P4) dan kelompok stres + fluoksetin dosis 10 mg/kgBB sebagai kontrol positif (P5). Jaringan otak tersebut dibuat preparat yang dicat secara imunohistokimia untuk penghitungan prosentase sel yang mengekspresikan Bcl-2. Data yang diperoleh dianalisis menggunakan uji one way ANOVA dilanjutkan analisis post-hoc test.

HASIL

Terdapat perbedaan bermakna pada rerata prosentase ekspresi Bcl-2 antara kelompok yang mendapat terapi *Centella asiatica* dan fluoksetin dibandingkan dengan stres ($p < 0,05$). Ekspresi Bcl-2 pada kelompok yang mendapat terapi *Centella asiatica* dan kelompok fluoksetin lebih tinggi dibanding kelompok stres.

KESIMPULAN

Ekstrak etanol *Centella asiatica* dapat meningkatkan ekspresi protein antiapoptosis Bcl-2 di cortex prefrontalis tikus Sprague Dawley yang diberi perlakuan stres restraint kronik. *Centella asiatica* dapat berperan sebagai pengobatan pada restraint stress yang dialami tikus.

Kata Kunci : *Centella asiatica*, B-cell lymphoma 2, cortex prefrontalis, stres, tikus

INTRODUCTION

Stress can cause disorders of cell homeostasis, because it increases the secretion of glucocorticoids. In the central nervous system, an increase in glucocorticoids will trigger release of glutamate. The latter binds to N-methyl-D-aspartate (NMDA) receptors, resulting in the opening of calcium channels, with subsequent calcium entry into the cytoplasm and increase in cytoplasmic free calcium. The entry of calcium and water into the mitochondria results in mitochondrial edema, impaired balance between

Bcl-2-associated X protein (Bax) and B-cell lymphoma 2 (Bcl-2) expression, increase in mitochondrial membrane permeability and release of cytochrome c, triggering apoptosis.⁽¹⁾

Stress affects several areas of the prefrontal cortex, hippocampus, amygdala and brainstem. In the prefrontal cortex, stress results in a decrease in the activity of neurons, a reduction in the number of dendrites and shortening of dendrites in the medial prefrontal cortex.⁽²⁾ Chronic stress results in retraction of the apical dendrites of pyramidal cells in layer II / III and V of the medial prefrontal cortex.⁽³⁾ Therefore,

individuals who experience stress should be given therapy to prevent apoptosis and morphological changes in the prefrontal cortex.

Drugs that can be used to inhibit apoptosis in the central nervous system caused by stress belong to the class of antidepressant drugs. Olanzapine, imipramine, amitriptyline, citalopram and fluoxetine can inhibit apoptosis by increasing the expression of Bcl-2 and B-cell lymphoma-extra large (Bcl-xl) and decreasing the expression of Bax.^(4,5) Although they may inhibit apoptosis, antidepressant drugs have many side effects. Side effects of tricyclic antidepressant drugs include dry mouth, dizziness, decreased vision, constipation, sedation, orthostatic hypotension and tachycardia. One class of antidepressant drug, the selective serotonin reuptake inhibitors (SSRIs), have side effects such as nausea, sleep disturbances, sexual dysfunction, changes in appetite, headache and dry mouth.⁽⁶⁾ Therefore, it is necessary to find an alternative drug to inhibit apoptosis, one of which can be found in the class of herbs such as pegagan (*Centella asiatica*).

Centella asiatica has a wide therapeutic dose range. *Centella asiatica* extract at a dose of 100-1000 mg/kg body weight did not cause any side effects, and had a median lethal dose (LD50) of more than 4000 mg/kg body weight. *Centella asiatica* extract up to a dose of 4000 mg/kg did not result in impaired liver and kidney function as well as changes in the macroscopic picture of the liver, lung and kidney.⁽⁷⁾ *Centella asiatica* contains madecassoside, asiaticoside, asiatic acid and madecassic acid which have antioxidant and neuroprotectant effects.⁽⁸⁾ Administration of *C. asiatica* extract in mice with stress can improve spatial memory, increase hippocampal *Cornu Ammonis* (CA) 1 pyramidal layer thickness, and increase the number of neuroglia in the hippocampus.^(9,10) The objective of this study was to evaluate the anti-apoptotic effects of *C. asiatica* ethanol extract, in particular on the expression of Bcl-2 in the prefrontal cortex of rats exposed to chronic restraint stress.

METHODS

Research design

This study is part of an umbrella research using an experimental design to evaluate the effect of *Centella asiatica* ethanol extract on memory, apoptosis and blood biochemistry (corticosteroid hormone levels and interleukins). The present study was limited to the examination of Bcl-2 protein expression in the prefrontal cortex of Sprague Dawley rats. The study period was from August 2012 until September 2013.

Research subjects

The sample size was determined from a mean difference in Bcl-2 expression between groups of 9.0%, with $\sigma=0.05$ and $\beta=0.2$, yielding 6 subjects per group. Therefore the total sample size for 6 groups was 36.⁽¹¹⁾ The subjects were 34 brain tissue samples from male Sprague Dawley rats exposed to chronic restraint stress for 21 days. The samples were taken from 6 experimental groups consisting of a non-stress group (K), a negative control group exposed to stress + arabic gum powder (P1), 3 groups exposed to stress + *C. asiatica* at dosages of 150 mg/kg body weight (P2), 300 mg/kg body weight (P3), and 600 mg/kg body weight (P4), respectively, and the positive control group exposed to stress + fluoxetine at a dosage of 10 mg/kg body weight (P5). In the present study chronic restraint stress was used, by placing the rats into plastic transparent acrylic restraint tubes of 15 cm length and 5.5 cm diameter, with a number of 3 mm diameter holes at their closed ends. This treatment caused the rats to be immobilized, resulting in psychological stress.

Starting from day 1, the stress groups were exposed to restraint stress by placing them into the restraint tubes daily for 6 hours, from 09.00 to 15.00, for 21 days. In the stress groups receiving *C. asiatica* or fluoxetine, the treatments were given by gavage at predetermined dosages, 30 minutes before being stressed. The non-stress and stress control groups were given arabic gum powder for 21 days. On day 23 the rats were

terminated and perfused, followed by taking brain tissue samples. Subsequently the samples were made into paraffin blocks, sliced on a microtome, and stained immunohistochemically with Bcl-2 antibody.

Preparation of *Centella asiatica* extract

C. asiatica was obtained in simplicia form from CV. Merapi Farma Herbal (a commercial herbal manufacturer). *C. asiatica* identification was performed in the Plant Systematics Laboratory, Faculty of Biology, Gadjah Mada University. The *Centella asiatica* extract was standardized nonspecifically by examination of its water content and specifically by determining its asiaticoside content by thin layer chromatography. The *C. asiatica* extract contained $4.16 \pm 0.51\%$ asiaticoside. The *C. asiatica* extract was diluted to give three concentrations of 30 mg/ml, 60 mg/ml and 120 mg/ml, respectively. These concentrations were used to prepare the three treatment dosages of 150 mg/kg, 300 mg/kg body weight and 600 mg/kg, respectively. Dilution of the extract was performed once in three days at the Laboratory of Pharmacology, Medical Faculty, Gadjah Mada University. The dilutions were stored in the refrigerator. To calculate the volume of the *C. asiatica* extract for each of the rats, the following formula was used:

$$\text{Volume} = \text{dose (mg/kgBW)} \times \text{weight (kg)} / \text{concentration (mg/mL)}.$$

Immunohistochemical staining

Paraffin blocks were made from the brain tissue and then cut into 4 μm -thick sections. The cut brain areas comprised the prelimbic, infralimbic and cingulate cortex at Bregma 3.72 mm to 2.52 mm.⁽¹²⁾ Five sections were made from each rat brain at 50 μm spacings. The first step in immunohistochemical staining consisted of deparaffinizing the sections using xylol and descending alcohol series. The tissue sections were then incubated in 3% H_2O_2 in 10% methanol for 20 minutes and rinsed 3 times with phosphate buffered saline (PBS).

The tissue sections were blocked by Background Sniper blocking reagent for 10 minutes. Without rinsing, the tissue sections were incubated overnight at 4^o C with anti-Bcl-2 primary antibody (1:400) (Bioworld No. BS1511). The following day, the sections were rinsed using PBS 2 times, 0.2% Tween 2 times and PBS 2 times. The sections were incubated in Trekki Universal Link, followed by incubation with horseradish peroxidase-conjugated streptavidin for 10 minutes. The tissue sections were again rinsed using PBS 2 times, 0.2% Tween 2 times and PBS 2 times. The final color product for single labeling was typically visualized by immersion in 3,3'-diaminobenzidin (1:100) solution for 5 minutes, rinsed 5 times with distilled water, counterstained with Meyer's hematoxylin for 1 minute, and rinsed with tap water for 2 minutes. The sections were then dehydrated in an ascending alcohol series (70%, 80%, 90%, 95% and 100%) for 1 minute each, cleared with xylene, and then covered with canada balsam and cover slip.

Measurements

For each slide, cell counts were performed on 500 or more cells. The percentage of cells expressing Bcl-2 was calculated as the number of cells expressing Bcl-2 divided by the total number of cells multiplied by 100%.

Ethical clearance

This study has received ethical clearance from the Ethics Committee, Faculty of Medicine, Gadjah Mada University, under number KE/FK/399/EC.

Statistical analysis

Tests of normality showed that the percentage data were normally distributed, therefore one-way ANOVA was performed to compare the mean number of cells between groups, followed by a post-hoc test to determine differences between specific groups. The critical level for rejection of the null hypothesis was considered to be a p value of 0.05.

Table 1. Mean Bcl-2 expression by treatment groups

	Treatment groups					P	
	K (n=5)	P1 (n=6)	P2 (n=6)	P3 (n=6)	P4 (n=6)		P5 (n=5)
Bcl-2 expression (%)	1.41 ± 0.32	5.00 ± 1.12	24.89 ± 5.11	21.91 ± 4.52	25.31 ± 5.84	21.84 ± 4.37	0.000

K : control; P1: stress only; P2: stress + *C. asiatica* dosage of 150 mg/kg body weight; P3: stress + *C. asiatica* dosage of 300 mg/kg body weight; P4: stress + *C. asiatica* dosage of 600 mg/kg body weight; P5: stress + fluoxetine dosage of 10 mg/kg body weight

RESULTS

A total of 34 rats were analyzed, because two rats died during the study. The mean expression of the anti-apoptotic protein Bcl-2 in the non-stress group was $1.41 \pm 0.32\%$. Mean Bcl-2 expression in this group was the lowest compared to the other groups. Mean Bcl-2 expression in the stress-only group (P1) was $5.00 \pm 1.12\%$, which was higher than that in the non-stress group, but lower than in the groups receiving therapy (group P2, P3, P4, P5). Groups P2, P3, and P4, receiving *C. asiatica* ethanol extract at a dose of 150 mg, 300 mg, and 600 mg, respectively, had mean Bcl-2 expression values of $24.89 \pm 5.11\%$, $21.91 \pm 4.52\%$ and $25.31 \pm 5.84\%$, respectively. The mean expression of Bcl-2 in the group receiving fluoxetine (P5) was $21.84 \pm 4.37\%$. Thus, the groups that received *C. asiatica* ethanol extract and fluoxetine showed higher Bcl-2 expression than the non-stress group K and the stress-only group P1. The results of the one-way ANOVA test showed significant differences in the expression of Bcl-2. To find out between which groups there was a significant difference, follow-up analysis using a post-hoc test was performed. The latter showed significant differences between the stress-only group P1 on the one hand, and the groups treated with *C. asiatica* ethanol extract and fluoxetine (P2, P3, P4, P5) on the other hand, with p-values of 0.000 to 0.002 ($p < 0.05$) (Table 1).

Results of multiple comparisons analysis showed that the control group (K) was significantly different from the groups exposed to respectively stress + *C. asiatica* dosage of

150 mg/kg body weight (P2), stress + *C. asiatica* dosage of 300 mg/kg body weight (P3), stress + *C. asiatica* dosage of 600 mg/kg body weight (P4) and stress + fluoxetine dosage of 10 mg/kg body weight (P5). Group P1 was significantly different from the groups exposed to respectively stress + *C. asiatica* dosage of 150 mg/kg body weight (P2), stress + *C. asiatica* dosage of 300 mg/kg body weight (P3), stress + *C. asiatica* dosage of 600 mg/kg body weight (P4) and stress + fluoxetine dosage of 10 mg/kg body weight (P5). There were no significant differences between the groups exposed to stress + *C. asiatica* dosage of 150 mg/kg body weight (P2), stress + *C. asiatica* dosage of 300 mg/kg body weight (P3), stress + *C. asiatica* dosage of 600 mg/kg body weight (P4) and stress + fluoxetine dosage of 10 mg/kg body weight (P5) (Table 2).

Table 2. Post-hoc analysis of Bcl-2 expression percentages between groups

	95% significance index		p
	Lower bound	Upper bound	
K vs P1	-18.32	6.63	0.483
K vs P2	-33.70	-13.26	0.001*
K vs P3	-30.73	-10.28	0.001*
K vs P4	-34.13	-13.68	0.001*
K vs P5	-31.11	-9.76	0.001*
P1 vs P2	-29.63	-10.14	0.002*
P1 vs P3	-26.66	-7.16	0.001*
P1 vs P4	-20.06	-10.56	0.004*
P1 vs P5	-27.06	-6.62	0.001*
P2 vs P3	-6.77	12.72	0.547
P2 vs P4	-10.17	9.32	0.932
P2 vs P5	-7.18	13.27	0.556
P3 vs P4	-13.15	6.35	0.483
P3 vs P5	-10.15	10.30	0.994
P4 vs P5	-6.75	13.70	0.496

*The mean difference is significant at the 0.05 level

DISCUSSION

In this study, stress-exposed rats showed higher expression of Bcl-2 than did rats not exposed to stress. Differences in Bcl-2 expression between both groups were not significant. Research conducted by Kim et al.⁽¹³⁾ found that rats exposed to restraint stress, noise stress and cold stress for 10 days showed impaired memory and higher expression of Bcl-2, Bax and caspase-3 than non-stressed rats. Increased Bcl-2 expression was presumably a compensatory mechanism for neuronal protection from exposure to stress.⁽¹⁴⁾

In the present study, the expression of Bcl-2 in the stress-only group (P1) was lower than in the groups treated with *C. asiatica* and fluoxetine (group P2, P3, P4 and P5). Treatment with ethanol extract of *C. asiatica* at a dose of 150 mg/kg body weight, 300 mg/kg body weight and 600 mg/kg body weight could increase the expression of Bcl-2 protein. Dose differences did not result in significantly different expression of Bcl-2 protein. This result shows that the ethanol extract of *C. asiatica* can provide neuroprotective effects by inhibiting apoptosis through increased expression of Bcl-2 protein. This result is consistent with research conducted by Uvarajan et al.⁽¹⁵⁾ and Omar et al.⁽¹⁶⁾ who reported that the administration of asiaticoside at a dose of 50 mg/kg for 21 days can increase the expression of Bcl-2 protein and decrease the expression of Bax protein in the striatum and midbrain in rats with induced 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) neurotoxicity. In our study, there were no significant differences in the mean expression of the Bcl-2 protein between the groups that received ethanol extract of *C. asiatica* and the group on fluoxetine. Administration of fluoxetine at a dose of 10 mg/kg body weight could increase the expression of Bcl-2 protein in the prefrontal cortex of rats exposed to chronic restraint stress.

The results are consistent with research by Kosten et al.⁽¹⁷⁾ which states that the administration of fluoxetine at a dose of 5 mg/

kg for 21 days in rats exposed to repeated unpredictable stress increases Bcl-2 mRNA in the cingulate, frontal, and lateral parietal cortex. The study by Guirado et al.⁽¹⁸⁾ showed that in middle-aged rats chronic fluoxetine treatment induces changes in the expression of molecules related to neuronal structural plasticity. A previous study has clearly shown that intraperitoneal injections per se of fluoxetine in rats have effects on the structure of neurons after chronic stress.⁽¹⁹⁾

In the present study some of the rats died from gavaging trauma and some paraffin blocks were damaged due to blunted microtome knives and fatigue of the investigators. We also experienced difficulties in determining the limits of the medial prefrontal cortex, although we were guided by the atlas of the rat brain by Paxinos and Watson.⁽¹²⁾


Further researches are required to determine the expression of the Bax pro-apoptotic protein, and the Bax/Bcl-2 ratio. In addition, further researches are needed to examine other parameters such as the expression of active caspase-3 and TUNEL assay.

CONCLUSION

Ethanol extract of *C. asiatica* could increase the expression of Bcl-2 protein in the prefrontal cortex of Sprague Dawley rats exposed to chronic restraint stress. This study suggests that *Centella asiatica* may be useful in the treatment of cerebral stress.

ACKNOWLEDGMENTS

The authors would like to thank the Faculty of Medicine, Islamic University of Indonesia, Prof. Dr. Mustofa, M.Kes. Apt and dr. Dwi Cahyani Ratnasari, M.Kes. PA(K) for all advice given. We also thank Rina Susilowati M.D. Ph.D as laboratory supervisor for her support. Thanks are also due to Anggraini Janar Wulan, Rizal Adi Kusnomo, and Suparno of the Department of Physiology, Faculty of Medicine, Gadjah

Mada University, and to Sumaryati of the Department of Histology and Cellular Biology, Faculty of Medicine, Gadjah Mada University, for their technical assistance. 

REFERENCES

- Chen SD, Yang DI, Lin TK, et al. Roles of oxidative stress, apoptosis, PGC-1 α and mitochondrial biogenesis in cerebral ischemia. *Int J Mol Sci* 2011;12:7199-215.
- Wilber AA, Walker AG, Southwood CJ, et al. Chronic stress alters neural activity in medial prefrontal cortex during retrieval of extinction. *Neuroscience* 2011;174:115–31.
- Goldwater DS, Pavlides CP, Hunter RG, et al. Structural and functional alterations to rat medial prefrontal cortex following chronic restraint stress and recovery. *Neuroscience* 2009;164:798–808.
- Murray F, Hutson PH. Hippocampal Bcl-2 expression is selectively increased following chronic but not acute treatment with antidepressants, 5-HT1A or 5-HT2C/2B receptor antagonists. *Eur J Pharmacol* 2007;569: 41–7.
- Shiskina GT, Kalinina TS, Berezova IV, et al. Stress-induced activation of the brainstem Bcl-xl gene expression in rats treated with fluoxetine: correlation with serotonin metabolism and depressive-like behavior. *Neuropharmacology* 2012;62:177-83.
- Ferguson JM. SSRI antidepressant medications: adverse effects and tolerability. *J Clin Psychiatry* 2001;3:22-7.
- Chauhan PK, Singh V. Acute and subacute toxicity study of the acetone leaf extract of *Centella asiatica* in experimental animal models. *Asian Pac J Trop Bio* 2012;2:511-3.
- Kumar A, Dogra S, Prakash A. Neuroprotective effects of *Centella asiatica* against intracerebroventricular colchicine-induced cognitive impairment and oxidative stress. *Int J Alzheimer Dis* 2009. Article ID 972178, 8 pages, doi:10.4061/2009/972178.
- Sari DCR. The relationship between memory, CA1 hippocampus and neuroglia after *Centella asiatica* leaf extract's administration in stress-induced rat (*Rattus norvegicus*), Proceeding of 2nd International Joint Symposium Frontier In Biomedical Sciences: From Genes to Applications Faculty of Medicine Gadjah Mada University. Yogyakarta;2011.
- Sari DCR, Aswin S, Suharmi S, et al. The effect of *Centella asiatica* ethanolic extract's administration duration on spatial memory in rat (*Rattus norvegicus*) after electric-stress induced. *Mutiara Medika Jurnal Kedokteran Kesehatan* 2011;3:1-10.
- Charan J, Kantharia ND. How to calculate sample size in animal studies? *J Pharmacol Pharmacother* 2013;4:303–6. doi: 10.4103/0976-500X.119726
- Paxinos G, Watson C. The rat brain. 6th ed. Philadelphia: Elsevier Inc.;2007.p.11-8.
- Kim BK, Ko IG, Kim SE, et al. Impact of several types of stresses on short-term memory and apoptosis in the hippocampus of rats. *Int Neurorol J* 2013;17:114-20.
- Tan ML, Ooi JP, Ismail N, et al. Programmed cell death pathways and current antitumor targets. *Pharm Res* 2009;26;7:1547-56.
- Uvarajan S, Janardhanam VA, Gopinath G. Neuro-shielding effect of asiaticoside against MPTP induced variations in experimental mice. *IJEST* 2012;4:960-5.
- Orhan IE. *Centella asiatica* (L.) Urban: from traditional medicine to modern medicine with neuroprotective potential. *Evid Based Complement Alternat Med* 2012;10:1-8.
- Kosten, TA, Galloway MP, Duman RS, et al. Repeated unpredictable stress and antidepressants differentially regulate expression of the Bcl-2 family of apoptotic genes in rat cortical, hippocampal and limbic brain structures. *Neuropsychopharmacology* 2008;33: 1545-58.
- Guirado R, Matarredona DS, Varea E, et al. Chronic fluoxetine treatment in middle aged rats induces changes in the expression of plasticity related molecules and in neurogenesis. *BMC Neurosci* 2012;5:1-13.
- Huang GJ, Herbert J. Stimulation of neurogenesis in the hippocampus of the adult rat by fluoxetine requires rhythmic change in corticosterone. *Biological Psychiatry* 2006;59:619-24.