

Biochemical changes in *Rumex acetosa* infected with Mancozeb resistant *fusarium proliferatum*

Ambuse MG and Bhale UN

Department of Botany, Shrikrishna Mahavidyalaya Gunjoti Tq. Omerga Dist Osmanabad-413613 (M.S.) India.
Research laboratory, Dept. of Botany, Arts, science and Commerce College Naldurg, Tq. Tuljapur. Dist. Osmanabad
413602 (M.S.)
Email: unbhale2007@rediffmail.com; mallama.10@gmail.com

Manuscript Details

Available online on <http://www.irjse.in>
ISSN: 2322-0015

Editor: Dr. Arvind Chavhan

Cite this article as:

Ambuse MG and Bhale UN.
Biochemical changes in *Rumex acetosa* infected with
Mancozeb resistant *fusarium proliferatum*, Int. Res.
Journal of Science & Engineering, 2018; Special Issue
A6: 1-5.

© The Author(s). 2018 Open Access

This article is distributed under the terms
of the Creative Commons Attribution
4.0 International License

(<http://creativecommons.org/licenses/by/4.0/>),
which permits unrestricted use, distribution, and
reproduction in any medium, provided you give
appropriate credit to the original author(s) and the
source, provide a link to the Creative Commons
license, and indicate if changes were made.

ABSTRACT

Biochemical changes were observed from healthy and artificially inoculated mancozeb resistant (At_6) and sensitive (At_5) isolates of Chuka (*Rumex acetosa* L.) caused by *Fusarium proliferatum*. There was found significant variation between healthy and infected leafy vegetables showed significant changes with respect to estimation of Moisture content, total sugar reducing sugar, non-reducing sugar, total ash, crude fiber, phosphorus, fat, ascorbic acid, calcium, iron and crude protein. Among them, moisture in the *Rumex acetosa* infected with sensitive and resistant strains was decreased. Among them total ash was increased in healthy leaf, sensitive and resistant followed by Crude protein, Phosphorus and Ascorbic acid, healthy leaf sensitive and resistant. In case of total sugar it was in reduced infected leaf. But Calcium and Fat however was decreased due to infection of both isolates.

Key word: *Rumex acetosa*, mancozeb, *Fusarium proliferatum*, biochemical changes.

INTRODUCTION

Vegetables are very important ingredients in human diet. The nutritive values of healthy vegetables are altered because of fungal attack and sometimes fungi produce certain mycotoxins in them and make them unsuitable for human consumption. Leafy vegetables have more nutrition per calorie than any other food. Chuka (*Rumex acetosa* L.) of family Polygonaceae often simply called sorrel is a perennial herb that is cultivated as leafy vegetable and medicinal properties [1]. In small quantities of sorrel is harmless, in large quantities it can be fatal. Several subspecies have been named not all are cultivated i.e. *Rumex acetosa* sp. *acetosa*, *R. acetosa* sp. *ambiguous*, *R. acetosa* sp. *arifolius*, *R. acetosa* sp. *hibernicus*, *R. acetosa* sp. *hirtulus* and *R. acetosa* sp. *Vinealis* [2,3].

Children and pregnant and nursing women should not use sheep sorrel. The herb is not recommended for people with kidney stones arthritis, endometriosis, gout and hyperacidity, since it may aggravate those conditions, due to its high acidic content and large amounts of sorrel should be avoided [4]. Many examples of fungicide resistance including mancozeb in pathogen of various crops have been reported [5,6,7]. The present investigation was made to evaluate the biochemical changes due to *Fusarium proliferatum* inciting leaf spot of *Rumex acetosa*.

MATERIAL AND METHODS

Isolation of *Fusarium proliferatum*: A survey which was commenced from June 2008 to July 2009 in growing season conducted in field and markets of Marathwada region of Maharashtra state. Delicious leafy vegetable crops are attacked by fungal diseases like, leaf spot showing symptom of fungal infection were collected and symptomatology of the disease was studied under natural and artificial conditions. Microscopic slides were prepared from the infected leafy vegetable to observe mycelium, conidiophores and conidia of the pathogens. Isolation of pathogens was done from each distinct type of symptom produced on leafy vegetable. Infected leaf spots were cut into small pieces by

sterilized blade then surface sterilized with mercuric chloride (0.1%) for 1 min. The pieces were then washed with sterilized water and dried by sterilized blotting paper. These pieces were placed on potato dextrose agar (PDA) medium and incubated at 28±1°C. The pathogens were isolated and maintained on PDA medium. The fungi were identified *Fusarium proliferatum* (Fries) Wiltshire. The cultures were deposited at Agharkar Research Institute (ARI), Pune and Department of Botany, Arts, Science and Commerce College Naldurg. Morphological characters of the pathogen were studied and pathogens were identified by Ellis [8].

Sensitivity of the pathogens against fungicide (MIC):

Sensitivity of ten isolates was tested against mancozeb (75%WP) by food poisoning technique [9]. Czapek Dox agar plates were prepared containing different concentration of (100 to 5000µg/ml) fungicide for treating both isolates. Eight days fresh culture disc (8mm) of isolates were inoculated at the centre of plates in triplicate. The plates were incubated at 28±1°C in the dark and radial growth was measured at different intervals. Plates without fungicides treated as control.

Biochemical analysis: Biochemical analysis of sensitive and resistant isolates by inoculating *Rumex acetosa* leaves were inoculated with spore suspension (105/ml) of pathogen. Uninoculated plants served as control. After 40 days the infected and healthy leaves were collected and dried in oven at 60°C for 24 hrs. The dried samples were crushed in grinder. 10gms of samples were extracted in 100 ml ethanol and analyses for 12 parameters such as Moisture, Fat, crude protein [10] Crude fiber [11], Total sugar [12], Reducing sugar [13], Non reducing sugar, Total ash [14] and [15], Ascorbic acid [16], Phosphorus, Iron and Calcium [17, 18].

RESULTS AND DISCUSSION

Isolation of *Fusarium proliferatum* (Fries) Wiltshire:

After two-three days of inoculation, ash green colonies with whitish peripheral concentric rings are formed. Isolated fungus was identified as *Fusarium proliferatum*. On the basis of cultural and morphological

characteristics the radial growth of the fungus in culture was uniform.

Sensitivity of the pathogens against fungicide MIC:

Sensitivity (MIC) *Fusarium proliferatum* was tested against Mancozeb, The results indicated in (Table 1) that there was large variation in the sensitivity of these isolates. Some isolates were sensitive At₅ (100 µg/ml) while others were resistant Fp₃(1900 µg/ml). Sensitivity ranged from (200 to 700 µg/ml) and sensitive isolates are Fp₁, Fp₄ Fp₂ and Fp₃ resistant isolates Fp₅ (3900 µg/ml). Among them, Fp₃ for resistant isolate were used for biochemical analysis.

Table1: MIC of carbendazim against *Fusarium proliferatum* isolates causing leaf spot of *Rumex acetosa*.

Isolates	MIC µg/ml
Fp ₁	1500
Fp ₂	200
Fp ₃	1900
Fp ₄	1000
Fp ₅	1600

Biochemical analysis:

Biochemical analysis was determined from leafy vegetable *Rumex acetosa* are shown in (Table.2).). It was noted that the content of all parameters in the pathogen varied in sensitive and resistant strains. Moisture in the *Rumex acetosa* infected with sensitive and resistant strains was decreased. Among them total ash was increased in healthy leaf, sensitive and resistant followed by Crude protein, Phosphorus and Ascorbic acid, healthy leaf sensitive and resistant. In case of total sugar it was in reduced infected leaf. But Calcium and Fat however was decreased due to infection of both isolates.

MIC:Minimum Inhibitory Concentration.*- Sensitive. +- Resistant.

There are reports supporting the characteristic of resistant isolates. According to Sindhan and Parashar [19] the concentration of total sugars, reducing sugars, non reducing sugars, nitrogen and phosphorus was decreased due to infection by early and late leaf spot pathogens of groundnut but the phenol content was increased. The changes of total sugar, reducing sugar, non-reducing sugar, starch, protein, ash and lipid contents decreased remarkably but those of Vitamin-C,

Table 2: Biochemical characteristics of the *Rumex acetosa* infected with mancozeb resistant and sensitive isolates of *Fusarium proliferatum*

Sr. No.	Estimation	Healthy	Sensitive (Fp5)	Resistant (Fp3)
1	Moisture (%)	49.34	35.40	30.20
2	Fat (%)	0.92	0.86	1.06
3	Crude protein (%)	20.81	13.20	20.93
4	Crude fiber	0.6	0.4	0.5
5	Total sugar (mg/g)	7.5	3.4	2.7
6	Reducing sugar (mg/g)	2.20	2.0	2.7
7	Non reducing sugar (mg/g)	5.3	1.4	0.55
8	Total Ash (%)	26.11	20.63	27.96
9	Ascorbic acid	12	7.30	8.5
10	Phosphorus	18	12.35	16.4
11	Iron (mg/100gm)	9.80	4.1	4.1
12	Calcium	0.7	0.3	0.4

phenol, pectin and mineral contents increased significantly in leaves of disease affected mulberry as compared to those in healthy mulberry [20]. Biochemical changes in pomegranate fruit infected with carbendazim resistant of *Fusarium proliferatum* showed more loss of reducing and non-reducing sugar as compared to healthy one [21]. Bhale [22] reported total sugars were reduced in infected leaves of spinach when compared with healthy one. Tofazzal Hossain [23] reported that the content of total sugars, reducing sugars was reduced due to infection by *Colletotrichum gloeosporioides*. Khillare [24] reported total sugars, total amino acids, crude protein DNA and RNA contents increased in their quantity due to infection by both the isolates of fruit rot of grape.

Acknowledgment

The authors are thankful to Dr. S.K. Singh, Scientist, ARI, Pune for identification of the fungal pathogen.

REFERENCES

1. Naik VN. Flora of Marathwada, Ammit Publications, Aurangabad 1998, 1&2.
2. Brinker FJ. Herb contraindications and drug interactions: with appendices addressing specific conditions and medicines. Sandy, Or Eclectic Institute, 1997, 146.
3. Lewis WH. Medical botany: plants affecting man's health. New York: Wiley, 1977.
4. Orlandini C. The natural poisoning of hens by sorrel (*Rumex acetosa*) and experiments on poisoning by oxalic acid and potassium binoxalate. Boll Soc Eustachiana, 1933, 31:217.
5. Gangawane LV and Saler. Resistance to fungicides in *Aspergillus flavus*. Neth. J. Pl. Path. 1981, 87- 25.
6. Hollomon DW. Mode of action of hadroxy pyrimidine fungicides. Fungicides Resistance in protection (Dekker, J. and S.G. Geogopoulous Eds.) CAPD wegeningen, Netherlands. *Horticultural* Vol. 10 (Eds: Chadha K.L. and Rethinam P.), Malhotra Publishing House, New Delhi. 1981, 1113-1138.
7. Jones AL and Ehret GR. Tolerance to fungicides in *Venturia* and *Monilinia* of tree fruits. *Proc. Ann. Phytopath. Soc.*, 1976, 3: 84-90.
8. Ellis MB. 'Dematiaceous Hyphomycetes' (Commonwealth Mycological Institute: Kew, UK) 1971.
9. Dekker J and Gielink AJ. Acquired resistance to pimaricin in *Cladosporium Cucumerinum* and *Fusarium oxysporum* sp. *narcissi* associated with decreased virulence. *Neth. J. Plant Pathol.* 1979, 85: 67-73.
10. Lowry OH, Rosebrough NJ, Farr AL, Randall RJ. Protein measurement with Folin Phenol reagent. *J. Biol. Chem.*, 1951, 193: 265-275.
11. Maynard AJ. *Methods in Food Analysis*, Academic Press, New York, 1970, 176.
12. Krishnaveni S, Theymoli Balasubramanian and Sadasivam S. *Food Chem.*, 1984, 15:229.
13. Miller GL. Use of Dinitrosalicylic Acid Reagent for Determination of Reducing Sugar. *Anal Chem.* 1959, 31:426-428.
14. AOAC. Official and tentative methods of analysis "Association of official agricultural chemist", Inc. Washington., 1975, 4 D.C.
15. Mungikar AM. *Intercropping fodder crops*. Sarswati press, Aurangabad 1999.
16. Sadasivam S and Theymoli Balasubramanian. *Practical Manual in Biochemistry*. Tamil Nadu Agricultural University Coimbatore, 1987, 14.
17. Toth SJ, Prince AL, Wallance A and Mikkelsen DS. Rapid qualitative determination of eight mineral elements in plant tissues by systematic procedure involving use of flame photometer. *Soil Sci.*, 1948, 66: 459 - 466.
18. Fiske Cyrus H and subbarao Yellapragada. The colorimetric determination of phosphorus. *Journal of Biological Chemistry*. 1925, 66 (2): 375- 400.
19. Sindhan GS and Prashar RD. Biochemical changes in groundnut leaves due to infection by early and late leaf spot pathogen. *Indian J. Mycol. Pl. pathol.* 1996, Vol 26 (2), 210- 212.
20. Ghosh L, Alam MS, Ali MR, Shohael AM, Alam F and Islam R. Changes in some biochemical parameters of Mulberry (*Morus* sp.) leaves after

- infected with leaf spot disease. *Online Journal of Biological Sciences*, 2003, 3(5): 508-514.
21. Bhale UN and Kamble SS, Gangawane LV. Biochemical changes in spinach infected with carbendazim resistant *Alternaria spinaciae*. *Indian Phytopath.* 2010, 63(2):230-231.
 22. Bhale UN, Bharade SS and Kamble SS. Biochemical changes in Pomegranate (*Punnica granatun* L.) fruit infected with Carbendazim resistant of *Alternaria alternata* (Fr.) Keissler. *Biotechnology and Bioinformatics.*, 2009, 1 (1) : 109-112.
 23. Tofazzal Hossain, MM Zahangir Alam and Absar. Changes in different nutrients and enzymes contents in mango leaves infected with *Colletotrichum gloesporiodes*. *Indian Phytopath.*, 1999, 52(1):75-76.
 24. Khillare VC, Kadam KS and Gangawane LV. Biochemical changes in host by Aureofungin resistant *Alternaria tenuis* causing fruit rot of grape. *Bioinfolet*, 2005, 2 (2): 77-80.

© 2018 | Published by IRJSE

**Submit your manuscript to a
IRJSE journal and benefit from:**

- ✓ Convenient online submission
- ✓ Rigorous peer review
- ✓ Immediate publication on acceptance
- ✓ Open access: articles freely available online
- ✓ High visibility within the field

Email your next manuscript to IRJSE
: editorirjse@gmail.com
