



Salubrious Effect of *Tylophora Asthamatica* on Paracetamol Hepatotoxicity

S. Ahmed John¹ and R. Malathi²

¹ Dept. of Botony, J.M.College, Trichy

² Dept. of Biotechnology, Bharathidasan University College, Orathanadu

Abstract. In this communication, we document the hepatoprotective effect of methonolic extract of *Tylophora asthmatica*. Over dosage of paracetamol induces hepatotoxicity. The data obtained in the present study illustrates asthmatica possess hepatoprotective effect an the paracetamol-induced hepatotoxicity in wister rats.

Keywords: *Tylophora asthmatica*, Hepatotoxicity, paracetamol

Query: (1) Ref. [7] is uncited. Please cite it in the text.

(2) “shivpuri *et al.*(1972)” is cited in the text, but not provided.

(Received: 27 April 2010)

1 Introduction

Paracetamol (Acetaminophen) is an antipyretic and analgesic drug that is available over the counter and overuse of paracetamol can cause overproduction of ROS during formation of N-acetyl-P-benzoquinoneimine (NAPQI) by cytochrome P450 [1]. This mecha-

nism has been suggested to participate in the development of oxidative stress and injury in paracetamol-induced hepatotoxicity [2].

Tylophora asthmatica, a wild indigenous plant, belongs to the family asclepiadaceae and is commonly called as Indian ipecae. It is an endangered plant species, endemic to the state of Tamilnadu in India. The powdered leaves, stems and root contain 0.2–0.3% alkaloid; of these Tylophorine, tylophorinine and tylophoridine are important alkaloids. Various studies have confirmed the anti-inflammatory activity [3] direct stimulant of adrenal cortex [4] anti-asthmatic [shivpuri *et al.* 1972] and the treatment of bronchitis, rheumatism and dermatitis [5]. The extract from leaves and stems of *T. asthmatica* have been reported to possess antedecent, insecticide and antitumour properties [6].

The aim of the present study was to evaluate the hepato protective effect of the methanolic leaves extracts of *tylophora asthmatica* using rat model.

2 Materials and Methods

Adult healthy male wistar strain rats weighing 180 to 200 g were maintained under uniform laboratory conditions in standard steel cages and provided with food [Hindustan lever Ltd, Bangalore] and water *ad libitum*.

After two weeks the animals were grouped into 4, each group contains 6 animals. Group I animals were given normal saline and fed with standard diet serve as control. Group II animals were administered a single dose of paracetamol suspension [1 g kg^{-1} body weight, *i.p.*]. This dose is known to cause liver damage in animal received META extract [200 mg kg^{-1} body weight]. Group IV animals' co treated with META extract prior to induction of liver damage with 2ml of paracetamol suspension. On day 9 which in 36 hr after paracetamol treatment, all the animals were sacrificed and the blood

samples were collected for investigations. The animals were sacrificed by cervical dislocation after anesthesia. Liver was excised, weighed. WBC and RBC were counted in the whole blood. Blood urea and cholesterol were estimated using serum. Data are expressed as mean \pm SD and subjected to Student's-*t* test for statistical significance at the level of significance $P \leq 0.05$ (Table 1).

Table 1: Effect of T. Asthmatic on Paracetamol Induced Hepatotoxicity.

Parameters	Group-I Control	Group-II Paracetamol	Group-III META	Group-IV Paracetomal META
Liver Weight (g)	3.6 \pm 0.16	2.7 \pm .16*	3.5 \pm 0.16	3.3 \pm 0.19*
WBC Count (Cells/cumm)	6055 \pm 146	6840 \pm 143*	6720 \pm 104*	6090 \pm 216
RBC Count (Cells/cumm)	18,70,000 \pm 20,000	21,50,000 \pm 15,811*	24,50,000 \pm 22,361*	19,10,000 \pm 27,386*
Urea (mg/dl)	25.5 \pm 0.86	8.4 \pm 0.4*	25.8 \pm 0.4	19.9 \pm 0.63*
Cholesterol (mg/dl)	102 \pm 3.87	135 \pm 3.39*	093 \pm 3.39	120 \pm 2.45*

* denotes statistical significance in comparison to the control at $p < 0.05$. Liver weight was determined by gravimetric method. WBC and RBC counts were by haemocytometric method. Urea was determined by DAM method and cholesterol was determined by wybenga and pilleggi.

From the Table 1, it is clear that T.asthmatica influences the activity of paracetamol at the organ, hematological and biochemical levels. The normal liver weight, which is 3.6 g, was decreased 2.7 g due to paracetamol toxicity. But when T.asthmatica is supplemented along with paracetamol, the liver weight raise up to 3.3 g.

In case of RBC and WBC the significant increase due to paracetamol treatment, from the normal value has been reduced back to almost normal condition due to supplementation of *T.asthmatica* along with paracetamol. Due to paracetamol toxicity, a drastic decrease in the blood urea level from 25.5 mg/dl to 8.4 mg/dl is observed. *META* is able to increase the urea level [19.9 mg/dl] significantly. Similarly the normal cholesterol level, 102 mg/dl which is significantly increases up to 135 mg/dl due to paracetamol, it has been decreased to 120 mg/dl in group IV. Thus *T.asthmatica* seems to reduce the hepatotoxicity produced by the paracetamol. These need further confirmation with histopathological studies.

Acknowledgements

The author wishes to acknowledge Dr.S.Ahamed John, Head, Dept. of Botany, J.M. College, Trichy, for the guidance and valuable suggestions and support.

References

- [1] Dahlin, D.C., G.T. Miwa, A.Y.H.Lu and S.D.Nelson. 1984. N-acetyl-p-benzoquinonimine. A cytochrome p. 450 mediated oxidation product of acetaminopen PNAS 81: 1327–1331.
- [2] James, L.P., P.R. Mayeux and J.A. Minson 2003. Acetaminophen induced hepatotoxicity Drug metabolism and Disposition, 31: 1499–1506.
- [3] Gopalakrishnan, C.D., Shankaranarayanan, S.K. Nazimudeen and L. Kameswaran, 1980. Effect of tylophorine, a major alkaloid of *T.India*, on immunopathological and inflammatory reactions, Indian J. Med. Res, 71: 940–948.
- [4] Udupa, A.L., S.L. Udupa and M.N.Gurusamy, 1991. The Possible site of anti asthmatic action of *T.asthmatica* on pituitary adrenal in albino rats, Planta Med., 57: 409–413.

-
- [5] Nadkarni, K., 1976, Indian Materia Medica. Vol I, Bombay, India, Popular Prakashan, p. 1252.
 - [6] Verma, G.S., V. Ramakrishnan, N.B.Mulchandani and M. Chandha, 1986. Entomol Exp. Applic., 40: 99.
 - [7] Akintonwa, A., and A.R. Essien, 1990. Protective effects of Garcinia kola seed extract against paracetamol-induced hepatotoxicity in rats. J. Ethnopharmacol. 29: 207–211.

