ISSN: 2322-0015

Chemical Analysis of Summer Honeys collected from Apis dorsata hives of Bhadrawati Tahsil of Chandrapur District of Maharashtra State (India)

Borkar Laxmikant^{1*} and Mate Devendra²

1Department of Botany, S. S. Jaiswal Arts, Comm. and Science College, Arjuni (Mor), Gondia, India.

2Department of Botany, Nutan Adrash Arts, Commerce and M.H. Wegad Science College, Umrer, Dist - Nagpur.

*Corresponding Author E-mail: borkar laxmikant@rediffmail.com

Manuscript Details

Received: 05 April, 2014 Revised: 10 May, 2014 Revised received: 29 June, 2014 Accepted: 03 July, 2014

Published: 09 July, 2014

ISSN: 2322-0015

Editor: Dr. Arvind Chavhan

Cite this article as:

Borkar Laxmikant and Mate Devendra-Chemical Analysis of Summer Honeys collected from Apis dorsata hives of Bhadrawati Tahsil of Chandrapur District of Maharashtra State (India), Int. Res. J. of Sci. & Engg., 2014; 2 (4):139-141.

Copyright: @ Author(s), This is an open access article under the terms of the Creative Commons Attribution Non-Commercial No Derivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is noncommercial and no modifica- tions or adaptations are made.

ABSTRACT

The present investigation was undertaken to determine the chemical analysis of 5 summer honey samples (CHN - BHA - KHA, CHN - BHA -CHA, CHN - BHA - MUD, CHN - BHA - KOL, CHN - BHA - BHN) collected from forest area of Bhadrawati Tahsil of Chandrapur District of Maharashtra State (India). These samples were analysed for several parameters such as moisture, total reducing sugar, Levulose or Fructose, Dextrose or Glucose, L/D ratio, sucrose, Acidity. This type of chemical analysis favours the utilization of the honey for good quality in this area.

Key words: Chemical Analysis, Summer Honey, Bhadrawati Tahsil.

INTRODUCTION

Honey is a carbohydrate rich naturally complex product produced by honey bees from floral nectar. Honey has been used by all civilization as nutrient food and in traditional medicine. The quality of honey depends on various physiological factors such as climate, soil, etc. Honey contains sugar, protein, moisture, vitamins, minerals, enzymes, polyphenols and flavonoids (Al - Manary et al., 2002) because of this unique and complex nature, honey is proved to be useful in the treatment of burns, wounds, skin ulcers as an antioxidant and in the treatment of external eye diseases (Balasubramanyam, 2011). Furthermore, honey is a highly valuable ingredient in condiments, beverage, sauces and sweets. In fact numerous studies have been reported on physical, chemical and melissopalynological parameter of honeys from all over the world. (Adenken et al., 2010; Anklam, 1998; Cherian et al., 2011; Downey et al., 2005; Ramnath and Shivaramm, 2012, Terrab et al., 2002; Xesus et al., 2010).

The scientific literature revealed that the information is not available with respect to chemical characteristics of honeys from Bhadrawati Tahsil of Chandrapur District of Maharashtra State in India. The purpose of this study was to investigate some chemical parameters such as moisture, total reducing sugar, levulose or fructose, dextrose or glucose, levulose/ dextrose, sucrose, acidity and microscopical analysis of honey collected from different regions of Bhadrawati Tahsil of Chandrapur District of Maharashtra State in India.

MATERIALS AND METHODS

Chemical analysis of the honeys are carried out by using Indian standard Specification, IS: 4941 (1974) and IS: 8464 (1977). The percentage of total reducing sugar, (Levulose or Fructose + Dextrose or Glucose), Levulose, Dextrose, Sucrose, Acidity, Moisture and L/D ratio were estimated.

RESULTS AND DISCUSSION

The chemical properties of the 5 summer honey samples (Viz. CHN – BHA - KHA, CHN – BHA – CHA, CHN – BHA – MUD, CHN – BHA – KOL, CHN – BHA - BHN) from Bhadrawati Tahsil of Chandrapur District of Maharashtra State are reported in Table.

In the present study moisture content in the samples ranges from 19 - 24.6.

Increase the temperature moisture is low and decrease the temperature moisture is high. Increase in moisture content of honey is also indicative of adultration. The low moisture content of honey forms an important part of the system which protect honey for attack by microorganism.

Sugars:

Honey consists of mostly glucose and fructose. The actual proportion of fructose to glucose, in any particular honey, depends largely on the sources of the nectar. All samples contained more fructose than glucose.

This indicates that Bhadrawati honeys would be less prone to granulation fructose level in honey is higher than that of glucose. Honey with high fructose to glucose ratio would remain liquid for longer period. The fructose/glucose ratios may have an impact or honey flavour, since fructose is much sweeter than glucose.

Acidity:

Acidity of the honey sample ranges by 0.1748 to 0.3473 respectively. Acidity values may indicate the fermentation of honey sugar by yeast.

Table: Chemical Analysis of honey samples obtained from Bhadrawati Tahsil of Chandrapur District.

Sr		Parameter						
N o	Location Of Parameter	Moisture %	Total Reducing Sugar %	Levulose or Fructose %	Dextorse or Glucose %	L/D Ratio	Sucrose %	Acidity %
1	CHN -BHA - MUD	24.4	72.81	39.467	33.314	1.317	1.675	0.2392
2	CHN – BHA – KOL	22.5	64.736	33.673	31.673	1.203	2.005	0.3473
3	CHN – BHA – KHA	23.2	71.511	39.097	32.414	1.341	0.798	0.2369
4	CHN - BHA – BHN	24.6	73.214	38.999	34.215	1.226	2.575	0.1748
5	CHN – BHA – CHN	19	72.781	40.592	32.189	1.403	2.545	0.2323

REFERENCES

- 1. Adenekan, MO, Amusa NA, Lawal AO, Okpeze VE. Physicochemical and microbiological properties of honey samples obtained from Bada, *Journal of Microbiology and Antimicrobials*, 2010; 2(8):100-104
- Al ML, Danial DJ, Moise A, Bebis O, Lasio L, Bogedanov S. Phycochemical and bioachive properties of different floral originhoneys from Romdnia. *Food Chemistry*, 2002; 112, 863-867.
- 3. Anklam EA. A review of the analogical and botanical origine of honey, *Food Chemistry*, 1998; 63, 549 562.

140 www.irjse.in

- 4. Balsubramanyam MV. Chemical Characteristics of much floral wild and apiary honeys from Western Ghats of Karnataka. *The Bioscan*, 2011: 6, 467 469.
- Cherian KJ Bhowal M and Godghate SD. Pollen and Physiochemical analysis of honey preduceed by *Apis* cerena indica of Nagpur, Maharashtra (INDIA). *Journal of* Environmental Research and Development, 2011; 5(3): 542-550.
- Downey GJ, Hussey K, Kelly JD, Walshe TF and Martin PG. Preliminary contribution to the characteristics of artisanal honey produced on the island of Ireland by palynological and physico – chemical data. *Food Chemistry*, 2005; 91, 347-354.
- 7. IS: 4941-1974 Indian Standard Specification for extracted honey (First Revision), Indian Standards Institution, 1974, New Delhi: 1-16.
- 8. IS: 8464-1977 Indian Standard Specification for Squeezed honey, Indian Standards Institution, 1977, New Delhi: 1-8.
- Ramnath Subharani and Venkataramegouda Sivaramm physicochemical and pollen analysis of Western ghats honey of Karnataka south, India. I.J. S.N.,2012: 3(4):831-835
- 10. Terrab AJ, Diez MJ and Heredia FJ. Characterization of Moroccan unifloral honeys by theis physicochemical characteristics. *Food Chemistry*, 2002: 79, 373 379.
- 11. Xesus FJ, Jose P, Maria LE, Antonio I and Jose PA. Palynological and physicochemical data charctrazation of honeys produced in the Entre Douro e Munho region of Portugal, *International Journal of food Science and Technology*, 2010; 45,: 1255-1262.

© 2014 | Published by IRJSE