# Journal of Coastal Life Medicine

journal homepage: www.jclmm.com



Original article doi: 10.12980/jclm.4.2016J6-176

©2016 by the Journal of Coastal Life Medicine. All rights reserved.

Screening of marine algae (*Padina* sp.) from the Lengeh Port, Persian Gulf for antibacterial and antifungal activities

Azadeh Taherpour<sup>1</sup>, Bita Archangi<sup>2\*</sup>, Sadraddin Ghaemmaghami<sup>3</sup>, Hossein Zolgharnein<sup>4</sup>, Kamal Ghanemi<sup>5</sup>

<sup>1</sup>Department of Marine Science and Oceanography, Faculty of Marine Biology, Khorramshahr University of Marine Science and Technology, Khorramshahr, Iran

### ARTICLE INFO

Received 5 Sep 2016 Accepted 12 Sep 2016

Article history:

Available online

Keywords:

Padina sp.

Antibacterial

Antifungal

Lengeh Port

Persian Gulf

ABSTRACT

**Objective:** To evaluate the antibacterial efficacy of different solvent extracts of *Padina* sp. against selected human pathogenic bacteria and fungi species such as *Escherichia coli*, *Shigella* sp., *Staphylococcus aureus* (*S. aureus*), *Pseudomonas aeruginosa*, *Aspergillus flavus* and *Candida albicans*.

**Methods:** Various solvents including methanol, ethyl acetate, chloroform and hexane were used to acquire crude extracts from marine algae *Padina* sp. After crude preparation, antibacterial and antifungal activities were screened against clinically important human pathogenic bacteria using disc and well diffusion methods. For all the bacterial species used in this research, minimum inhibitory concentration was undertaken considering various solvent extracts of *Padina* sp. To ensure the accuracy of experiments, a positive control was also included.

**Results:** Confirmed that hexane is the best solvent to extract antimicrobial agents from *Padina* sp. Among selected bacteria, *S. aureus* was the most sensitive test microorganism. While, all other microorganisms showed resistance against methanol, ethyl acetate, chloroform extracts. In fact, by increasing concentration of hexane extract, inhibition of *S. aureus* growth or antimicrobial activity was increased. Growth inhibition zone in well method showed better results compared to disc diffusion method. The minimum inhibitory concentration and minimum bactericidal concentration of hexane extract were 15 and 30 mg/mL against *S. aureus*, respectively. All *Padina* sp. extracts did not reveal any antifungal activities against fungi species in this study.

**Conclusions:** Brown algae extracts showed sufficient antibacterial properties against *S. aureus*. Therefore, *Padina* sp. in this research can be a good candidate to design and manufacture novel antibacterial agents used in pharmaceutical industries.

### 1. Introduction

To date, infectious diseases are one of the main causes of high mortality in global human societies. Synthetic drugs and medicines ultimately are led to increase antibiotic resistance. Infectious diseases caused by bacteria and fungi have raised awareness to improve public health and protect global economies. Drug resistant infections especially with bacterial involvement are still a major challenge. Therefore, it is essential to discover new antibiotic compounds from alternative sources including oceans and marine organisms. Marine natural sources such as seaweeds and marine algae can be excellent alternative to remedy the situation by screening, developing and manufacturing antibacterial and antifungal compounds from novel bioactive substances. There is an increasing demand for therapeutic drugs from natural products. Recently, the high potential to contribute marine organisms, especially algae to the discovery of new bioactive substances is being increased[1]. The new therapeutic agents should be effective and have a novel mechanism of action regarding resistance behavior[2]. Marine algae are potential renewable resources in the world. Algae are considered as diverse source of secondary metabolites having a broad spectrum of biological activities such as antimicrobial[3], antiviral[4], antifungal[5], anti-allergic[6], anticancer[7], anti-fouling[8] and antioxidant[9] that have been used to develop the pharmaceutical industries.

Despite the importance of medicinal properties and pharmaceutical applications of marine algae, few pharmacological researches undertaken to detect novel therapeutic agents for infectious diseases treatments<sup>[10]</sup>. There are many scientific records with reference

<sup>\*</sup>Corresponding author: Bita Archangi, Department of Marine Science and Oceanography, Faculty of Marine Biology, Khorramshahr University of Marine Science and Technology, Khorramshahr, Iran.

E-mail: bita.archangi@gmail.com

The journal implements double-blind peer review practiced by specially invited international editorial board members.

to a wide spectrum of pathogen inhibitory compounds extracted from marine algae against human bacterial and fungi pathogens worldwide<sup>[11]</sup>. Fortunately, the Iranian southeast coasts are covered by a wide variety of marine algae. The main objective of the present study was to examine the antimicrobial and antifungal effects of brown algae extracts. For this reason, *Padina* sp. was selected to evaluate bioactive compounds using different solvent systems against four human pathogenic bacteria and two fungus species.

## 2. Materials and methods

## 2.1. Algal material

The brown alga, *Padina* sp. were collected from the intertidal regions of the Lengeh Port (latitude  $54^{\circ}$  52' N; longitude  $26^{\circ}$  32' E) southeast coast of Iran, during low tide. The collected marine algae were first identified morphologically using valid identification keys. Samples rinsed thoroughly with sea water to remove sands and epiphytes and then washed with distilled water. Algae samples were shade dried for 2 weeks and then were crushed by an electric grinder, the obtained powder was stored at -20 °C until the extraction process.

### 2.2. Preparation of algal extracts

The extraction was carried out using a Soxhlet extractor for 10 h. Different solvents were used successively with gradient polarity (methanol, ethyl acetate, chloroform and hexane). The extracts were concentrated under reduced pressure by using rotary vacuum evaporator in 40 °C and then stored in refrigerator for further use[12].

#### 2.3. Microorganisms and media

Strains of bacteria [*Staphylococcus aureus* (*S. aureus*) and *Shigella* sp., *Escherichia coli* (*E. coli*), and *Pseudomonas aeruginosa* (*P. aeruginosa*)] and fungus [*Candida albicans* (*C. albicans*), and *Aspergillus flavus* (*A. flavus*)] were obtained from the Institute of Razi, Iran. The species of bacteria for susceptibility test were grown in Muller-Hinton agar (Merk, New Jersey, United States) and species of fungus were grown in Potato Dextrose agar (Oxoid, Hampshire, United Kingdom). The concentrations of bacterial and fungal suspensions were standardized by matching to the 0.5 McFarland standard. All solvents applied in this research were of high performance liquid chromatography grade.

### 2.4. Antimicrobial activity

Antimicrobial activity of crude extracts was undertaken by two ways of the agar diffusion method (well and disc). Overnight nutrient broth culture of the test organisms which adjusted to 0.5 McFarland were firmly seeded over the agar plate using sterile cotton swab. The amount of 25  $\mu$ L of each extract (100, 300 mg/mL) was poured into a 6 mm diameter well or loa ded on paper discs. Standard antibiotic, gentamicin (10 mg/disc) and nistatin (300 mg/disc) were used as positive controls and solvents were used as negative control. Inhibition zone diameters were measured in mm after incubation for 24 h at 37  $^{\circ}C[13,14]$ . Statistical analysis for all the data was undertaken.

# 2.5. Minimal inhibitory concentration (MIC) and minimal bactericidal concentration (MBC)

MIC is the lowest concentration of an antimicrobial agent (extract) which can inhibit the growth of bacteria. The MIC of extract was determined by a macrodilution broth assay which is performed according to standard National Committee for Clinical Laboratory Standards (NCCL, 2006)[15]. Two-fold serial dilution of algal extract was prepared in nutrient broth to obtain 30.00-0.46 mg/mL concentration. Briefly, added 0.5 mL nutrient broth to 1-10 tubes and 2 tubes contain 1 mL nutrient broth. Then added 30 mg/mL extract to 1, 2, 10 tubes. Dilutions were made by transferred 0.5 mL from 2 tubes to the next tube and discarded 0.5 mL from 7th tube. After completing the dilutions, added 0.5 mL of a standardized inoculums (1/500 diluted 0.5 McFarland) to 1-8 tubes. The 8th tube serves as control for growth of the organisms, 9th tube serves as control for medium and 10th tube serves as control of extract. After overnight incubation, each tubes that inhibited visible growth was recorded as MIC. A total of 100 µL of contents of each tubes that showing inhibition of growth was subcultured onto a nutrient agar plate and treatments that killing 99.9% bacteria was recorded as MBC. Statistical analyses for all the data were carried out.

### 2.6. Antifungal activity of A. flavus

Antifungal susceptibility test was carried out by two technical methods. To determine the antifungal activity of extracts by diffusion method, 6 mm of two-day culture of *A. flavus* was cultured on Potato Dextrose agar and incubated in 28 °C until 3 cm diameter of fungus was achieved. Then wells and discs containing extracts were placed with 1 cm distance, separately. After 2 days incubation, antifungal activity of extracts was evaluated and diameter of inhibition zone was recorded.

In the next method, 500  $\mu$ L of extracts was poured in plates and Potato Dextrose agar medium was added with subsequently spinning in clockwise. After colling, 6 mm *A. flavus* fungus by pipet pasteur was cultured on Potato Dextrose agar plates. Samples without any extract treatment were considered as controls. Plates were incubated at 28 °C for 2 days. The results were determined from the presence or absence of growth and size of *A. flavus*.

### 2.7. Statistical analysis

All tests were performed in triplicate and expressed as mean  $\pm$  SD. Statistical analysis was carried out using SPSS version 16 software<sup>[16]</sup>. A One-way ANOVA was used to determine the difference in treatments when significant differences between means were applied by Tukey test.

Microorganism	Methanol	Ethyl acetate	Chloroform	Hexane	Gentamicin	Nitroforantuin	Nistatin
S. aureus	-	-	-	$9.0 \pm 0.5^{*}$	$23.0 \pm 0.5^{\circ}$	$22.0 \pm 0.5^{b}$	Nt
P. aeruginosa	-	-	-	_a	$17.0 \pm 0.5^{b}$	_a	Nt
E. coli	-	-	-	_a	$9.3 \pm 0.3^{a}$	$19.0 \pm 1.1^{b}$	Nt
Shigella	-	-	-	_a	$18.0 \pm 0.5^{b}$	$21.0 \pm 0.5^{b}$	Nt
C. albicans	-	-	-	_a	Nt	Nt	$26.6 \pm 1.2^{a}$
A. flavus	-	-	-	_a	Nt	Nt	$16.3 \pm 0.8^{b}$

Results of well diffusion were quoted in parenthesis. Different letters represent the statistical comparisons between groups by using ANOVA followed *post*hoc Tukeys test (P < 0.05).

-: No activity; Nt: Not tested; \*: 300 mg of extracts.

### 3. Results

The antibacterial activities of different extracts of *Padina* sp. against 4 bacterial and 2 fungal strains are shown in Table 1. Methanol, ethyl acetate and chloroform extracts did not show any antimicrobial effects against the tested microorganisms in both disc and well diffusion assays. However, hexane extract exhibited antibacterial effect against only Gram-positive bacteria, *S. aureus*. As observed, inhibition zone of *S. aureus* were  $7.6 \pm 0.3$  and  $8.3 \pm 0.3$  in disc diffusion assay and were  $8.6 \pm 0.3$  and  $9.0 \pm 0.5$  in well diffusion test in 100 mg/mL and 300 mg/mL concentrations, respectively.

Results of antifungal activity of extracts on *A. flavus* by mixture extracts with Potato Dextrose agar medium method did not reveal any significant inhibitory effects.

### 4. Discussion

The aim of this research was to evaluate the activity of *Padina* sp. from coastal regions of Lengeh Port in northern area of Persian Gulf to produce novel bioactive agents with potential pharmaceutical applications. In fact, the capability of marine algae to produce bioactive compounds has introduced them with a potential biomedicine interest. Worldwide records proved that, marine algae possess unique structure bioactive compounds with special physiological and metabolic properties<sup>[11]</sup>. Among marine algae examined in this regard, *Padina* sp. has revealed effective antibacterial activities against most human pathogenic bacteria<sup>[11]</sup>.

Previous studies suggested different results from the impact of effective solvents on algal extracts and antimicrobial substances. Although, a variety of solvents have been applied for extracting antimicrobial compounds of seaweed, it is still uncertain what kind of solvent would be the most effective for extraction. A few researchers used different solvents for extraction of seaweed and mentioned different points. For example, Sangeetha *et al.*[17] reported that methanolic extract was the best solution. Patra *et al.*[18] indicated that chloroform and ethyl acetate were the best solutions for extracting the antibiotic principle. Our present investigation showed that marine algae extract prepared using hexane showed higher activity compared to other extracts. Our results were similar with previous investigations suggesting significant antibacterial activity in hexane extract undertaken by Ozdemir *et al.*[19] in

2006 and Demirel *et al.*<sup>[20]</sup> in 2009. These findings indicated that extraction method had definite effect on the isolation of bioactive compounds.

Antibacterial activity was exhibited with the presence of hexane extract. Hexane extract of *Padina* sp. showed inhibiting activity against *S. aureus*. Similar to this study Lima-Filho *et al.*[21] in 2002 reported hexane extract of *Ulva fasciata* inhibited the growth of *S. aureus*. However, Ozdemir *et al.*[19] in 2006, reported hexane extract of *Dictyopteris membranaceae* and *Cystoseria barbata* had no antibacterial activity against *S. aureus*.

In the current study, hexane extract was most effective against Gram-positive bacteria compared to Gram-negative bacteria. These results confirmed earlier investigations on antibacterial activity of seaweeds that established by many scientists such as Kandhasamy and Arunachalam<sup>[22]</sup> in 2008; Kolanjinathan *et al.*<sup>[23]</sup> in 2013 and Krishnapriya *et al.*<sup>[24]</sup> in 2013.

The resistance of Gram-negative bacteria towards antibiotic compounds might be due to the hydrophilic surface of outer membrane which is composed of high amounts of lipopolysaccharide molecules. This fact can act as a barrier to the penetration of numerous antibiotic molecules. Therefore, bacterial membrane are capable of breaking down the molecules introduced from outside by the enzymes located in the periplasmic space. Although, the Grampositive bacteria do not possess such outer membrane and cell wall structures[2].

In other study conducted by Patra *et al.*[25] in 2008, the antibacterial activity of methanol extract of *Sargassum* sp. in different concentrations was evaluated and reported that the high concentrations of extracts have higher antibacterial activity against *S. aureus* and *E.coli* compared to low concentrations. In addition, Demirel *et al.*[20] in 2009, showed antibacterial activity of hexane extract of brown algae is increased with increasing concentrations against *S. aureus*. These results are in accordance with present study and suggest that by increasing concentration of hexane extract, inhibition of *S. aureus* growth or antimicrobial activity is increased. Therefore, increasing the concentration of extract is led to the higher potential of antimicrobial activities.

In this research, MIC and MBC values of hexane extract were obtained 15 and 30 mg/mL, respectively. However, it should be noted that the extracts obtained in this work, were not purified and consisted of a variety of compounds. Hence, high value of purified extract is required for testing MIC and MBC to show exact

correlation between undertaken studies in this regard[26,27].

In this study, none of the examined extracts showed toxicity against *A. flavus*. This result was previously obtained by Padmakumar and Ayyakkannu<sup>[28]</sup>, which confirmed resistance of *A. flavus* against seaweed extracts. Therefore, the results of present study were similar to Tuney *et al.*<sup>[29]</sup> investigation that evaluated antibacterial and antifungal activities of methanol, acetone and diethyl ether extracts of *Padina pavonica (P. pavonica*).

There have been reports on significant antimicrobial activities of marine macroalgae by several research teams. In Vijayabaskar and Shiyamala<sup>[30]</sup> and Kolanjinathan<sup>[23]</sup> studies methanol and hexane extracts of macroalgae showed significant antimicrobial activity against *E. coli*, *P. aeruginosa*, *Shigella* and *S. aureus*. Kayalvizhi *et al.*<sup>[31]</sup> tested methanol and chloroform extracts of *Padina boergessenii* against *C. albicans*, *A. flavus*, *E. coli*, *S. aureus*. The results showed *C. albicans* was resistant against chloroform extract whereas other microorganisms were sensitive against examined extracts.

Variable toxicity of seaweed related to some factors such as habitat and seasonal time of algal collection, different growth stages of algae, variation in algal genus and species, methods of extraction and type of solvents, screening tests and the types of tested bacterial strains.

Vijayabaskar and Shiyamala<sup>[30]</sup> indicated that the methanol extracts of brown algae, *Sargassum wightii*, *Turbinaria ornata* possess strong antimicrobial activity against tested microorganisms. These differences with our study can be related to the genus and species of algae.

Ben Ali *et al.*<sup>[32]</sup> evaluated secondary metabolites and antibacterial activity of *P. pavonica* in variable seasons and demonstrated that *P. pavonica* have higher antibacterial activity in summer. The stronger activity in summer may be due to rapid growth of the algae and high rate of photosynthesis. Ktari *et al.*<sup>[33]</sup>, Marti *et al.*<sup>[34]</sup>, and Ibrahim *et al.*<sup>[35]</sup> revealed variable cytotoxic and antitumor activities of algal in variable seasons and geographic locations. Therefore, depending on the collection of samples carried out in winter, the low antimicrobial activities may be related to the harvesting season.

Geographic location also is an important factor in algal toxicity. In Ktari *et al.*<sup>[36]</sup> study, *P. pavonica* collected in spring from the eastern coast of Tunisia did not exhibit any effects against *S. aureus* while in Ben Ali *et al.* study, the same algae species collected in spring from northern coast of Tunisia showed a significant antibacterial activity against *S. aureus*.

In some studies carried out on antibacterial activity of seaweed from Oman Sea and Persian Gulf coasts, no antibacterial activity was observed from algal extracts. Movahhedin *et al.*<sup>[37]</sup> screened methanol, dichloromethane and hexane extracts of marine algae from Oman Sea coasts as antimicrobial substances and reported that the tested bacteria were resistant to extracts of algae. In another study was done by Jassbi *et al.*<sup>[38]</sup> in the Persian Gulf coast, was also showed resistance of most of the microorganisms to algal extracts. Similar to our study, microorganism were resistance to methanol, ethyl acetate and chloroform extract. This similarity may be due to geographical zone. On the other hand, producing of secondary metabolites levels in seaweed depends on ecological parameters such as salinity, temperature, nutrients, UV radiation[39].

Valgas *et al.*<sup>[40]</sup> used variable screening methods to determine antibacterial activity of natural products. The well variant of the diffusion method was more sensitive than the disc variant. The disc is composed of cellulose and many free hydroxyl groups are put out on their glucose. Thus, polar cationic antibacterial substances or compounds adsorb to the disc and cannot diffuse into the medium while apolar compounds do not adsorb on the surface of the discs and diffuse into the media. Also, the extracts that possess heavy molecular compounds cannot diffuse in agar and therefore cannot show their activity.

In current study, different extracts were used with gradient polarity (methanol, ethyl acetate, chloroform, and hexane) and only hexane extract exhibited antibacterial activity. This would be the reason that methanolic, ethyl acetatic and chloroformic extracts are able to dissolve more compounds compared to hexane extract.

It is evident from the present study that antibiotic substances of *Padina* sp. may possibly be apolar and hydrophobic compounds. However, in order to exactly clarify the issue, it is necessary to purify the antibiotic compounds and conduct subsequent isolation and purification methods to obtain fractions from *Padina* species algae which already showed a good potential of antimicrobial activities against selective bacteria.

### **Conflict of interest statement**

We declare that we have no conflict of interest.

### References

- Gonzalez del Val A, Platas G, Basilo A, Cabello A, Gorrochategui J, Suay I, et al. Screening of antimicrobial activities in red, green and brown macroalgae from Gran Canaria (Canary Islands, Spain). *Int Microbiol* 2001; 4: 35-40.
- [2] Adaikalaraj G, Patric RD, Johnson M, Janakiraman N, Babu A. Antibacterial potential of selected red seaweeds from Manapad coastal areas, Thoothukudi, Tamil Nadu, India. *Asian Pac J Trop Biomed* 2012; 2(Suppl 2): S1077-80.
- [3] Salvador N, Gomez Garreta A, Lavelli L, Ribera MA. Antimicrobial activity of Iberian macroalgae. *Sci Mar* 2007; 71: 101-14.
- [4] Bouhlal R, Haslin C, Cherman JC, Colliec-Jouault S, Sinquin C, Silmon G, et al. Antiviral activities of sulfated polysaccharides isolated from *Sphaerococcus coronopifolius* (Rhodophytha, Gigartinales) and *Boergeseniella thuyoides* (Rhodophyta, Ceramiales). *Mar Drugs* 2011; 9: 1187-209.
- [5] Guedes EA, Araujo MA, Souza AK, de Souza LI, de Barros LD, Maranhão FC, et al. Antifungal activities of different extracts of marine macroalgae against dermatophytes and *Candida* species. *Mycopathologia* 2012; **174**: 223-32.
- [6] Na HJ, Moon PD, Lee HJ, Kim HR, Chae HJ, Shin T, et al. Regulatory effect of atopic allergic reaction by *Carpopeltis affinis*. J

Ethnopharmacol 2005; 101: 43-8.

- [7] Murugan K, Iyer VV. Differential growth inhibition of cancer cell lines and antioxidant activity of extracts of red, brown, and green marine algae. *In Vitro Cell Dev Biol Anim* 2013; 49: 324-34.
- [8] Bhadury P, Wright CP. Exploitation of marine algae: biogenic compounds for potential antifouling applications. *Planta* 2004; 219: 561-78.
- [9] Kumar Chandini S, Ganesan P, Bhaskar N. *In vitro* antioxidant activities of three selected brown seaweeds of India. *Food Chem* 2008; **107**: 707-13.
- [10] Mosaddegh M, Gharanjik BM, Naghibi F, Esmaeili S, Pirani A, Eslami Tehrani B, et al. A survey of cytotoxic effects of some marine algae in the Chabahar coast of Oman Sea. *Res J Pharm* 2014; 1: 27-31.
- [11] Rosaline XD, Sakthivelkumar S, Rajendran K, Janarthanan S. Screening of selected marine algae from the coastal Tamil Nadu, South India for antibacterial activity. *Asian Pac J Trop Biomed* 2012; 2(Suppl 1): S140-6.
- [12] Sujatha L, Lalitha Govardhan T, Subba Ranaiah G. Antibacterial activity of green seaweeds on oral bacteria. *Indian J Nat Prod Resour* 2012; 3(3): 328-33.
- [13] Kolanjinathan K, Stella D. Antibacterial activity of marine macro algae against human pathogens. *Recent Res Sci Technol* 2009; 1: 20-2.
- [14] Seenivasan R, Indu H, Archana G, Geetha S. The antibacterial activity of some marine algae from south east coast of India. *Am-Eurasian Agric Environ Sci* 2010; **9**: 480-9.
- [15] Wikler MA. Methods for dilution antimicrobial susceptibility tests for bacteria that grow aerobically; Approved Standard. 7 ed. Wayne: Clinical and Laboratory Standards Institute; 2006.
- [16] Alghazeer R, Whida F, Abduelrhman E, Gammoudi F, Naili M. *In vitro* antibacterial activity of alkaloid extracts from green, red and brown macroalgae from western coast of Libya. *Afr J Biotechnol* 2013; **12**(51): 7086-91.
- [17] Sangeetha S, Dhayanithi NB, Sivakumar N. Antibacterial activity of sargassum longifolium and *Gracilaria corticata* from Gulf of Mannar against selected common shrimp pathogens. *Int J Pharm Bio Sci* 2014; 5(2): 76-82.
- [18] Patra JK, Patra AP, Mahapatra NK, Thatoi HN, Das S, Sahu RK, et al. Antimicrobial activity of organic solvent extracts of three marine macroalgae from Chilika Lake, Orissa, India. *Malaysian J Microbiol* 2009; **5**: 128-31.
- [19] Ozdemir G, Horzum Z, Sukatar A, Karabay-Yavasoglu NU. Antimicrobial activities of volatile components and various extracts of *Dictyopteris membranaceae*. And *Cystoseira barbata*. From the coast of Izmir, Turkey. *Pharm Biol* 2006: 44: 183-8.
- [20] Demirel Z, Yilmaz-Koz FF, Karabay-Yavasoglu UN, Ozdemir G, Sukatar A. Antimicrobial and antioxidant activity of brown algae from the Aegean Sea. J Serb Chem Soc 2009: 74: 619-28.
- [21] Lima-Filho JVM, Carvalho AFFU, Freitas SM, Melo VMM. Antibacterial activity of extracts of six macroalgae from the northeastern Brazilian coast. *Braz J Microbiol* 2002; 33: 311-4.
- [22] Kandhasamy M, Arunachalam KD. Evaluation of *in vitro* antibacterial property of seaweeds of southeast coast of India. *Afr J Biotechnol* 2008; 7: 1958-61.

- [23] Kolanjinathan K, Ganesh P, Saranraj P and Sekar D. Antimicrobial activity of *Gracilaria folifera* extracts against pathogenic microorganisms. *Int J Curr Biochem Biotechnol* 2013; 2: 6-9.
- [24] Krishnapriya M, john Milton M, Arulvasu C, Gajendran B. Evaluation of antibacterial properties of selected red seaweeds from Rameshwaram, Tamil Nadu, India. *J Acad Ind Res* 2013; 1(11): 667-70.
- [25] Patra JK, Rath SK, Jena K, Rathod VK, Thatoi H. Evaluation of antioxidant and antimicrobial activity of seaweed (*Sargassum* sp.) extract: a study on inhibition of Glutathione-S-transferase activity. *Turk J Biol* 2008; **32**: 119-25.
- [26] Bansemir A, Blume M, Schroder S, Lindequist U. Screening of cultivated seaweeds for antibacterial activity against fish pathogenic bacteria. *Aquaculture* 2005; 252: 79-84.
- [27] Dashtiannasab A, Kakoolaki S, Sharif Rohani M, Yeganeh V. In vitro effects of Sargassum latifolium (Agardeh, 1948) against selected bacterial pathogens of shrimp. Iran J Fish Sci 2012; 11: 765-75.
- [28] Padmakumar K, Ayyakkannu K. Seasonal variation of antibacterial and antifungal activities of the extracts of marine algae from Southern coasts of India. *Bot Mar* 1997; 40: 507-15.
- [29] Tuney I, Cadirci BH, Ünal D, Sukatar A. Antimicrobial activities of the extracts of marine algae from the coast of Urla (Izmir, Turkey). [Turk J Biol 2006; 30: 171-5.
- [30] Vijayabaskar P, Shiyamala V. Antibacterial activities of brown marine algae (*Sargassum wightii* and *Turbinaria ornata*) from the Gulf of Mannar Biosphere Reserve. Adv Biol Res 2011; 5: 99-102.
- [31] Kayalvizhi K, Subramanian V, Anantharaman P, Kathiresan K. Antimicrobial activity of seaweeds from the Gulf of Mannar. Int J Pharm Appl 2012: 3: 306-14.
- [32] Ben Ali AI, Ktari L, Boudabbous A, El Bour M. Seasonal variation of antibacterial activity of the brown alga *Padina pavonica* (L) thivy collected from northern coast of Tunisia. *Bull Inst Natn Scien Tech Mer de Salammbô* 2010; **37**: 111-6.
- [33] Ktari L, Guyot M. A cytotoxic oxysterol from the marine alga Padina pavonica (L.) Thivy. J Appl Phycol 1999; 11: 511-3.
- [34] Marti R, Uriz MJ, Turon X. Seasonal and spatial variation of species toxicity in Mediterranean seaweed communities: correlation to biotic and abiotic factors. *Mar Ecol Prog Ser* 2004; 282: 73-85.
- [35] Ibrahim AMM, Mostafa MH, El-Masry MH, El-Naggar MMA. Active biological materials inhibiting tumor initiation extracted from marine algae. *Egypt J Aquat Res* 2005; **31**: 146-55.
- [36] Ktari L, Longeon A, Guyot M. Biological activities of marine algae from Tunisian coasts. J Res Oceanogr 2001; 26: 37-41.
- [37] Movahhedin N, Barar J, Fathi Azad F, Barzegari A, Nazemiyeh H. Phytochemistry and biologic activities of Caulerpa peltata native to Oman Sea. *Iran J Pharm Res* 2014; 13: 515-21.
- [38] Jassbi AR, Mohabati M, Eslami S, Sohrabipour J, Miri R. Biological activity and chemical constituents of red and brown algae from the Persian Gulf. *Iran J Pharm Res* 2013; 12(3): 339-48.
- [39] Plaza M, Cifuentes A, Ibanez E. In the search of new functional food ingredients from algae. *Trends Food Sci Technol* 2008; 19: 31-9.
- [40] Valgas C, de Souza SM, Smania EFA, Smania Jr A. Screening methods to determine antibacterial activity of natural products. *Braz J Microbiol* 2007; 38: 369-80.