

Advancement in Medicinal Plant Research Vol. 2(2), pp. 24-26, May 2014 ISSN: 2354-2152

Short Communication

Effect of aqueous extract of *Hibiscus sabdariffa* L. on the weight and the activity of superoxide dismutase (SOD) and malondialdehyde (MDA) in the brain of *Rattus norvegicus*

E. Jeroh¹*, E. P. Awhin², C. A. Alalor³ and F. C. Chukwudebeli²

¹Department of Physiology, Faculty of Postgraduate Studies, Delta State University, Abraka, Nigeria.

Accepted 2 May, 2014

ABSTRACT

The effect of aqueous extract of *Hibiscus sabdariffa* L. on the weight and the activity of superoxide dismutase (SOD) and malondialdehyde (MDA) in the brain of *Rattus norvegicus* was studied in this research. The results from this study revealed that the consumption of *zobo* drink *ad libitum* increased the weight of the treated animals so much (223.00 \pm 8.30*) when compared with that of the control (176.00 \pm 3.76) at P < 0.05. The result of this study also revealed that the consumption of *zobo* drink *ad libitum* increased the enzyme activity of the enzyme superoxide dismutase (SOD) of the treated rats (8.78 \pm 0.15) when compared with that of the control (6.83 \pm 0.88) at P < 0.05. This was not the same for malondialdehyde (MDA) as the treated had a reduced enzyme activity (3.90 \pm 0.34) when compared with that of the control (4.51 \pm 0.25) at P > 0.05. The result from this study has shown that *zobo* drink, if taken in adequate manner serve as an antioxidant which reduces the activity of oxidants within the cells of the brain of the treated rats thus, the consumption of *zobo* drink *ad libitum* reduced oxidation reaction in the brain cells which is good for health.

Keywords: Zobo drink, Hibiscus sabdariffa L., superoxide dismutase, malondialdehyde, Rattus novergicus.

*Corresponding author. E-mail: onome4jeroh@yahoo.com.

INTRODUCTION

²Department of Medical Biochemistry, College of Basic Medical Sciences, Delta State University, Abraka, Nigeria.

³Department of Pharmaceutics and Industrial Pharmacy, Delta State University, Abraka, Nigeria.

REFERENCES

- Alscher RG, Erturk N, Heath LS, 2002. Role of superoxide dismutases (SODS) in controlling oxidative stress in plants. J Expt Biol, 53(372):1331-1341.
- Awhin EP, 2012. The effect of aqueous extract of *Habiscus sabdariffa* L. on the activity of alkaline phosphatase and uric acid in the Liver of *Rattus novergicus*. Cont J Med Res, 6:12-15.
- Carvajal-Zarrabal O, Waliszewski SM, Barradas-Dermitz DM, Orta-Flores Z, Hayward-Jones PM, Nolasco-Hipólito C, Angulo-Guerrero O, Sánchez-Ricaño R, Infanzón RM, Trujillo PR, 2005. The consumption of *Hibiscus sabdariffa* dried calyx ethanolic extract reduced lipid profile in rats. Plant Foods Hum Nutr, 60(4):153-159.
- Dahiru D, Obi OJ, Umaru H, 2003. Hibiscus sabdariffa calyx extract on carbon tetrachloride induced liver damage. BIOKEMISTRI, 15(1):27-33.
- Ezugwu CO, 2003. Haemostatic activity of the leaf extract of *Habiscus* sabdariffa L. J Trop Med Plant, 3:175-180.
- Marnett LJ, 1999. Lipid peroxidation DNA damage by malondialdehyde. Mutat Res, 424(1-2):83-95.
- McKay DL, Chen CY, Saltzman E, Blumberg JB, 2010. *Habiscus* sabdariffa L. tea (Tisane) lowers blood pressure in prehypertensive and mildly hypertensive adults. J Nutr, 140(2):298-303.
- Mohammed R, Fernadez J, Pineda M, Aguilar M, 2007. Roselle *Hibiscus sabdariffa* seed oil is a rich source of γ-tocopherol. J Food Sci, 72:S207-S211.
- Odigie IP, Ettarh RR, Adigun SA, 2003. The effect of aqueous extract of petals of *Hibiscus sabdariffa* L on the established stages of 2 kidney 1 cap removascular hypertension. J Ethnopharmacol, 86:181-185.
- Okasha MAM, Abubaka MS, Bako IG, 2008. Study of the effect of aqueous *Hibiscus sabdariffa* Linn seed extract on serum prolactin level of lactating female albino rats. Euro J Sci Res, 22(4):575-583.
- Pelvig DP, Pakkenberg H, Stark AK, Pakkenberg B, 2008. Neocortical glial cell numbers in human brains. Neurobiol Aging, 29:1754-1762.
- Poon HF, Calabrese V, Scapaginini G, Butterfield DA, 2004. Free radicals and brain aging. Clin Geriat Med, 20:329-359.
- Suboh SM, Bilto YY, Aburjai TA, 2004. Protective effects of selected medicinal plants against protein degradation, lipid peroxidation and deformability loss of oxidatively stressed human erythrocytes. Phytother Res, 18(4):280-284.
- Tseng TH, Hsu JD, Lo MH, Chu CY, Chou FP, Huang CL, Wang CJ, 1998. Inhibitory effect of Hibiscus protocatechuic acid on tumor promotion in mouse skin. Cancer Lett,126: 199-207.
- Vilasine H, Anocha U, Noppawan PM, Nuntavan B, Hitoshi S, Angkana H, Chuthamanee S, 2005. Antioxidant effects of aqueous extracts from dried calyx of *Hibiscus sabdariffa* L (Roselle) *in vitro* using rat low density lipoprotein (LDL). Biol Pharm Bull, 28:481-484.
- Wang CJ, Wang JM, Lin WL, Chu CY, Chou FP, Tseng TH, 2000. Protective effect of Hibiscus anthocyanins against tert-butl hydroperoxides-induced hepatic toxicity in rats. Food Chem Toxicol, 38:411-416.