

# Antiamoebic activity and cytotoxicity of ethanolic extract of *Cyperus rotundus* L.

Ahmed S. Kabbashi<sup>1</sup>, El-badri E. Osman<sup>2\*</sup>, Amel M. Abdrabo<sup>1</sup>, Nadir Abuzeid<sup>3</sup>, Mohammed I. Garbi<sup>4</sup>, Waleed S. Koko<sup>5</sup> and Mahmoud M. Dahab<sup>6</sup>

<sup>1</sup>Medicinal and Aromatic Plants and Traditional Medicine Research Institute (MAPTMRI), P.O. Box 2404, National Center for Research, Khartoum, Sudan.

<sup>2</sup>Elsheikh Abdallah Elbadri University, Berber, Sudan.

<sup>3</sup>Department of Clinical Microbiology, Faculty of Medical Laboratory, Omdurman Islamic University, Omdurman, Sudan.

<sup>4</sup>Department of Microbiology, Faculty of Medical Laboratory Sciences, International University of Africa, P.O. Box 2469 Khartoum, Sudan.

<sup>5</sup>College of Science and Arts in Ar Rass, University of Qassim, K.S.A.

<sup>6</sup>Department of Microbiology, Faculty of Pure and Applied Sciences, International University of Africa, P.O. Box 2469 Khartoum, Sudan.

Accepted 21 October, 2015

---

## ABSTRACT

*Cyperus rotundus* L. (Family: Cyperaceae), has a number of pharmacological and biological activities including anti-inflammatory, antidiabetic, antidiarrhoeal, cytoprotective, antimutagenic, antimicrobial, antioxidant, cytotoxic and apoptotic, anti-pyretic and analgesic activities. The World Health Organization (WHO) estimates that the protozoan *Entamoeba histolytica* is a major cause of morbidity worldwide, causing approximately 50 million cases of dysentery and 100,000 deaths annually. Intestinal amoebiasis due to the infection of *E. histolytica* is ranked third on the list of parasitic protozoan infections leading to death behind malaria and schistosomiasis. The purpose of the paper was to investigate the *in-vitro* antiamoebic activity and cytotoxicity (MTT assay) of ethanol extract of *C. rotundus* L. (whole plant). The ethanol extract of *C. rotundus* (whole plant), with different concentration (500, 250 and 125 ppm) and metronidazole concentration (312.5 µg/ml) was investigated *in vitro* against *E. histolytica* trophozoites. The result was obtained from *C. rotundus* whole plant ethanol extract which exhibited 100% mortality within 96 h, at a concentration 500 ppm; this was compared with Metronidazole which gave 96% inhibition at concentration 312.5 µg/ml at the same time. In addition cytotoxicity (MTT assay) with different concentration (500, 250 and 125 ppm) in comparison to triton-x100 (the reference control) which verified the safety of the examined extract with an IC<sub>50</sub> less 100 µg/ml. In conclusion, these studies prove the potent activity of *C. rotundus* against *E. histolytica* trophozoites *in vitro* with verified safety evidence for use.

**Keywords:** *In vitro*, antiamoebic, *Entamoeba histolytica*, Metronidazole, cytotoxicity (MTT-assay), *C. rotundus* (whole plant).

---

\*Corresponding author. E-mail: elbadri150@yahoo.com.

---

## INTRODUCTION

Medicinal plants are still invaluable source of safe, less toxic, lower price, available and reliable natural resources of drugs all over the world. People in Sudan and in other developing countries have relied on traditional herbal preparations to treat themselves. Therefore, it is useful to investigate the potential of local plants against these

disabling diseases (Amaral et al., 2006; Koko et al., 2008).

*Cyperus rotundus* L.; (Family: Cyperaceae) is the largest family in the monocotyledons consisting of 109 genera and approximately 5,500 species (Govaerts et al., 2007). Also known as purple nut, sedge or nut grass is a

common perennial weed with slender, scaly creeping rhizomes, bulbous at the base and arising singly from the tubers which are about 1 to 3 cm long. The tubers are externally blackish in colour and reddish white inside, with a characteristic odour. The stems grow to about 25 cm tall and the leaves are linear, dark green and grooved on the upper surface. Inflorescences are small, with 2 to 4 bracts, consisting of tiny flowers with a red-brown husk. The nut is three-angled, oblong-ovate, yellow in colour and black when ripe. *C. rotundus* is indigenous to India, but are now found in tropical, subtropical and temperate regions (Uddin et al., 2006).

In Asian countries, the rhizomes of *C. rotundus*, which are used as traditional folk medicines for the treatment of stomach and bowel disorders, and inflammatory diseases, have been widely investigated (Dang et al., 2011; Gupta et al., 1971; Won-Gil et al., 2001). *C. rotundus* is a traditional herbal medicine (Weenen et al., 1999; Zhu et al., 1997). The tuber part of *C. rotundus* is one of the oldest known medicinal plants used for the treatment (Yu et al., 2004; Zeid et al., 2008). Infusion of this herb has been used (Umerie and Ezeuzo, 2000). It is a multipurpose plant, widely used in traditional medicine around the world as herbal remedies (Oliver-Bever, 1986; Puratuchikody et al., 2006; Joshi and Joshi, 2000; El-Kamali and El-Khalifa, 1999). A number of pharmacological and biological activities have been reported for this plant (Zhu et al., 1997; Durate et al., 2005; Sundaram et al., 2008; Raut et al., 2006; Kilani et al., 2005; Kilani et al., 2007; Kilani et al., 2008; Dhillon et al., 1993; Pal and Dutta, 2006; Neffatti et al., 2008).

*C. rotundus* has a broad spectrum of applications as herbal remedies in China, Africa, Latin America, India, Saudi Arabia and Sudan (Nandihalli and Bendixen, 1987). In Chinese pharmacopoeia, it was described as an agent to regulate circulation, normalize menstruation, and relieve pain (Huang, 1999).

In Sudan the tubers of *C. rotundus* L. are used in stomach disorders and bowels irritation. An infusion of the tubers is used in dyspepsia, diarrhea, dysentery, as cures, vomiting, cholera and fevers. The tubers are given in large doses as an anthelmintic. A poultice of the fresh tubers is used to cure wounds, ulcers and sores; it is also applied to the breast to promote the flow of milk. Paste is used in scorpion stings (El-Ghazali et al., 1994).

The methanolic extract of the tubers showed an anti-inflammatory effect for the treatment of inflammatory diseases mediated by over production of nitric oxide and superoxide (Seo et al., 2001). Moreover, it showed significant antidiarrhoeal activity in castor oil induced diarrhea in mice (Seo et al., 2001). The dried tubers are used to treat dysmenorrhea and other menstrual irregularities. The aqueous extract of the dried tubers has an inhibitory effect on the uterus, (uterine relaxation) in both pregnant and non-pregnant women, and relieving pain. The herb can stimulate gastric and salivary secretion. In addition, the aqueous extract of the dried tubers has antibacterial and anti-malarial effects (Huang,

1999).

The World Health Organization (WHO) estimates that the protozoan *Entamoeba histolytica* is a major cause of morbidity worldwide, causing approximately 50 million cases of dysentery and 100,000 deaths annually (WHO, 1997; Ravdin and Stauffer, 2005). Intestinal amoebiasis due to the infection of *E. histolytica* is ranked third on the list of parasitic protozoan infections leading to death behind malaria and schistosomiasis (Farthing et al., 1996). This infection remains a significant cause of morbidity and mortality world-wide (Stanley and Reed, 2001). Amoebiasis is a rare occurrence in developed countries of the world, but only found in travelers, immigrants, homosexuals and institutionalized persons. *E. histolytica*-associated dysentery is a common occurrence in the less developed and developing countries of the world, but is more common in areas of low socioeconomic status, poor sanitation and nutrition especially in the tropics (Ravdin and Stauffer, 2005).

Thus the majority of *E. histolytica* infections, morbidity and mortality occur in Africa, Central and South America and the Indian sub-continent (Haque et al., 2000). Metronidazole is the drug now widely used and recommended in the treatment of amoebiasis (Townson et al., 1994). But it is less effective in the tissue than in the gut lumen (Bhopale et al., 1995). In addition, it can eradicate only up to 50% of laminae infections (Tierney et al., 1998). Metronidazole sometimes causes adverse effects, example, myoplasia, neuralgia, and allergic dermatitis (Upcroft et al., 2006). Hence, the present study was conducted to investigate the anti-amoebic activity and cytotoxicity of *C. rotundus* (whole plant) in Sudan.

## MATERIALS AND METHODS

### Plant materials

*C. rotundus* (whole plant) was collected from Central Sudan during the period of January to February 2014. The plant was identified and authenticated by the taxonomists of Medicinal and Aromatic Plants and Traditional Medicine Research Institute (MAPTMRI), Khartoum, Sudan.

### Preparation of crude extract

Extraction was carried out for the whole plant of *C. rotundus* by using overnight maceration techniques according to the method described by Harbone (1984). About 50 g of powdered material was macerated in 250 ml of ethanol for 3 h at room temperature. Occasional shaking for 24 h at room temperature was performed and, the supernatant was decanted. Thereafter, the supernatant was filtered under reduced pressure by rotary evaporator at 55°C. Each residue was weighed and the yield percentage was calculated and then stored at 4°C in tightly sealed glass vial ready for use. The remaining extracts which were not soluble were successively extracted using ethanol with the described technique. The extracts were kept in freeze dryer for 48 h, (Virtis, USA) until they were completely dried. The residue was weighed and the yield percentage was calculated. The extracts were kept and stored at 4°C until required.

### **In vitro testing of extract for antiameobic activity**

#### **Parasite isolate**

*E. histolytica* used in all experiments were taken from patients of Ibrahim Malik Hospital (Khartoum). All parasite samples were examined by wet amount preparation; the positive samples were transported to the laboratory in nutrient broth medium. Trophozoites of *E. histolytica* were maintained in RPMI 1640 medium containing 5% bovine serum at  $37 \pm 1^\circ\text{C}$ . The parasite trophozoites were maintained for the assays and were employed in the log phase of growth.

#### **Inoculum**

*E. histolytica* was inoculated in the RPMI-1640 medium and incubated at  $37 \pm 1^\circ\text{C}$  for 48 h. parasites were counted under the microscope by haemocytometer chamber.

#### **In vitro susceptibility assays**

*In vitro* susceptibility assays used the sub-culture method Cedillo-Rivera et al. (2002), which is being described as a highly stringent and sensitive method for assessing the anti-protozoal effects (gold standard) particularly in *E. histolytica*, *Gairdia intestinalis* and *Trichomonas vaginalis* (Arguello-Garcia et al., 2004). 5 mg from

each extract and compound was dissolved in 50  $\mu\text{l}$  of dimethyl sulfoxide (DMSO) at Eppendorf tube containing 950  $\mu\text{l}$  D.W in order to reach concentration of 5 mg/ml (5000 ppm). The concentrates were stored at  $-20^\circ\text{C}$  for further analysis. Sterile 96-well microtitre plate was used for different plant extracts, positive control and negative control.

Three columns of a microtitre plate wells [8 columns (C)  $\times$  12 rows (R)] were chosen for each extract, 40  $\mu\text{l}$  of an extract solution (5 mg/ml) were added to the first column wells C-1: On the other hand, 20  $\mu\text{l}$  of complete RPMI medium were added to the other wells of the second column and third column (C-2 and C-3). Serial dilutions of the extract were obtained by taking 20  $\mu\text{l}$  of extract to the second column wells and taking 20  $\mu\text{l}$  out of the complete solution in C-2 wells to C-3 wells and discarding 20  $\mu\text{l}$  from the total solution of C-3 to the remaining 20  $\mu\text{l}$  serial solutions in the successive columns. 80  $\mu\text{l}$  of culture medium was complemented with parasite and added to all wells. The final volume in the wells was 100  $\mu\text{l}$ .

In each test, Metronidazole (a trichomonocide) pure compound [(1-(2-hydroxyethyl)-2-methyl-5 Nitroimidazole], was used as positive control in concentration 312.5  $\mu\text{g/ml}$ , whereas untreated cells were used as a negative controls (culture medium plus trophozoites). For counting, the samples were mixed with Trypan blue in equal volume. The final number of parasites was determined with haemocytometer three times for counting after 24, 48, 72 and 96 h. The mortality % of parasite for each extracts activity was carried out according to the following formula:

$$\text{Mortality of parasites (\%)} = \frac{\text{Negative control} - \text{tested sample with extract}}{\text{Negative control}} \times 100\%$$

Only 100% inhibition of the parasite was considered, when there was no motile parasite observed.

#### **Cytotoxicity screening**

Microculture tetrazolium MTT assay was utilized to evaluate the cytotoxicity of the *C. rotundus*.

#### **Microculture tetrazolium (MTT) assay**

##### **Principle**

This colorimetric assay is based on the capacity of mitochondria succinate dehydrogenase enzymes in living cells to reduce the yellow water soluble substrate 3- (4, 5-dimethyl thiazol-2-yl)-2, 5-diphenyl tetrazolium bromide (MTT) into an insoluble, blue colored formazan product which is measured spectrophotometrically. Since reduction of MTT can only occur in metabolically active cells, the level of activity is a measure of the viability of the cells (Patel et al., 2009).

##### **Preparation of *C. rotundus* extract**

Using a sensitive balance 5 mg of each extracts were weighed and put in eppendorf tubes. 50  $\mu\text{l}$  of DMSO were added to the extract and the volume was completed to 1 ml with distilled water obtaining a concentration of 5 mg/ml. The mixture was vortexed and stirred by magnetic stirrer to obtain a homogenous solution.

#### **Cell line and culturing medium**

Vero (Normal, African green monkey kidney) cells were cultured in a culturing flask containing a complete medium consisting of 10% fetal bovine serum and 90% minimal essential medium (MEM) and then incubated at  $37^\circ\text{C}$ . The cells were sub cultured twice a week.

#### **Cell line used**

Vero cells (Normal, African green monkey kidney).

#### **Cell counting**

Cell counts were done using the improved Neubauer chamber. The cover slip and chamber were cleaned with detergent, rinsed thoroughly with distilled water and swapped with 70% ethanol, then dried. An aliquot of cell suspension was mixed with equal volume of 0.4% trypan blue in a small tube. The chamber was charged with cell suspension. After cells had settled, the chamber was placed under light microscope. Using 40 X objective, cells in the 4 large corner squares (each containing 16 small squares) were counted. The following formula was used for calculating cells:

$$(\text{Cells/ml}) N = \frac{\text{Number of cells counted} \times \text{Dilution factor} \times 10^4}{4}$$

**Table 1.** Preliminary quantitative data on yield percentage (%) of *C. rotundus* (whole plant) used for the antiameobic activity and cytotoxicity study.

Scientific name of plant	Family name	Part used	Yield (%)
<i>Cyperus rotundus</i> L.	Cyperaceae	Whole plant	10.5

### Procedure

The monolayer cell culture formed in the culturing flasks was trypsinized and the cells were put in centrifuging tube and centrifuged for 5 min separating the cells from the supernatant that flicked out. 1 ml complete medium was added to the cells and all the cell suspension was contained in a basin. In a 96-well microtitre plate, serial dilutions of each extracts were prepared. 3 duplicated concentrations for each extracts, that is, 6 wells for each of 8 extracts. All wells in rows A, B and C were used in addition to first 4 wells from each rows D, E and F. The first 2 wells of row G were used for the negative control and the first 2 wells of row H were used for the positive control Triton X. 20 µl complete medium pipetted in all wells in rows B, C and mentioned wells of rows E and F. Then 20 µl from each extracts were pipetted in rows A and B and first 4 wells of rows E and F. 20 µl taken from row B were pipetted and mixed well in row C from which 20 µl were taken and flicked out. The same was done from E to F. After that 80 µl complete medium were added to all used wells. Then adjusting the cell account to 3000 cell/well, 100 µl of cell suspension were added completing all wells to the volume 200 µl. Now, we have duplicated three concentrations 500, 250, 125 µg/ml for each extract. Then the plate was covered and incubated at 37°C for 96 h.

After four days, the supernatant was removed from each well without detaching the cells. MTT suspension stock (5 mg/ml) prepared earlier in 100 ml phosphate buffer solution (PBS) was diluted (1:3.5) in a culture medium. To each well of the 96-well plate, 50 µl of diluted MTT were added. The plate was incubated for further 4 h at 37°C. MTT was removed carefully without detaching cells, and 100 µl of DMSO were added to each well. The plate was agitated at room temperature for 10 min then read at 540 nm using microplate reader. The percentage growth inhibition was calculated using the formula below:

$$\% \text{ Cell inhibition} = 100 - \left\{ \frac{(Ac - At)}{Ac} \right\} \times 100$$

Where, At = Absorbance value of test compound; Ac = Absorbance value of control.

### Statistical analysis

All data were presented as means ± S.D. Statistical analysis for all the assay results were done using Microsoft Excel program (2007).

## RESULTS AND DISCUSSION

The whole plant of *C. rotundus* family (Cyperaceae) was screened for antiameobic activity against (*Entamoeba histolytica*) trophozoites *in vitro*, and tested for cytotoxicity using 3-(4, 5-dimethyl thiazol-2-yl)-2, 5-diphenyl tetrazolium bromide (MTT) Vero cell line after the evaluation of the yield percentage of the plant which was found to be 10.5% (Table 1).

The antiameobic potential of the ethanolic extract of *C. rotundus* (whole plant), with different concentrations (500,

250 and 125 ppm) and Mertronidazole (the reference control) with concentration (312.5 µg/ml) was investigated against *E. histolytica* trophozoites *in vitro*. However, the ethanol extract of *C. rotundus* (whole plant) showed 100% inhibition at a concentration 500 µg/ml after 96 h; which was compared with Metronidazole giving 96% inhibition at concentration 312.5 µg/ml at the same time against *E. histolytica* (Figure 1).

The whole plant of *C. rotundus* which was screened for antiameobic activity against (*E. histolytica*) trophozoites *in vitro* showed antiameobic activity with an inhibition concentrations (IC) more than 1.32 µg/ml and increasing successively during the days of the screening (Table 2).


Moreover, these results obtained in this study are similar to studies carried out for antimalaria activity by Thebtaranonth and Thebtaranont (1995) and the previous comprehensive screening of Sudanese medicinal plants for their antiprotozoal activity (Samia et al., 2004; Ali et al., 2002; Koko, 2005).

The promising activity of the plant in this study may be explained for the chemical constituents- Cyprotene, cypera-2, 4-diene, a-copaene, cyperene, aselinene, rotundene, valencene, ylanga-2, 4- diene, g-gurjunene, trans-calamenene, d-cadinene, g-calacorene, epi-aselinene, a-murolene, g-murolene, cadalene, nootkatene which the plant proved to contain (Joulain and Konig, 1998), in addition to mustakone, cyperol (Nyasse et al., 1988), isocyperol (Hikino and Takemoto, 1967) cyperotundone (Hikino et al., 1966), and acyperone (Howe and Mc Quillin, 1955; Haaksma et al., 1992).

Interestingly, the cytotoxicity assays were conducted in this study to evaluate the cytotoxicity effects of ethanolic extract of *C. rotundus* (whole plant) by using MTT-assay including (Vero cell line). Table 3 indicated the inhibition percentage (%) of Vero cell line growth *in vitro* by ethanolic extract of *C. rotundus* (whole plant) for different concentrations 125 to 500 µg/ml and showed an IC<sub>50</sub> > 100 (µg/ml) which is verifying the plant safety. This result was similar to that produced by Ahmed et al. (2012), who found that the plant extract gave similar result using the Brine Shrimp Bioassay.

## CONCLUSION

This result enhances the ethno botanical uses of *C. rotundus* (whole plant) as antiameobic in cases associated with amoebiasis in Sudan. Further investigations regarding the mode of action and other



**Figure 1.** *In vitro* activity of *C. rotundus* ethanol extract against *E. histolytica*.

**Table 2.** Inhibition concentration (IC) *C. rotundus* (whole plant) ethanol extract against *E. histolytica*.

IC ( $\mu\text{g/ml}$ )	IC <sub>50</sub>	IC <sub>90</sub>	IC <sub>95</sub>	IC <sub>99</sub>
After one day	1.32	918.97	2081.79	4004.52
After two days	1.43	432.61	979.14	1882.12
After three days	1.58	266.59	506.19	845.45
After four days	4.97	207.44	330.76	480.40

Key: IC<sub>50</sub> Inhibition concentration 50%, IC<sub>90</sub> Inhibition concentration 90%, IC<sub>95</sub> Inhibition concentration 95%, IC<sub>99</sub> Inhibition concentration 99%.

**Table 3.** Cytotoxicity of *C. rotundus* extract on normal cell lines (Vero cell line) as measured by the MTT assay.

No.	Name of sample	Concentration ( $\mu\text{g/ml}$ )	Absorbance	Inhibition (%) $\pm$ SD	IC <sub>50</sub> ( $\mu\text{g/ml}$ )
1	<i>C. rotundus</i> (whole plant)	500	0.98	40.39 $\pm$ 0.07	> 100
		250	1.04	30.80 $\pm$ 0.09	
		125	1.34	16.84 $\pm$ 0.01	
2	*Control		0.09	95.96 $\pm$ 0.01	

Key: \*Control = Triton-x100 was used as the control positive at 0.2  $\mu\text{g/ml}$ .

related pharmacological studies such as *in vivo* investigation, drug formulation and clinical trials are highly recommended.

## ACKNOWLEDGEMENTS

The authors are grateful to Prof. Dr. Aisha Zohair Al-Magboul, Department of Microbiology and Parasitology, Medicinal and Aromatic Plants and Traditional Medicine Research Institute (MAPTMRI) Khartoum, Sudan.

## REFERENCES

- Ahmed M, Mahayrookh, Mehjabeen, Rehman AB, Jahan N, **2012**. Analgesic, antimicrobial and cytotoxic effect of *Cyperus rotundus* ethanol extract. Pak J Pharmacol, 29(2):7-13.
- Ali H, Konig GM, Khalid SA, Wright AD, Kaminsky R, **2002**. Evaluation of selected Sudanese medicinal plants for their *in vitro* activity against *hemoflagellates*, selected bacteria, HIV-1-RT and tyrosine kinase inhibitory, and for cytotoxicity. J Ethnopharmacol, 83:219-228.
- Amaral FMM, Ribeiro MNS, Barbosa-Filho JM, Reis AS, Nascimento FRF, Macedo RO, **2006**. Plants and chemical constituents with giardicidal activity. Braz J Pharmacogn, 16:696-720.
- Arguello-Garcia R, Cruz-suto M, Romero-Montoya L, Ortega-Pierres G, **2004**. Variability and variation in drug susceptibility among *Giardia*

- duodenalis* isolates and clones exposed to 5-nitromidazoles and benzimidazoles *in vitro*. J. Antimicrob. Chemother, 54:711-721.
- Bhopale K, Pradhan K, Masani K, Karl C, 1995.** Additive effect of diloxanidefuroate and Metronidazole (Entamizole R) in mouse caecal amoebiasis. Indian J Exp Biol, 33:73-74.
- Cedillo-Rivera R, Chave B, Gonzalez-Robles A, Tapia-Contreras A, Yopez-Mulia L, 2002.** *In vitro* effect of nitazoxanide against *Entamoeba histolytica*, *Giardia lamblia* and *Trichomonas vaginalis* trophozoites. J Eukaryotic Microbiol, 49:201-208.
- Dang GK., Parekar RR, Kamat SK, Scindia AM, Rege NN, 2011.** Anti-inflammatory activity of *Phyllanthus emblica*, *Plumbago zeylanica* and *Cyperus rotundus* in acute models of inflammation. Phytother Res, 25(6):904-908.
- Dhillon RS, Singh S, Kundra S, Basra AS, 1993.** Studies on the chemical composition and biological activity of essential oil from *Cyperus rotundus* Linn. Plant Growth Regul, 13:89-93.
- Durate MCT, Figueira GM, Sartoratto A, Rehder VLG, Delarmelina C, 2005.** Anti-Candida activity of Brazilian medicinal plant. J Ethnopharmacol, 97:305-311.
- El-Ghazali GEB, El-Tohami MS, El-Egami AAB, 1994.** Medicinal plants of the White Nile provinces. Medicinal plants of the Sudan.
- El-Kamali HH, El-Khalifa KF, 1999.** Folk medicinal plants of riverside forests of the Southern Blue Nile district, Sudan. Fitoterapia, 70:493-497.
- Farthing MS, Cavellos AM, Kelly P, Cook GC, 1996.** Intestinal Protozoa; In Manson's Tropical Disease. 20<sup>th</sup> Edition, London W.B. Saunder Company. 1255-1267.
- Govaerts R, Simpson DA, Goetghebeur P, Wilson K, Egorova T, Bruhl JJ, 2007.** World checklist of Cyperaceae. The Board of Trustees of the Royal Botanic Gardens, Kew. Available at <http://www.kew.org/wcsp/monocots/> Accessed on 1 October 2007.
- Gupta MB, Palit TK, Singh N, Bhargava KP, 1971.** Pharmacological studies to isolate the active constituents from *Cyperus rotundus* possessing anti-inflammatory, anti-pyretic and analgesic activities. Ind J Med Res, 59:76-82.
- Haaksma AA, Jansen BJM, de Groot A, 1992.** Lewis acid catalysed DielsAlder reactions of S-(+)-carvone with silyloxy dienes. Total synthesis of (+)-a-cyperone. Tetrahedron, 48:3121-3130.
- Haque R, Mollah NU, Ali IK, Alam K, Eubanks A, Lyerly D, Petri Jr WA, 2000.** Diagnosis of amebic liver abscess and intestinal infection with the TechLab *Entamoeba histolytica* II antigen detection and antibody tests. J Clin Microbiol, 38(9):3235-3239.
- Harbone B, 1984.** Phytochemical methods. 2nd. New York, Champan., Hall, pp.4, 4-7.
- Hikino H, Ito K, Aota K, 1966.** Structure and absolute configuration of cyperotundone. Chem Pharmaceut Bullet, 14:890-899.
- Hikino H, Takemoto T, 1967.** Structure and absolute configuration of cyperol and isocyperol. Chem Pharmaceut Bullet, 15(12):1929-1933.
- Howe R, Mc Quillin FJ, 1955.** The structure of cyperone. Part IV. The synthesis of natural (+)-a-cyperone, its enantiomorph and epimer. J Chem Soc, 2423-2428.
- Huang KC, 1999.** The dried tuber of *Cyperus rotundus* L. The Pharmacology of Chinese Herbs 2nd Ed., pp.320-321.
- Joshi AR, Joshi K, 2000.** Indigenous knowledge and uses of medicinal plants by local communities of the Kali Gandaki Watershed Area, Nepal. J Ethnopharmacol, 73:175-183.
- Joulain D, Konig WA, 1998.** The Atlas of Spectral Data of Sesquiterpene Hydrocarbons. E. B.-Verlag, Hamburg.
- Kilani S, Ben Ammar R, Bouhlel I, Abdelwahed A, Hayder N, Mahmoud A, Ghedira K, Chekir-Ghedira L, 2005.** Investigation of extracts from (Tunisian) *Cyperus rotundus* as antimutagens and radical scavengers. Environ Toxicol Pharmacol, 20:478-484.
- Kilani S, Bouhlel I, Ben Ammar R, Ben Sghair M, Skandrani I, Boubaker J, Mahmoud A, Dijoux-Franca MG, Ghedira K, Chekir-Ghedira L, 2007.** Chemical investigation of different extracts and essential oil from the tubers of (Tunisian) *Cyperus rotundus*. Correlation with their antiradical and antimutagenic properties. Ann Microbiol, 57:657-664.
- Kilani S, Ledauphin J, Bouhlel I, Ben Sghair M, Boubaker J, Skandrani I, Mosrati R, Ghedira K, Barillier D, Chekir-Ghedira L, 2008.** Comparative study of *Cyperus rotundus* essential oil by a modified GC/MS analysis method. Evaluation of its antioxidant, cytotoxic, and apoptotic effects. Chem Biodivers, 5:729-742.
- Koko SW, 2005.** Antimalarial activity of *Xanthium brasiliicum* Vell\*. *In vitro*, *In vivo* and toxicological approaches. Recent Prog Med Plants, 15:1-10.
- Koko WS, Mesaik MA, Yousaf S, Galal M, Choudhary MI, 2008.** *In vitro* immunomodulating properties of selected Sudanese medicinal plants. J. Ethnopharmacol, 118:26-34.
- Nandihalli LE, Bendixen UB, 1987.** Worldwide distribution of purple and yellow (*Cyperus rotundus* and *C. esculentus*). Weed Technol, 1(1):61-65.
- Neffatti A, Ben Ammar R, Dijoux-Franca MG, Ghedira K, Chekir-Ghedira L, 2008.** *In vitro* evaluation of antibacterial, antioxidant, cytotoxic and apoptotic activities of the tubers infusion and extracts of *Cyperus rotundus*. Bioresour Technol, 99:9004-9008.
- Nyasse B, Ghogumu Tih R, Sodengam BL, Martin MT, Bodo B, 1988.** Mandassidione and other sesquiterpenic ketones from *Cyperus articulatus*. Phytochemistry, 27:3319-3321.
- Oliver-Bever B, 1986.** Medicinal Plants in Tropical West Africa. Cambridge University Press: Cambridge, UK. pp 200.
- Pal DK, Dutta S, 2006.** Evaluation of the antioxidant activity of the roots and rhizomes of *Cyperus rotundus* L. Ind J Pharm Sci, 68:256-258.
- Patel S, Gheewala N, Suthar A, Shah A, 2009.** *In-vitro* cytotoxicity activity of *Solanum nigrum* extract against Hela cell line And Vero cell line. Int J Pharm Pharmaceut Sci, 1(1):38-46.
- Puratuchikody A, Nithya DC, Nagalakshmi G, 2006.** Wound healing activity of *Cyperus rotundus* Linn. Indian J. Pharm. Sci., 68:97-101.
- Raut NA, Gaikwad NJ, 2006.** Antidiabetic activity of hydro-ethanolic extract of *Cyperus rotundus* in alloxan induced diabetes in rats. Fitoterapia, 77:585-588.
- Ravdin JI, Stauffer MM, 2005.** *Entamoeba histolytica* (Amoebiasis). In: Mendell, G. L., Benneth, J. E. Dolin, R. (ed) Mendell, Doglas and Benneth) Principles and Practice of Infectious Diseases. 6th ed. Philadelphia, P. A. Churchill Livingstone.
- Samia HA, Elmaliq KH, Khalid HS, Ashamat AM, Khojali SME, 2004.** Biochemical changes in rates experimentally infected with *Trypanosoma evansi*. J Anim Vet Adv, 3(7):483-486.
- Seo WG, Pae HO, Oh GS, Chai KY, Kwon TO, Yun YG, Kim NY, Chung HT, 2001.** Inhibitory effects of methanol extract of *Cyperus rotundus* rhizomes on nitric oxide and superoxide productions by murine macrophage cell line, RAW 264.7 cells. Ethno Pharmacol, 76:59-64.
- Stanley Jr SL, Reed SL, 2001.** Microbes and microbial toxin: paradigms for microbial-mucosal. Interactions VI. *Entamoeba histolytica*: parasite/hosts interactions. Am J Physiol Gastrointest Liver physiol, 280(6):1049-1054.
- Sundaram MS, Sivakumar T, Balamurugan G, 2008.** Anti-inflammatory effect of *Cyperus rotundus* Linn. Leaves on acute and subacute inflammation in experimental rat models. Biomedicine, 28:302-304.
- Thebtaranonth C, Thebtaranont Y, 1995.** Antimalarial sesquiterpenes from tubers of *Cyperus rotundus*: structure of 10,12-peroxycalamenene, a sesquiterpene endoperoxide. Phytochemistry, 40(1):125-128.
- Tierney LM, Mcphee JS, Papadakis MA, 1998.** Current medical diagnosis and treatment Appleton and Lange (Ed.), Stanford, Conn., 344-353.
- Townson SM, Boreham PF, Upcroft P, Upcroft JA, 1994.** Resistance to the nitroheterocyclic drugs. Acta Trop., 56:173-194.
- Uddin SJ, Mondal K, Shilpi JA, Rahman MT, 2006.** Antidiarrhoeal activity of *Cyperus rotundus*. Fitoterapia, 77(2):134-136.
- Umeric SC, Ezeuzo HO, 2000.** Physicochemical characterization and utilization of *Cyperus rotundus* starch. Bioresour Technol, 72:193-196.
- Upcrof JA, Dunn LA, Wriugh JM, Benakli K, Upcrof P, Vanelle P, 2006.** 5-nitroimidazole drugs effective against Metronidazole-resistant *Trichomonas vaginalis* and *Giardia dodonalis* spp. Antimicrob Agent Chemother, 50:344-347.
- Weenen H, Nkunya MH, Bray DH, Mwasumbi LB, Kinabo LS, Kilimali VA, 1999.** Antimalarial activity of Tanzanian medicinal plants. Planta Medica, 56:368-370.
- Won-Gil S, Hyun OP, Gi-Su O, Kyu-Yun C, Tae-Oh K, Young-Gab Y, Na-Young K, Hun-Taeg C, 2001.** Inhibitory effects of methanol extract of *Cyperus rotundus* rhizome son nitric oxide and superoxide

- productions by murine macrophage cell line, RAW 264.7 cells. J Ethno Pharmacol, 76(1):59–64.
- World Health Organization (WHO), 1997. Weekly Epidemiological Records, WHO 72:97-100.
- Yu J, Lei G, Cai L, Zou Y, 2004. Chemical composition of *C. rotundus* extract. J Phytochem, 65:881-889.
- Zeid AN, Majid SJ, Raghidah IW, Huda AAH, 2008. Extraction, identification and antibacterial activity of *Cyperus* oil from Iraqi *C. rotundus*. Eng Technol, 26(10):1156.
- Zhu M, Luk HH, Fung HS, and Luk CT, 1997. Cytoprotective effects of *Cyperus rotundus* against ethanol induced gastric ulceration in rats. Phytother Res, 11: 392–394.

---

---

**Citation:** Kabbashi AS, Osman EE, Abdrabo AM, Abuzeid N, Garbi MI, Koko WS, Dahab MM, 2015. Antiamoebic activity and cytotoxicity of ethanolic extract of *Cyperus rotundus* L.. Adv Med Plant Res, 3(4): 155-161.

---

---