



Contents lists available at ScienceDirect

Asian Pacific Journal of Tropical Medicine

journal homepage: www.elsevier.com/locate/apjtm

Document heading doi:

Design of a multiplex PCR method for detection of toxigenic–pathogenic in *Vibrio cholerae*

Imani Fooladi AA^{1*}, Iman Islamieh D¹, Hosseini Doust R², Karami A³, Marashi SM⁴

¹Applied Microbiology Research Center, Baqiyatallah University of Medical Sciences, Tehran, Iran

²Department of Microbiology, Faculty of Advance Sciences and Technology, Islamic Azad University–Pharmaceutical Sciences Branch, Tehran, Iran

³Molecular Biology Research Center, Baqiyatallah University of Medical Sciences, Tehran, Iran

⁴Department of Microbiology and Immunology, Babol University of Medical Sciences, Babol, Iran

ARTICLE INFO

Article history:

Received 10 October 2012

Received in revised form 15 November 2012

Accepted 15 December 2012

Available online 20 February 2013

Keywords:

Vibrio cholerae

Multiplex PCR

Rapid detection

ABSTRACT

Objective: To study virulence and regulatory genes (*hlyA*, *ctxB*, *tcpI*) in clinical strains of *Vibrio cholerae* (*V. cholerae*), simultaneously. **Methods:** Three important genes, *tcpI*, *hlyA* and *ctxB* were used for detection of toxigenic and pathogenic *V. cholera* by chain reaction assay method. **Results:** According to the results of the PCR, the incidence of *hlyA*, *tcpI*, and *ctxB* genes in clinical isolates was obtained as 94.7% (72 sample), 90.8% (69 sample), and 92.1% (70 sample), respectively. Five strains possessed all genes except *ctxB*, six strains possessed all genes except *tcpI*, four strains possessed all genes except *hlyA*, one strain possessed only *hlyA* and 60 strains contained a combination of three genes, including *hlyA*, *ctxB* and *tcpI*. **Conclusions:** Result show that this method could be reliable to detect toxigenic–pathogenic strains of *V. cholerae* in Iran.

1. Introduction

Cholera is a severe diarrhea that is caused by *Vibrio cholerae* (*V. cholerae*)^[1–3]. affects in an important way on the health and economy of the people^[1,3,4]. Information about presence of this bacterium in clinical samples can help in properly preventive actions^[1].

Additional, there are a high amounts of non molecular microbiological methods that has been used for detection of *V. cholerae*^[1] which have the disadvantages of laborious and boring^[1,3,4]. Since cholera detection requires urgent testing^[5], therefore applied molecular methods that are confident and rapid are needed^[1,4]. Among methods based DNA, monoplex PCR is imperfect because it don't give concurrent detection of the toxigenic–pathogenic potential and regulating factors^[3,6]. All these reasons have been encouraging us to explore rapid and affordable detection of

toxigenic and nontoxigenic *V. cholerae* in clinical samples and, consequently, estuarine environments and food products.

This organism enters in small intestine and produces cholera toxin (CT)^[7–9]. Toxin co-regulated toxin (TCP), related to *Vibrio* pathogenicity island (VPI)^[9], encode the *tcpA* gene^[2,10]. *tcpA* acts as a receptor for CTX Φ ^[1,9] and colonization of the gut is facilitated due to presence of *tcpA* on the surface of the bacterium^[2,7,8,10]. Therefore, any strain possessing *tcpA* gene is regarded as pathogenic and it can become toxigenic after obtaining the CTX cassette gene^[1]. On the other hands, *tcpI*—other member in VPI— is an integral inner membrane protein, engaged as environmental sensing and signal transduction and reversely regulates the synthesis of the *tcpA*^[7]. Hemolysin (*hlyA*) too is another virulence factor that with high frequency found in toxigenic and non-toxigenic strains^[8] and it causes diarrhea^[8,11].

In this research, we studied virulence and regulatory genes (*hlyA*, *ctxB*, *tcpI*) in clinical strains of *V. cholerae*, simultaneously.

*Corresponding author: Abbas Ali Imani Fooladi, Applied Microbiology, Research Center, Baqiyatallah University Of Medical Sciences, Tehran, Iran.

Tel / Fax :00982188039883

E-mail: imanifooladi.a@bmsu.ac.ir or imanifooladi.a@gmail.com

2. Material and methods

2.1. Sampling

A total of 76 stool samples were collected from different patients in diverse provinces of Iran such as Sistan and Baluchestan, Ghom, Tehran, Ghazvin, Kurdistan and Golestan during epidemics, between 2005 and 2010.

2.2. Identification and serogrouping of *V. cholerae*

First, samples were enriched in alkaline peptone water (APW). The tubes were then shaken (100 rpm) at 37 °C for 6 to 8 h. Approximately 5 µL of enriched APW broth was streaked onto TCBS agar (Merck, Germany) and incubated at 37 °C for 18–24 h. All the yellow colonies on TCBS plates were selected and assayed by biochemical tests including oxidase, sucrose, mannose and arabinose fermentation, motility, indol, methyl red Voges–Proskauer & resistant to poly–myxin B. Clinical isolates of the *V. cholerae* were serogrouped by using growth from TSI agar slants and slide agglutination test with polyvalent O₁, O₁₃₉ Inaba and Ogawa antisera (Mast Diagnostics Ltd., Bootle, Mersey side, UK).

Afterward, all strains were suspended in BHI broth (Pronadisa, Spain) supplemented with 30% glycerol and stored at –70 °C^[12].

2.3. DNA samples were prepared as follow

Before gene extraction, frozen store strains were subcultured streaked onto LB agar (Pronadisa, Spain) and then onto TCBS agar to confirm purity. For DNA extraction Bidinost *et al* method^[13] was modified and used.

2.4. Primers design

The oligonucleotide primers for *ctxB* and *tcpI* were designed based on existing GenBank sequences for all strains of *V. cholerae* by allele ID 6 software and oligonucleotide primer for *hlyA* was obtained of Rivera *et al* study^[7].

2.5. Chain reaction assay (PCR)

PCR methods were carried out using *V. cholerae* 62013 as standard strain and clinical samples were tested subsequently. *V. cholerae* 62013 is a toxigenic and pathogenic *V. cholerae* strain^[16,17].

In the first stage, primers were evaluated by uniplex PCR, When verified the correct functioning of each set, all primers were assayed together by multiplex PCR. One isolate of *V. cholerae* 62013 was used as the PCR positive control for *ctxB*, *hlyA*, *tcpI*, and *Salmonella typhi*, *Shigella dysenteriae*, *Escherichia coli* O₁₅₇:H₇ were used as negative controls.

PCR was carried out in 25 µL containing 2.5 µL of 10× PCR amplification buffer (500 mM KCl; 100 mM Tris–HCl pH 9.0; 1% Triton X–100), 1 µL dNTPs (2.5 mM each), 1 µL

(10 pmol) of each of the primers, 1 µL of MgCl₂ 50 mM, 0.20 µL of *Taq* DNA polymerase(1U) (Cinagen. Iran) and 5 µL of DNA template and double distilled water to a final volume of 25 µL. The PCR circumstances for all the reactions consisted of a first step of 5 min at 94 °C and a final step of 5 min at 72 °C. The middle step included 35 cycles of denaturation, annealing and extension times for: a) *ctxB*: 1 min at 94 °C, 1 min at 60 °C and 1 min at 72 °C; b) *hlyA*: 1 min at 94 °C, 1 min at 62 °C and 1 min at 72 °C. c) *tcpI*: 1 min at 94 °C, 1 min at 60 °C and 1 min at 72 °C respectively.

Amplified products (5 µL) were analyzed based on the appoint of their sizes by 1.5% agarose gel electrophoresis in TAE buffer. A 100 base pair ladder (Fermentas, Germany) was used as molecular size standard, Stained with ethidium bromide, and visualized under a Bio–Rad UV transilluminator (Hercules, CA, USA).

Relative bands from three different PCRs were extracted and purified from agarose gel (Cinagen. Iran) using the DNA extraction kit (Fermentas, Germany) and sequenced (Macrogen Research, Seoul, Korea).

2.6. Multiplex PCR assay

Briefly, all three primer pairs were added to the PCR mixture of a standard strain. The following reagents were added to achieve each 25 of PCR mixture: 2.5 µL of 10 × PCR amplification buffer (500 mM KCl; 100 mM Tris–HCl pH 9.0; 1% Triton X–100), 1 µL of MgCl₂ 50 mM, 1 µL dNTPs, 0.3 µL (10 pmol) of each of forward and reverse primers, 0.2 µL (1U) of *Taq* DNA polymerase and 5 µL of DNA template and double distilled water to a final volume of 25 µL. The automated thermal cycler (Master cycler gradient, Eppendorf, Germany) was programmed as follows: initial denaturation at 94 °C for 5 min, followed by 35 cycles consisting of 94 °C for 1 min, annealing at 58 °C for 1 min, and extension at 72 °C for 1 min and a final extension phase of 72 °C for 5 min. The PCR products were separated by 1.5% agarose gel and stained with ethidium bromide. The agarose gel was imaged with a Bio–Rad UV transilluminator.

2.7. Specificity of multiplex PCR technique for *ctxB*, *tcpI* and *hlyA* genes

To confirm the specificity of this study primers, we used 50 pg of genomic DNA of standard strain of *V. cholera* and 3 strains of other digestive system pathogenic bacteria's (*Salmonella typhi*, *Shigella dysenteriae*, and *Escherichia coli* O₁₅₇:H₇).

Primers applied in this research were specific and no amplification was seen with other strains of *Salmonella typhi*, *Shigella dysenteriae* and *Escherichia coli* O₁₅₇:H₇ and nonspecific bands were not seen .

2.8. Sensitivity of multiplex PCR technique for *ctxB*, *tcpI* and *hlyA* genes

Performing sensitivity test and determine the detection

Table 1

Primers characterization applied for this study.

Target genes	Nucleotide sequences 5'–3'	Accession no.	Tm ^a (°C)	Amplicon size(bp)	Source or reference
<i>ctxB</i> –F	ATG AGG CGT TTT ATT ATT CCA TAC AC	HQ599508.1	57.5	128	Original
<i>ctxB</i> –R	TAC CAG GTA GTC AAC ATA TAG ATT CA				
<i>tcpI</i> –F	TGC GTG ATG CTA ATT GGA CT	L25659.1	60.4	444	Original
<i>tcpI</i> –R	TTC GGT TTG TTT GCT TGA TG	–	–	–	–
<i>hlyA</i> –F	GGC AAA CAG CGA AAC AAA TAC C	M36855.1	59.0	738/727 (ET/Clas)	15
<i>hlyA</i> –R	CTC AGC GGG CTA ATA CGG TTT A	–	–	–	–

Tm^a, melting temperature.

limit for *V. cholerae* can define the strength of this test. The sensitivity of multiplex PCR assay was tested with standard strain of *V. cholerae* O₁. Standard strain was grown overnight in a BHI broth. Genomic DNA was extracted by the method described above and tenfold serial dilutions were prepared from 100 ng to 10 pg, and multiplex–PCR was carried out at each concentration.

3. Results

Biochemical and serological assays defined that all isolates belongs to the *V. cholerae* O₁ category. Extracting gene process was confirmed through observation of appropriate bands in gel agarose 1%. The simple and multiplex PCR products gained for each gene studied were shown in Figure 1 and size of each amplicon was confirmed. Primers applied in this research were specific and no amplification was seen with other strains of *Salmonella typhi*, *Shigella dysenteriae* and *Salmonella coli* O₁₅₇:H₇ and nonspecific bands were not seen (Figure 1).The sensitivity or minimum concentration of detection of *ctxB*, *tcpI*, and *hlyA* genes for O₁ strain were calculated 5, 50, and 50 pg, respectively.




Figure 1. Uniplex and Multiplex PCR for detection of virulence-associated genes in clinical strains of *V. cholerae* O₁ biotype El Tor. Lane 1, 2, 3, 7 multiplex PCR for the two clinical samples, positive and negative controls strains respectively, lane 4, 5, 6 uniplex PCR for *ctxB*, *tcpI* and *hlyA*, respectively, lane 8 molecular weight marker 100–bp DNA.

According to the results of the PCR, the incidence of *hlyA*, *tcpI*, and *ctxB* genes in clinical isolates was obtained as 94.7% (72 strains), 90.8% (69 strains), and 92.1% (70 strains), respectively. Five strains possessed all genes except *ctxB*, six strains possessed all genes except *tcpI*, four strains possessed all genes except *hlyA*, one strain possessed only *hlyA* and 60 strains contained a combination of three genes, including *hlyA*, *ctxB* and *tcpI* (Table 2). The *tcpI* and *hlyA* gene were frequently detected in toxigenic *V. cholera* O₁ serogroup (84.21% and (86.84%) respectively. The majority genotype observed in these samples was *ctxB hlyA ET tcpI* (78.95%).

4. Discussion

Formerly, many studies has been carried out with multiplex PCR method for study of virulence factors in *V. cholera*, but many of those studies were detecting virulence genes that are not frequently detected[6,8] or many of them were not searching toxicity and pathogenicity simultaneously[18]. Moreover, many of these methods are time consuming and unspecified.

Among diverse toxins generated by *V. cholerae*, cholera toxin is the most strong but a number of other genes such as *hlyA* and *tcpI* are concerned with cholera pathogenesis[1]. The sequence of *hlyA* gene in the classical biotype has an 11–bp deletion, contrasted to the El Tor biotype[7,11]. Despite exist differences in the nucleotide sequences of *hlyA* in El Tor and classical biotypes, the oligonucleotide primer applied in this study had been chosen from the fragment of the gene possessing the consensus sequence, make possible the detection of both biotypes[15]. Recently in Iran, Fallah and his group[5] applied multiplex PCR to detect *V. cholera* O₁ and non O₁ using the *ctxA*, *tcpA* and *ompW* genes on standard strains. All target genes were detected in *V. cholera* O₁ but only *ompW* gene was detected in *V. cholera* non O₁. Moreover, non–clinical samples were analyzed.

In other PCR multiplex studies that had been done on a variety of serogroups of *V. cholerae* by other researchers, these results were reported. In many studies, *hlyA* gene was found in all toxigenic and non toxigenic *V. cholerae* O₁ and O₁₃₉[7,8,11,14]. The presence of *hlyA* (100%) gene in non toxigenic isolates of our study with high frequency in above

studies, prove these genes can be used for detection of non toxigenic serogroups. The *tcpI* gene was reported in studies of Singh *et al*[11] and Rivera *et al*[7] with frequency of 100% and 84.2% for *V. cholerae* O₁ serogroup respectively.

Also, *ctx* gene frequency of our study (92.1%) was approximately equal acquired results of similar studies with O₁ and O₁₃₉ serogroups, for example, frequency of this gene in studies has been reported equally, 89.7%, 94.7% and 96.7%, respectively[7,11,19].

In the present study the results of acquired with Multiplex PCR were compared to monoplex PCR in 76 clinical isolates of the *V. cholerae* O₁. This analysis determined that there was perfect homology between these methods. Although six strains were belonging to the O₁ serogroup but don't containing the *ctx* gene cluster. Although, the *hlyA* gene used in this study has a common sequence in *V. cholerae* O₁, O₁₃₉, non-O₁ and non-O₁₃₉ serogroups[15], but since all isolates studied in this research had belonged to O₁ serogroup, and as regards classical biotype to be extinct nowadays[14, 20] therefore it can be concluded that strains possessor these gene (72 strains) are belonging to biotype Eltor. In addition, all these strains were Voges–Proskauer positive and resistant to polymixin B and these can confirm our hypothesis. We believe that our study is the first report for rapid detection of these genes by multiplex PCR from clinical isolates of *V. cholerae* in Iran.

Conflict of interest statement

We declare that we have no conflict of interest.

References

- [1] Goel AK, Ponmariappan S, Kamboj DV, Singh L. Single multiplex polymerase chain reaction for environmental surveillance of toxigenic–pathogenic O₁ and non-O₁ *Vibrio cholerae*. *Folia Microbiol (Praha)* 2007; **52**: 81–85.
- [2] Kumar P, Peter WA, Thomas S. Rapid detection of virulence-associated genes in environmental strains of *Vibrio cholerae* by multiplex PCR. *Curr Microbiol* 2010; **60**: 199–202.
- [3] Khuntia HK, Pal BB, Chhotray GP. Quadruplex PCR for simultaneous detection of serotype, biotype, toxigenic potential, and central regulating factor of *Vibrio cholerae*. *J Clin Microbiol* 2008; **46**: 2399–401.
- [4] Espineira M, Atanassova M, Vieites JM, Santaclara FJ. Validation of a method for the detection of five species, serogroups, biotypes and virulence factors of *Vibrio* by multiplex PCR in fish and seafood. *Food Microbiol* 2010; **27**: 122–131.
- [5] Mehrabadi J, Morsali P, Rohani nejad H, Imani Fooladi AA, Najafy S. Evaluation of new primers for detecting toxigenic *Vibrio cholerae* by multiplex PCR. *Microbiol Res* 2011; **3**: 3.
- [6] Singh DV, Isac SR, Colwell RR. Development of a hexaplex PCR assay for rapid detection of virulence and regulatory genes in *Vibrio cholerae* and *Vibrio mimicus*. *J Clin Microbiol* 2002; **40**: 4321–434.
- [7] Rivera IN, Chun J, Huq A, Sack RB, Colwell RR. Genotypes associated with virulence in environmental isolates of *Vibrio cholerae*. *Appl Environ Microbiol* 2001; **67**: 2421–249.
- [8] Kumar P, Peter WA, Thomas S. Detection of virulence genes in *Vibrio cholerae* isolated from aquatic environment in Kerala, Southern India. *Appl Biochem Biotechnol* 2008; **151**: 256–62.
- [9] Sarkar A, Nandy RK, Nair GB, Ghose AC. *Vibrio* pathogenicity island and cholera toxin genetic element-associated virulence genes and their expression in non-O₁ non-O₁₃₉ strains of *Vibrio cholerae*. *Infect Immun* 2002; **70**: 4735–4742.
- [10] Sharma A, Chaturvedi AN. Prevalence of virulence genes (*ctxA*, *stn*, *OmpW* and *tcpA*) among non-O₁ *Vibrio cholerae* isolated from fresh water environment. *Int J Hyg Environ Health* 2006; **209**: 521–526.
- [11] Singh DV, Matte MH, Matte GR. Molecular analysis of *Vibrio cholerae* O₁, O₁₃₉, non-O₁, and non-O₁₃₉ strains: clonal relationships between clinical and environmental isolates. *Appl Environ Microbiol* 2001; **67**: 910–921.
- [12] Islam MS, Mahmud ZH, Ansaruzzaman M. Phenotypic, genotypic, and antibiotic sensitivity patterns of strains isolated from the cholera epidemic in Zimbabwe. *J Clin Microbiol* 2011; **49**: 2325–2327.
- [13] Bidinost C, Saka HA, Aliendro O. Virulence factors of non-O₁ non-O₁₃₉ *Vibrio cholerae* isolated in Cordoba, Argentina. *Rev Argent Microbiol* 2004; **36**: 158–163.
- [14] Goel AK, Jain M, Kumar P, Jiang SC. Molecular characterization of *Vibrio cholerae* outbreak strains with altered El Tor biotype from southern India. *World J Microbiol Biotechnol* 2010; **26**: 281–287.
- [15] Panicker G, Call DR, Krug MJ, Bej AK. Detection of pathogenic *Vibrio* spp. in shellfish by using multiplex PCR and DNA microarrays. *Appl Environ Microbiol* 2004; **70**: 7436–7444.
- [16] Sharifnia A, Bakhshi B, Hashemi M, Pourshafie MR. Characterisation of diversity in wbeT region of the Oantigen biosynthesis gene cluster in Ogawa and Inaba serotypes of *Vibrio cholerae*. 19th European Congress of Clinical Microbiology and Infectious Diseases; 2009.
- [17] Zeighami H, Sattari M. Cloning and expression of a cholera toxin Beta subunit in *Escherichia coli*. Proceedings of the 3rd International Conference on Energy and Development –Environment–Biomedicine (EDEB '09), Athens, Greece; December 29–31, 2009.
- [18] Hoshino K, Yamasaki S, Mukhopadhyay AK. Development and evaluation of a multiplex PCR assay for rapid detection of toxigenic *Vibrio cholerae* O₁ and O₁₃₉. *FEMS Immunol Med Microbiol* 1998; **20**: 201–207.
- [19] Radu S, Vincent M, Apun K. Molecular characterization of *Vibrio cholerae* O₁ outbreak strains in Miri, Sarawak Malaysia. *Acta Trop* 2002; **83**: 169–176.
- [20] Safa A, Sultana J, Dac Cam P, Mwansa JC, Kong RY. *Vibrio cholerae* O₁ hybrid El Tor strains, Asia and Africa. *Emerg Infect Dis* 2008; **14**: 987–988.