



Contents lists available at ScienceDirect

## Asian Pacific Journal of Tropical Medicine

journal homepage: [www.elsevier.com/locate/apjtm](http://www.elsevier.com/locate/apjtm)

Document heading doi: 10.1016/S1995-7645(14)60088-2

# Prevalence of gastrointestinal helminthic infections and associated risk factors among schoolchildren in Tilili town, northwest Ethiopia

Alamneh Abera, Endalkachew Nibret\*

Biology Department, Bahir Dar University, P.O.Box 79, Bahir Dar, Ethiopia

## ARTICLE INFO

## Article history:

Received 6 January 2014

Received in revised form 15 March 2014

Accepted 28 May 2014

Available online 20 July 2014

## Keywords:

Intestinal helminthes

Prevalence

Risk factors

Tilili town

Ethiopia

## ABSTRACT

**Objective:** To determine the prevalence of gastrointestinal helminthes and associated risk factors among schoolchildren in Tilili town, northwest Ethiopia. **Methods:** A cross-sectional study involving 385 schoolchildren was conducted between November 2011 to February 2012. Each student was selected using systematic random sampling method. Questionnaire and observation were used to identify socio-demographic and associated risk factors. Fresh stool samples were observed using formal-ether concentration technique. Data were analyzed using SPSS 16.0 statistical software. **Results:** Four species of intestinal helminthes were identified with an overall prevalence of 44.2% (170 of 385 schoolchildren). The predominant parasites were *Ascaris lumbricoides* (*A. lumbricoides*) 153 (39.7%) and *Trichuris trichiura* (*T. trichiura*) 30 (7.8%). One hundred thirty five (35.1%) had single infections and 35 (9.2%) were infected with more than one helminthic parasites in which 32 (8.4%) were double infections and 3 (0.8%) were triple infections. Significant associations were observed between intestinal helminth infection and those of age, grade level, and school variables. Prevalence of hookworm infection was significant in children who did not wear shoes regularly ( $P < 0.05$ ). **Conclusions:** intervention programs and education on personal and environmental hygiene should be implemented for the prevention and control of helminthic infections in the study area.

## 1. Introduction

Intestinal helminthes are intestinal parasites that inhabit the human gastrointestinal tracts and are one of the most prevalent forms of parasitic disease causing organisms. The soil transmitted helminthes [*Ascaris lumbricoides* (*A. lumbricoides*), hookworm and *Trichuris trichiura* (*T. trichiura*)] are parasitic nematode worms causing human infection through contact with parasite eggs or larvae that thrive in the warm and moist soil of the world's tropical and subtropical countries[1]. They are one of the world's most important causes of physical and intellectual growth retardation.

Ascariasis caused by *A. lumbricoides* causes widespread morbidity due to a variety of medical and surgical

complications. It has been estimated that 1.471 billion cases of infection globally and 65 000 deaths occur due to *A. lumbricoides*[2, 3]. Hookworm in human is caused by helminthic parasites, *Necator americanus* (*N. americanus*) and *Ancylostoma duodenale* (*A. duodenale*) and is transmitted through contact with contaminated soil. It is one of the most common chronic infections with an estimated 1.3 billion cases globally and directly accountable for 65 000 deaths annually[4]. The two hookworm species that infect human exhibit differences in their pathogenicity, mode of transmission, geographical distribution and these differences may influence the morbidity of hookworm disease[5]. Clinical manifestations of hookworm disease are the consequences of chronic intestinal blood loss and iron deficiency anemia. In children, chronic hookworm disease retards physical growth which is sometimes most apparent at puberty[6]. Likewise, *A. lumbricoides* and hookworm, *T. trichiura* considerably affect the physical and mental

\*Corresponding author: Endalkachew Nibret, Biology Department, Bahir Dar University, P.O.Box 79, Bahir Dar, Ethiopia.  
E-mail: [endt2002@yahoo.com](mailto:endt2002@yahoo.com)

development in children[7]. About 1.049 billion cases of infection occurred due to *T. trichiura* and 70 000 deaths occur due to *T. trichiura* annually[8–10]. *Hymenolepis nana* (*H. nana*) infection has cosmopolitan distribution and most commonly infects humans, especially school-aged children (aged 4–10 years)[11].

Although several studies have been conducted on the distribution and prevalence of intestinal parasitic infections in Ethiopia, still there are several localities lacking epidemiological information on intestinal helminthic infections. Tilili town, located in northwestern Ethiopia, is one of such localities without report on gastrointestinal helminthic infections. Therefore, the objective of the present study was to assess the prevalence of gastrointestinal helminthic infections and associated risk factors among schoolchildren in the study area.

## 2. Materials and methods

### 2.1. Study design and area

A cross-sectional study was conducted in Tilili, town, northwest Ethiopia, from November 2011 to February 2012. Tilili is located at a distance of 420 km from Addis Ababa. The town is also located at 2 450 meter above sea level. The mean annual temperature ranges from 10–25 °C and the average 30 years annual rainfall is 1 396mm[12]. There are three elementary and junior secondary schools in Tilili town. The studied schools were Hibret, Tilili and Kili-Mesk elementary and junior secondary schools. In the town, the total student population of the schools was 4 431 (1 148 from Hibret, 1 871 from Kili-Mesk and 1 412 from Tilili) and among these 2 153 and 2 278 were male and female students, respectively.

The study population consisted of 403 schoolchildren and among these, 104, 129 and 170 children were selected from Hibret, Tilili and Kili-Mesk primary schools, respectively. The study population included all age groups, both sexes and students from both urban and rural settings. Students who took anthelmintic drugs at time of the study or two weeks prior to the study were excluded.

Questionnaire was used to gather information on socio-demographic and sanitary facilities of students. The study participants were also examined physically for the presence or absence of dirt materials in their fingernails. The sample size was determined using a general formula  $Z^2p(1-p)/d^2$  considering the level of significance at 5% and assuming the prevalence of intestinal helminthes in the study area to be 50%[13]. To lesson errors arising from likelihood of noncompliance or possible dropout, 5% of the sample size

was added to the normal sample size. Allocation of students was made proportional to the number of students for each school. To select the study subjects, the students were first stratified according to their educational level (grade 1 to 8) and they were taken from each class category by systematic random sampling using class roster as a sampling frame.

### 2.2. Ethical consideration and stool sample processing

Prior to stool sample collection, verbal consent was obtained from both schoolchildren and their guardians. Schoolchildren were guided on how to bring their stool samples and then they were provided with clean plastic stool vials with toilet tissue paper. After collecting sufficient fresh stool sample with plastic vial from each participant, codes were given for all on the plastic vials and then each child was interviewed using a structured questionnaire. All stool samples were preserved in 10% formalin solution. A specimen was then transported to Biomedical and Microbiology Laboratory, Biology Department, Bahir Dar University. Part of each individual stool sample was processed using formal-ether concentration method and examined under microscope to identify helminth eggs. The remaining part of the stool sample was preserved in 10% formalin solution for further reference.

### 2.3. Data analysis

The data collected through interview, questionnaire, and parasitological examination were analyzed using SPSS 16.0 statistical software. *Chi-square* ( $\chi^2$ ) test was used to observe the association between risk factors and prevalence of intestinal helminthes infections. Crude Odds Ratio (*COR*) and 95% confidence interval (*CI*) were performed to determine the strength of the association between infection and risk factors. *P*-values less than 0.05 were considered to be statistically significant.

## 3. Results

Out of 403 who were willing to participate in the study, only 385 students who took part in the study were included in the present analysis. Of these, 193 (50.1%) were males and 192 (49.9%) were females. The mean age of the study subject was 11.17 years old. According to their residence, 218 (56.6%) were from urban and 167 (43.4%) schoolchildren were from rural areas. The study population consisted of students from Hibret, Tilili and Kili-Mesk primary schools with 104 (27.0%), 122 (31.7%) and 159 (41.3%) students, respectively (Table 2).

Of the total 385 stool samples examined, 170 (44.2 %) were positive for one or more of intestinal helminthes. Among these, a total of 88 males and 82 females were positive for one or more helminth infections. Ninety five (43.6%) urban schoolchildren and seventy five (44.9%) rural schoolchildren were infected with one or more than one intestinal helminthes but, there was no significance difference between urban and rural dwellers ( $P>0.05$ ) (Table 2). The prevalence of intestinal helminthic infections were statistically significant ( $P<0.05$ ) among the schools (Table 2). Four major intestinal helminthes, namely, *A. lumbricoides*, *T. trichiura*, hookworm and *H. nana* were identified with

an overall prevalence of 44.2% in this study area. The most frequent helminth infection was *A. lumbricoides* (39.7%), and the least encountered helminth was *H. nana* (1%) (Table 3). Among 170 positive individuals, majority of the students (35.1%), had single infection. Thirty five (9.2%) of the students were infected with more than one intestinal helminth parasites, in which 32 (8.4%) were double intestinal helminthes infections and 3 (0.8%) were triple infections. Only female students were infected by triple infections (Table 1).

Among age groups, the most and the least affected age groups were 6–8 and 12–14 years old with a prevalence of 55.7% and 34.6%, respectively (Table 2).

**Table 1**

Prevalence of helminth infections among Tilili town primary school children, 2011/12.

Intestinal helminthes	No. of infected (%)	Male (%)	Female (%)
Single infection			
<i>A. lumbricoides</i>	121 (31.4)	69 (57.0)	52 (43.0)
<i>T. trichiura</i>	11 (2.9)	4 (36.4)	7 (63.6)
Hookworm	2 (0.5)	0 (0.0)	2 (100.0)
<i>H. nana</i>	1 (0.3)	1 (100.0)	0 (0.0)
Double infection			
<i>A. lumbricoides</i> + hookworm	15 (3.9)	6 (40.0)	9 (60.0)
<i>A. lumbricoides</i> + <i>T. trichiura</i>	13 (3.4)	7 (53.8)	6 (46.2)
<i>T. trichiura</i> + <i>H. nana</i>	2 (0.5)	1 (50.0)	1 (50.0)
<i>A. lumbricoides</i> + <i>H. nana</i>	1 (0.3)	0 (0.0)	1 (100.0)
Hookworm + <i>T. trichiura</i>	1 (0.3)	0 (0.0)	1 (100.0)
Triple infection			
<i>A. lumbricoides</i> + hookworm + <i>T. trichiura</i>	3 (0.8)	0 (0.0)	3 (100.0)
Total	170 (44.2)	88 (22.9)	82 (21.3)

**Table 2**

Overall prevalence of intestinal helminthic infections in relation to sex, age, residence, and school of children in Tilili town, 2011/12.

Characteristic	No of examined (%)	Positive(%)	Negative (%)	$\chi^2$ (P-value)
Sex				
Male	193 (50.1)	88 (45.6)	105 (54.4)	0.33 (0.568)
Female	192 (49.9)	82 (42.7)	110 (57.3)	
Age (years)				11.90 (0.008)
6–8	88 (22.3)	49 (55.7)	39 (44.3)	
9–11	113 (29.4)	48 (42.5)	65 (57.5)	
12–14	136 (35.3)	47 (34.6)	89 (65.4)	
15–18	48 (12.5)	26 (54.2)	22 (45.8)	
Residence				0.08 (0.790)
Urban	218 (56.6)	95 (43.6)	123 (56.4)	
Rural	167 (43.4)	75 (44.9)	92 (55.1)	
School of students				23.13 (0.000)
Hibret	104 (27.0)	61 (58.7)	43 (41.3)	
Tilili	122 (31.7)	61 (50.0)	61 (50.0)	
Kili–Mesk	159 (41.3)	48 (30.2)	111 (69.8)	

**Table 3**

Prevalence of helminth infections among sex, age groups, and school in Tilili town schoolchildren 2011/12.

Characteristics	No. of examined	Helminthes				P-value
		<i>A. lumbricoides</i> (%)	<i>T. trichiura</i> (%)	Hookworm (%)	<i>H. nana</i> (%)	
Sex						
Male	193 (50.1)	82 (42.5)	13 (6.7)	5 (2.6)	2 (1.0)	0.570
Female	192 (49.9)	71 (37.0)	17 (8.9)	16 (8.3)	2 (1.0)	
Age (years)						0.008
6–8	88 (22.9)	44 (50.0)	6 (6.8)	7 (7.9)	1 (1.1)	
9–11	113 (29.4)	42 (37.2)	12 (10.6)	5 (4.4)	1 (0.9)	
12–14	136 (35.3)	42 (30.9)	6 (4.4)	4 (2.9)	2 (1.5)	
15–18	48 (12.5)	25 (52.1)	6 (12.5)	5 (10.4)	0 (0.0)	
School						0.000
Hibret	104 (27.0)	58 (55.8)	7 (6.7)	10 (9.6)	0 (0.0)	
Tilili	122 (31.7)	56 (45.9)	13 (10.7)	7 (5.7)	0 (0.0)	
Kili–Mesk	159 (41.3)	39 (24.5)	10 (6.3)	4 (2.5)	4 (2.5)	
Total	385	153 (39.7)	30 (7.8)	21 (5.5)	4 (1.0)	

There was significant difference between intestinal helminthic infections with age-groups and school of students ( $P < 0.05$ ) (Table 3).

In univariate logistic regression analysis of factors potentially associated with gastrointestinal helminthic infections (Table 4), significant associations were observed between intestinal helminthic infections with family size, grade of students, hand washing habit after toilet, habit of cutting fingernail, dirty materials in fingernail and shoes wearing habit. The odds of being infected with gastrointestinal parasites for children from a family size of 7 and above was higher than those from a family size of less than 3 ( $COR: 1.85; 95\% CI: 0.29–1.98$ ). Children in grade level of 1–4 were more likely to be infected with gastrointestinal helminthes than those in grade level of 5–8 ( $COR: 1.66; 95\% CI: 1.08–2.55$ ). Children that did not wash their hands after using toilet were two times more likely to be infected than those who washed their hands ( $COR: 2.06; 95\% CI: 1.13–3.77$ ). Similarly, children who did not have the habit of cutting their fingernails were two times more likely to be infected than those who did cut their fingernails ( $COR: 2.30; 95\% CI: 1.16–4.57$ ). The odds of being infected with gastrointestinal helminthes in those children whose fingernails contained dirty materials were higher than those who did not have

dirty materials in their fingernails ( $COR: 1.63; 95\% CI: 1.02–2.60$ ). Those children who did not wear shoes regularly were almost two times more likely to be infected with hookworm than those who wore shoes regularly ( $COR: 1.89; 95\% CI: 1.13–3.17$ ).

#### 4. Discussion

In this study, the overall prevalence of intestinal helminth infection was 44.2%. It was a bit lower when compared to other studies which were done among schoolchildren in different countries. Prevalence of 53% from Oaxaca region of Mexico, and 47.0% from Mozambique were reported[14,15]. The difference in prevalence could be attributed to timing and seasonal differences in conducting the survey, environmental conditions and other geographical factors in the study areas. The prevalence of helminthic infection in the study area of current study was also relatively higher compared to other studies conducted on primary schoolchildren (32.0%) in western Tajikistan, in Ashanti region of Ghana (11.1%), and in Saudi Arabia (1.2%)[16–18]. The relatively higher rate of helminthic infection in schoolchildren of Tilili town might be due to the lack of hand washing habit after toilet, habit of

**Table 4**

Univariate logistic regression analysis for socio-demographic factors potentially associated with intestinal helminthic infections among Tilili town primary schoolchildren, 2011/12.

Risk factors		n	Positive (%)	Negative (%)	Crude odd ratio, 95% CI, P-value
Parents educational level	Illiterate	95	45 (47.4)	50 (52.6)	0.82, 0.38–1.79, 0.59
	Literate	150	64 (42.3)	86 (57.7)	1.47, 0.71–3.05, 0.26
	Completed	96	38 (39.6)	58 (60.4)	0.60, 0.27–1.30, 0.16
	Certificated and above	44	23 (52.3)	21 (47.7)	1.00
Occupation of parents	Merchants	154	60 (39.0)	94 (61.0)	0.98, 0.49–1.67, 0.73
	Farmers	153	76 (49.7)	77 (50.3)	1.40, 0.76–2.57, 0.25
	Others	8	5 (62.5)	3 (37.5)	2.36, 0.44–13.73, 0.25
	Government employees	70	29 (41.4)	41 (58.6)	1.00
Family size	7 and above	115	58 (50.4)	57 (49.6)	1.85, 0.29–1.98, 0.04*
	6	112	47 (42.0)	65 (58.0)	1.31, 0.69–2.50, 0.38
	5	82	38 (46.3)	44 (53.7)	1.57, 0.79–3.13, 0.17
	below 5	76	27 (35.5)	49 (64.5)	1.00
Grade of students	1–4	211	105 (49.8)	106 (50.2)	1.66, 1.08–2.55, 0.02*
	5–8	174	65 (37.4)	109 (62.6)	1.00
Availability of latrine	No	59	24 (40.7)	35 (59.3)	0.85, 0.46–1.54, 0.56
	Yes	326	146 (44.8)	180 (55.2)	1.00
Habit of using latrine	No	75	34 (45.3)	41 (54.7)	1.06, 0.62–1.82, 0.82
	Yes	310	136 (43.9)	174 (56.1)	1.00
Hand washing after toilet	No	59	35 (70.0)	24 (30.0)	2.06, 1.13–3.77, 0.01*
	Yes	326	135 (41.4)	191 (58.6)	1.00
Habit of cutting fingernails	No	45	28 (62.2)	17 (37.8)	2.30, 1.16–4.57, 0.01*
	Yes	340	142 (41.8)	198 (58.2)	1.00
Dirty materials in fingernails	Yes	263	126 (47.9)	137 (52.1)	1.63, 1.02–2.60, 0.03*
	No	122	44 (36.1)	78 (63.9)	1.00
Shoes wearing habit	No	85	48 (56.5)	37 (43.5)	1.89, 1.13–3.17, 0.01*
	Yes	301	123 (40.9)	178 (59.1)	1.00

\*Statistically significant at  $P < 0.05$ .

cutting fingernails, presence of dirty materials in fingernails, and shoes wearing habit.

The result of present study is also comparable with previous studies conducted in different parts of Ethiopia. High prevalence (92.7%) was recorded among schoolchildren at Lake Awassa<sup>[19]</sup>. The lower prevalence obtained in the present study in comparison with the above report could be attributed to availability and usage habit of latrine, bathing places and educational level of parents. Nevertheless, the percentage of helminthic infections in the current study was higher than other studies reported in Hintalo–Wajerat (26.53%) and in Babile town (27.2%)<sup>[20,21]</sup>. This might be due to availability of tap water for drinking, habit of cutting fingernails, family size and shoes wearing habit of students in those areas. However, the finding in current study area, Tilili town, was in agreement with other studies in Ethiopia, including eastern Ethiopia 45.9% and south Wello 43.3%<sup>[22,23]</sup>.

Among 170 students who were found positive for any helminthic infections, schoolchildren with single, double and triple infections were, 35.1%, 8.4% and 0.8%, respectively. It is common to find multiple infections per an individual in Ethiopia<sup>[24]</sup>. For example, 23.2% double and 5.3% triple infections were observed among school children in Zarima town, northwest Ethiopia<sup>[25]</sup>. In the current study, three common STH and one tapeworm (*H. nana*) infections were identified in 385 schoolchildren. The most frequent helminth infection was by *A. lumbricoides* followed by *T. trichiura* and hookworm infections with prevalence of 39.7%, 7.8%, and 5.5%, respectively. Similar result was reported from Abosa around Lake Ziway<sup>[26]</sup>. The least encountered helminth was *H. nana* with prevalence of 1%.

According to this study, there was significant association between grade level of students and rate of gastrointestinal helminthic infections. This was in contrast to other studies<sup>[27,28]</sup>. This finding might be due to lack of regular health education program in the school which can enhance their awareness in the transmission and control mechanisms. When we see the prevalence and associated risk factors of intestinal helminthes among schoolchildren in this study area, 84.7% of schoolchildren had toilet in their houses. But, the rate of intestinal helminth infection was not related to the presence or absence and usage of latrines ( $P>0.05$ ). Three hundred twenty six (84.7%) students had hand washing habit after toilet and there was significant difference between the rate of helminth infection and hand washing after toilet. Forty five (11.7%) students had not habit of cutting their fingernail regularly and among these, 62.2% were positive for one or more of helminth infections. There was a statistical significance difference between who did and did not cut their fingernails. This was probably due to lack of awareness,

poor hygiene practice and socio–demographic factor of schoolchildren. This result was also supported by earlier finding<sup>[29]</sup>. Two hundred sixty three (68.3%) student had dirty material in their right and left hand fingernails, out of these 47.9% were positive for one or more helminth infections. There was significant relation between helminth infection and dirty material in their fingernails. This is probably due to poor hygienic practice, socio–economic status and also playing habit of children with soil. This result was supported by previous study<sup>[30]</sup>.

Three hundred one (78.2%) children had shoes wearing habit. The absence of shoes wearing habit was the risk factor and statistically significant difference was observed between shoes wearing habit and intestinal parasitic infection. Those schoolchildren who had not shoes wearing habit were also more infected with hookworm than those schoolchildren who wore shoes regularly and this was statistically significant. This finding was similar to other studies<sup>[31,32]</sup>.

Provision of health education on transmission of hookworm to schoolchildren would motivate students to wear shoes regularly and thereby contributing a lot in the prevention and control of hookworm infection. Generally, the present findings in Tilili town strongly indicated the need for a comprehensive program to combat intestinal parasites and associated risk factors in the study area. Awareness creation by physicians, school teachers and mass media should be commenced promptly for preventing and controlling intestinal parasitic infections in general and helminthic infections in particular.

In conclusion, the current study has revealed a relatively high prevalence of intestinal helminthic infections among schoolchildren in Tilili town and this situation requires prompt implementation of interventional programs and education on personal and environmental hygiene. Physicians, school teachers and mass media should create awareness among schoolchildren and their parents for effective prevention and control of helminthic infections in the study area.

### Conflict of interest statement

We declare that we have no conflict of interest.

### References

- [1] Abdulhamid A, Hesham M, Johari S. Epidemiology of soil–transmitted helminthiasis in Malaysia. Reviewed. *Southeast Asian J Trop Med Public Health* 2011; **42**: 527–538.
- [2] Silva R, Brooker S, Hotez P, Montresor A, Engels D, Savioli L.

- Soil-transmitted helminth infections: updating the global picture. *Trends Parasitol* 2003; **19**: 547–551.
- [3] Drisdelle R. *Ascaris lumbricoides*: food borne pathogenic helminthes. *Intest Infect* 2007; **45**: 13–28.
- [4] Demissie F, Petros B, Kebede A. Hookworm species distribution among school children in Asendabo town Jimma zone, south Ethiopia. *Ethiop J Health Sci* 2008; **18**: 53–56.
- [5] Crompton D. The public health importance of hookworm disease. *Parasitology* 2000; **121**: 539–550.
- [6] Stephenson L, Latham M, Kury K, Brigham H. Treatment with a single dose of albendazole improves growth of Kenyan schoolchildren with hookworm, *T. trichiura* and *A. lumbricoides* infection. *Am J Trop Med Hyg* 1989; **41**: 78–87.
- [7] WHO. *Prevention and control of schistosomiasis and soil transmitted helminthiasis*. Geneva: WHO; 2002;
- [8] Chan M. The global burden of intestinal nematode infections—Fifty years on. *Parasitol Today* 1997; **13**: 438–443.
- [9] Ezeamama A, Friedman J, Acosta L, Bellinger D, Langdon G, Manalo D, et al. Helminth infection and cognitive impairment among Filipino children. *J Trop Med Hyg* 2005; **72**: 540–548.
- [10] Kwame D. *Infection disease: T. trichiura (Linnaeus, 1771)*. University of California; Los Angeles Medical Center; 2009.
- [11] King C. *Cestodes. Principles and practice of infectious diseases*. 6th ed. London: Elsevier Acad Press; 2005.
- [12] Bahir Dar Meteorology Branch Office. Ethiopian meteorology agency, Bahir Dar Branch; 2012.
- [13] Steven K. *Sampling*. 3rd ed. Hoboken: John Wiley and Sons Inc.; 2012.
- [14] Quihui L, Valencia M, Crompton D, Phillips S, Hagan P, Diaz S, et al. Prevalence and intensity of intestinal parasitic infections in relation to nutritional status in Mexican schoolchildren. *Trans R Soc Trop Med Hyg* 2004; **98**: 653–659.
- [15] Rassul N, Verónica C, Acácio S, Judite M. Geographic distribution and prevalence of schistosomiasis and soil transmitted helminthes among schoolchildren in Mozambique. *Am J Trop Med Hyg* 2009; **81**: 799–803.
- [16] Mathys B, Bobieva M, Karimova G, Mengliboeva Z, Jean R. Prevalence and risk factors of helminthes and intestinal protozoa infections among children from primary schools in western Tajikistan. *Parasit Vec* 2011; **4**: 195.
- [17] Sammy C, Gbedema Y, Gyampomah K. Accuracy of diagnosis of intestinal helminth parasites in a reference diagnostic laboratory in the Ashanti region of Ghana. *Int J Parasitol Res* 2011; **3**: 12–16.
- [18] Nagwa S, Magda M. Intestinal parasitic infection among children in the Kingdom of Saudi Arabia. *Aust J Basic Appl Sci* 2010; **4**: 4200–4204.
- [19] Merid Y, Hegazy M, Mekete G, Teklemariam S. Intestinal helminthic infection among children at Lake Awassa area, south Ethiopia. *Ethiop J Health Dev* 2001; **15**: 31–38.
- [20] Dejenie T, Asmelash T. *Schistosoma mansoni* among children of different water source user in Tigray, northern Ethiopia. *Momona Ethiop J Sci* 2010; **2**: 49–60.
- [21] Tadesse G. The prevalence of intestinal helminthic infections and association risk factors among school children in Babile town, Eastern Ethiopia. *Ethiop J Health Dev* 2005; **19**: 140–147.
- [22] Lo C, Ayele T, Birr H. Helminth and snail survey in Hargege region of Ethiopia with special reference to schistosomiasis. *Ethiop Med J* 1989; **27**: 73–83.
- [23] Tsehail A, Tilahun W, Amare D. Intestinal parasitism among students in three localities in south Wello, Ethiopia. *Ethiop J Health Dev* 1998; **12**: 231–235.
- [24] Legesse M, Erko B. Prevalence of intestinal parasites among school children rural area close to the southeast of Lake Langano, Ethiopia. *Ethiop J Health Dev* 2004; **18**: 116–120.
- [25] Alemu A, Atnafu A, Addis Z, Shiferaw Y, Teklu T, Mathewos B, et al. Soil transmitted helminthes and *S. mansoni* infections among school children in Zarima town, northwest Ethiopia. *BMC Infect Dis* 2011; **11**: 1–7.
- [26] Solomon G. Prevalence and associated risk factors of soil-transmitted helminthes among schoolchildren in Abosa around Lake Ziway, Southern Ethiopia. M. Sc thesis. Addis Ababa: Addis Ababa University; 2008.
- [27] Tshikuka J, Scott M, Gray-Donald K. *Ascaris lumbricoides* infection and environmental risk factors in an urban African setting. *Ann Trop Med Parasitol* 1995; **89**: 505–514.
- [28] Holland C, Asaolu S, Crompton D, Whitehead R, Coombs I. Targeted antihelminthic treatment of schoolchildren. *Parasitology* 1998; **113**: 87–95.
- [29] Ali I, Mekete G, Wodajo N. Intestinal parasitism and related risk factors among students of Asendabo elementary and junior secondary school, southwestern. *Ethiop J Health Dev* 1999; **13**: 157–161.
- [30] Pinar O, Sema E, Berna G, Ozlema O, Erdal B. Intestinal parasite prevalence and related factors in school children, Western city sample—Turkey. *BMC Public Health* 2004; **4**: 64.
- [31] Naish S, McCarthy J, Williams G. Prevalence, intensity and risk factors for soil transmitted helminth infection in a south Indian fishing village. *Acta Trop* 2004; **91**: 177–187.
- [32] Rebecca J, Ian D, Peter I, Norbert M, Andrew T. The prevalence, intensities and risk factors associated with geohelminth infection in tea-growing communities of Assam. *Indian Trop Med Int health* 2004; **9**: 688–701.