

HOSTED BY



ELSEVIER

Contents lists available at ScienceDirect

## Asian Pacific Journal of Tropical Medicine

journal homepage: <http://ees.elsevier.com/apjtm>Original research <http://dx.doi.org/10.1016/j.apjtm.2016.03.029>Antioxidant and hepatoprotective effects of *Ajuga nipponensis* extract by ultrasonic-assisted extractionChang-Wei Hsieh<sup>1,2</sup>, Wen-Ching Ko<sup>3</sup>, Wai-Jane Ho<sup>1</sup>, Chao-Kai Chang<sup>2</sup>, Guo-Jhou Chen<sup>3</sup>, Jen-Chieh Tsai<sup>1\*</sup><sup>1</sup>Department of Medicinal Botanicals and Health Applications, Da-Yeh University, 168 University Rd, Dacun, Chang-Hua, Taiwan, ROC<sup>2</sup>Biotechnology Research Center, Da-Yeh University, 168 University Rd, Dacun, Chang-Hua, Taiwan, ROC<sup>3</sup>Department of Bioindustry Technology, Da-Yeh University, 168 University Rd, Dacun, Chang-Hua, Taiwan, ROC

## ARTICLE INFO

## Article history:

Received 15 Jan 2016

Received in revised form 16 Feb 2016

Accepted 15 Mar 2016

Available online 23 Mar 2016

## Keywords:

*Ajuga nipponensis*

Ultrasonic-assisted extraction

Orthogonal test

Antioxidant

Hepatoprotection

## ABSTRACT

**Objective:** To investigate suitable condition for extraction of the active components from *Ajuga nipponensis* (*A. nipponensis*).**Methods:** Orthogonal experimental design was used to determine the optimal extraction parameters for ecdysterones and flavonoids. Finally, the hepatoprotective abilities of *A. nipponensis* extracts were evaluated by CCl<sub>4</sub>-induced animal models.**Results:** Maximum yields of flavonoids ( $7.87 \pm 0.10$ ) mg/g and ecdysterones ( $0.73 \pm 0.02$ ) mg/g could be obtained when the extraction time was 50 min, the extraction temperature was 60 °C, and the ratio of sample to 70% (v/v) ethanol was 1:20 (w/w). The antioxidant property of *A. nipponensis* was correlated to the concentration of its extracts. At 5 mg/mL, *A. nipponensis* extract scavenged 84.8% of DPPH radical and had absorbance values of  $2.43 \pm 0.04$  reducing power. Upon CCl<sub>4</sub>-induced liver injury, glutamic oxaloacetic transaminase and glutamic pyruvic transaminase decreased significantly after the mice were treated with *A. nipponensis*. Histological researches also explained that *A. nipponensis* reduced the extent of liver lesions induced by CCl<sub>4</sub>.**Conclusions:** *A. nipponensis* exhibited potent antioxidant activity in chemical experimental models and hepatoprotective effect against CCl<sub>4</sub>-induced liver damage.

## 1. Introduction

The liver is a vital organ in human which regulates metabolism, clears poisonous substances from blood and conversion of excess glucose to glycogen. Chronic liver failure affects many people globally with viral infection being the main cause. Nonetheless, alcohol and drugs abuse resulting in liver failure are also of high prevalence. Carbon tetrachloride (CCl<sub>4</sub>) and alcohol poisoning are commonly used to cause liver failure in animal models. The toxicity of CCl<sub>4</sub> depends on its reduction to form trichloromethyl

radicals via liver mitochondria, following by the production of more toxic trichloromethylperoxyl radical in the presence of oxygen [1]. On the other hand, ethanol feeding readily increases oxygen availability in liver lobules, resulting in hepatic hypoxia and greater O<sub>2</sub><sup>-</sup> and H<sub>2</sub>O<sub>2</sub> production. Besides, long-term alcohol consumption can up-regulate cytochrome P450 2E1 in hepatocytes and nicotinamide adenine dinucleotide phosphate oxidase in Kupffer cells, leading to alcoholic liver diseases [2].

It is believed that plants are the best sources of natural antioxidants, and these bioactive phytochemicals are valued as playing primary roles in relieving oxidation process by quenching free radicals, chelating metal ions, and scavenging oxygen in the biological systems [3], thus prevents free radical mediated oxidation of proteins, lipids and DNA which is implicated in the pathogenesis of many diseases. Flavonoids have multiple reduction capacities to facilitate the donation of electrons from hydroxyl moieties to oxidizing radical species [4]. On the other hand, the contribution of flavonoids to hepatoprotective properties has been clarified by several

\*Corresponding author. Jen-Chieh Tsai, Department of Medicinal Botanicals and Health Applications, Da-Yeh University, 168 University Rd, Dacun, Chang-Hua, Taiwan, ROC.

Tel: +86 886 4 8511888x1790

Fax: +86 886 4 8511349

E-mail: [jenchieh@mail.dyu.edu.tw](mailto:jenchieh@mail.dyu.edu.tw)

Peer review under responsibility of Hainan Medical College.

Foundation project: This work was in part supported by the Forestry Bureau of the Republic of China (Taiwan) (No. 99-06-5-02).

researches [1] that *Ajuga nipponensis* (*A. nipponensis*) is a widespread folk edible medicine in Asia for treating traumatic injury and inflammation [5,6]. Among numerous active compound contained in *A. nipponensis* and ecdysterones are unique for their great variety [7]. Although it lacks active hydroxyl groups which are critical characteristics of antioxidants such as tea polyphenols [8], the antioxidant properties of ecdysterones have been indicated by recent studies [4]. Further research indicates that ecdysteroids could also inhibit the peroxidation of rat liver microsomes induced by hydroxyl radicals, as monitored by the formation of thiobarbituric acid reactive substances, and prevent radical-induced decrease of membrane fluidity [9]. Apart from ecdysterones, flavonoids are another potential bio-constituents of *A. nipponensis* [10].

Extraction is a critical method for isolating the effective component in plants. There are several parameters concerned in the extraction of antioxidative components from plant tissues. This includes the types of solvent, the ratio of solvent-to-sample, the extraction temperature and the extraction time [11]. Antioxidants extracted by traditional techniques may undergo degradation during extraction, but the ultrasonic-assisted extraction (UAE) will enhance production yield as well as avoid thermal damages. UAE possesses greater solvent penetration into plant tissues due to the formation of cavitation bubble collapse, thus improving its solvent extraction ability [12].

Our Previous studies also show the UAE was the better way to isolate of flavonoids and ecdysteroids from *Ajuga* plants, and show the potential for hypoglycaemic health food [6]. This study was aimed to develop an optimum extraction method of *A. nipponensis* and evaluate its antioxidant and hepatoprotective effects. We used orthogonal experimental design to determine the optimal extraction parameters for ecdysterones and flavonoids of *A. nipponensis*. Several factors including extraction time, extraction temperature and the ratio of sample-to-solvent were taken into consideration. Finally, the hepatoprotective abilities of *A. nipponensis* extracts were evaluated by CCl<sub>4</sub>-induced animal models.

## 2. Materials and methods

### 2.1. Sample preparation of plant materials

*A. nipponensis* was provided by Taiwan Agricultural Research Institute, Council of Agriculture, Executive Yuan (Taiwan) with assistance from Instructor Te-Hsun Lin (Department of Medicinal Botany and Healthcare, Da-Yeh University, Taichung, Taiwan). Prior to extraction, the raw *A. nipponensis* was freeze-dried, then crushed by the homogenizer, and sieved over a 60-mesh screen.

In UAE, 3.0 g of *A. nipponensis* powders were mixed with 100 mL of 70:30 ratio of ethanol, and the mixture was sonicated in the ultrasonic extraction tank for 50 min at 50 °C with a frequency of 28 kHz. The extracts were suction filtered to discard the residue and the filtrates were then filtered through a 0.45 µm filter.

### 2.2. Orthogonal array experimental design

An orthogonal experiment [L<sub>9</sub>(3)<sup>4</sup>] test design was used to optimize the extraction conditions of *A. nipponensis* by UAE. In

the present study, extraction was accomplished with 70% ethanol according to the pre-test results. The effect of extraction time, the extraction temperature and the ratio of sample to extraction solvent on extraction yield was investigated. Each factor had three optimization levels. Levels for extraction time factor 1, 2, 3 representative of 45, 50 and 55 min, extraction temperature factor 1, 2, 3 representative of 40, 50 and 60 °C, and ratio of *A. nipponensis* to 70% ethanol factor 1, 2, 3 representative of 1:10, 1:20 and 1:30.

### 2.3. Determination of flavonoid contents

Flavonoid contents were determined using the aluminum chloride colorimetric method. The procedure was performed based on the protocol from the previous research method [13]. In brief, 0.5 mL of *A. nipponensis* extracts prepared at various concentrations (0.01, 0.05, 0.1, 0.5 and 1 mg/mL) were reacted with 1.5 mL of 95% ethanol, 0.1 mL of 10% aluminum chloride hexahydrate, 0.1 mL of 1 M potassium acetate (CH<sub>3</sub>COOK), and 2.8 mL of deionized water for 40 min at room temperature. Finally, the absorbance of the mixture was measured at 415 nm. Quercetin was used as the standard.

### 2.4. Determination of ecdysterone contents

The ecdysterone contents were analyzed with an RP-18 column by a Hitachi L-7400 HPLC system (Tokyo, Japan). The flow rate was 1.0 mL/min, and the composition ratio of the mobile phase was methanol/water at 40:60 (v/v). Ecdysterone was detected at 248 nm and 20-hydroxyecdysone was used as the standard [(0.01–0.1) mg/mL].

### 2.5. DPPH scavenging assay

The antioxidant abilities of *A. nipponensis* extracts were determined by their capacities to neutralize radicals of DPPH [di(phenyl)-(2,4,6-trinitrophenyl)iminoazanium]. Each 100 µL of 1 mM DPPH solution was placed into test tubes, and 100 µL varying *A. nipponensis* extracts concentrations in methanol each 20 µL of tested sample was dissolved in 80 µL of methanol were added. Reaction was completed at room temperature for 30 min, and the absorbance was measured at 517 nm. All experiments were performed in triplicate. Butylated hydroxyanisole (BHA) was used as a positive control. The capability to scavenge the DPPH radical was calculated by the following equation (%): = [(Ac – As)/Ac], where Ac is the absorbance of the control reaction and as is the absorbance in the present of samples or BHA. Therefore, lower values represent higher antioxidant ability of each tested sample.

### 2.6. Reducing power assay

An aliquot of sample (5 mL) at different concentrations was mixed with 2.5 mL of 0.2 M phosphate buffer and 2.5 mL of 1% potassium ferricyanide. The mixture was then reacted at 50 °C for 20 min. After cooling, 2.5 mL of 10% trichloroacetic acid was added. After that, 5 mL of the mentioned reaction liquid was mixed with 5 mL of distilled water and 1 mL of 0.1% iron (III) chloride. Finally, the absorbance was detected at 700 nm. All experiments were performed in triplicate. BHA was used as a positive control.

### 2.7. Experimental animals and hepatotoxic induction by CCL<sub>4</sub>

An acute CCL<sub>4</sub>-induced toxicity model was performed according to the previous research method [14]. Thirty ICR male mice weighting (20 ± 0.5) g were obtained from National Laboratory Animal Center, Taipei, Taiwan, and housed in a controlled environment at (23 ± 1) °C as well as a relative humidity of 40%–60% with a 12 h/12 h light/dark photoperiod. After adaption for a week, the mice were divided into four groups (*n* = 6 per group) and fed through gastrostomy tube. Normal control and hepatotoxic control groups were fed with saline (1 mL/100 g BW); positive control group was given silymarin (100 mg/kg BW); and treatment group was fed with 250 mg/kg and 500 mg/kg BW of *Ajuga nipponensis* extract respectively. Silymarin is a well-known liver protective agent [15]. On the eighth day, the normal control group was injected with soybean oil (0.3 mL/100 g BW) and the other groups were inject with 2% CCL<sub>4</sub>/soybean oil (0.3 mL/100 g BW) in the intraperitoneal area 6 h prior to blood collection. Blood was collected through tail bleeding. The activities of serum glutamic oxaloacetic transaminase (GOT) and glutamic pyruvic transaminase (GPT) were immediately determined after centrifugation of collected blood at 3 000×g for 10 min. The Liver tissues were collected from the same lobes and then fixed in a 10% buffered formaldehyde solution and embedded in paraffin. They were further cut into 2 μm sections, stained with hematoxylin and eosin, and then examined under light microscopy.

Liver tissues were collected from the same lobes and trimmed to 2 mm thickness. Then these tissues were fixed in 10% buffered formaldehyde solution and embedded in paraffin and further cut into 2 μm sections, stained with hematoxylin and eosin (H&E), and then examined under light microscopy.

### 2.8. Statistical analysis

Results were expressed as mean ± SD. Differences among the groups were subjected to a one-way ANOVA (analysis of variance) followed by Duncan's multiple range, Tukey's HSD or Fisher's tests. Statistical significance was accepted when a *P*-value was less than 0.01 or 0.05.

## 3. Results

### 3.1. Optimization of extracting flavonoids and ecdysterones from *A. nipponensis* by UAE

Optimization of the experimental condition is a critical step in an extraction process. UAE parameters were optimized to obtain the best extraction of flavonoids and ecdysterones from *A. nipponensis*. Thus, the extraction time, extraction temperature and the ratio of sample to solvent were considered as important factors during the UAE process of *A. nipponensis*. As shown in Table 1 given certain parameters, the results varied; and Table 2 showed that the parameters influence (R) on the amount of flavonoid contents in *A. nipponensis* extraction in the order of C > A > B, and parameters influence (R) on ecdysterone contents in the order of C > B > A. The ratio of sample to solvent was found to be the most important determinant of the contents

**Table 1**

Definition and level of factors in orthogonal array experimental design.

Levels	Factors		
	A	B	C
1	45	40	1:10
2	50	50	1:20
3	55	60	1:30

A: Extraction time (min); B: Extraction temperature (°C); C: Extraction proportions of sample/solvent.

of antioxidant compounds. In summary, the maximum yield of flavonoids (7.87 ± 0.10) mg/g and ecdysterones (0.73 ± 0.02) mg/g from *A. nipponensis* were obtained when the extraction time was 50 min, the extraction temperature was 60 °C, and the ratio of sample to 70% ethanol was 1: 20, as indicated in Table 3.

### 3.2. Effects of *A. nipponensis* extracts on DPPH scavenging and reducing power assays

After determining the most efficient extraction process of *A. nipponensis* extracts by UAE, we also detected the antioxidant abilities of *A. nipponensis* extracts processed by defined extraction conditions because the antioxidant bioactivities of plants might mainly contribute to their biological functions including hepatoprotective properties. The DPPH scavenging effects increased with increased *A. nipponensis* extraction concentration (Figure 1A). Results indicated that when the concentration of group 6 was 5 mg/mL, it exhibited the best antioxidant activities with a DPPH scavenging ability approximately 84.86% ± 1.76%, similar to BHA 90.08% ± 2.46%. Flavonoids have a diphenylpropane structure, in which B or A ring of the structure contains catechol, C-3 includes either hydroxy or gallic acid (galloyl group), C-2 and C-3 are linked by a double bond and the C-4 is in a ketoform. These provided flavonoid its free radical scavenging capacity [16].

As indicated in Figure 1B the higher concentration of BHA and *A. nipponensis* extracts the greater antioxidant property. Results indicated that when the concentration was 5 mg/mL, the absorbance values of BHA and *A. nipponensis* extracts reducing

**Table 2**

Results of orthogonal array design experiments by range analysis.

		A	B	C
Flavonoid	k1	5.80	6.08	5.21
	k2	6.92	6.14	6.97
	k3	6.20	6.70	6.75
	R	1.11	0.62	1.76
Ecdysterone	k1	0.62	0.57	0.53
	k2	0.61	0.59	0.66
	k3	0.61	0.67	0.65
	R	0.01	0.10	0.13


A: Extraction time (min); B: Extraction temperature (°C); C: Extraction proportions of sample/solvent. k1, k2, and k3 were average results corresponding to their respective factor group 1, 2, and 3. R was extreme difference of the maximum and minimum k1, k2, and k3 for each factor reflecting the contribution of the factor to the extraction result.

**Table 3**

Orthogonal array of the experiments on flavonoid and ecdysterone contents of *A. nipponensis* extracts by UAE.

Experiment no.	Factors			Results	
	A	B	C	Flavonoid content (mg/g)	Ecdysterone content (mg/g)
1	1	1	1	4.37 ± 0.14	0.51 ± 0.05
2	1	2	2	5.92 ± 0.18	0.65 ± 0.07
3	1	3	3	7.12 ± 0.21	0.69 ± 0.03
4	2	1	3	6.75 ± 0.11	0.61 ± 0.01
5	2	2	1	6.13 ± 0.16	0.49 ± 0.06
6	2	3	2	7.87 ± 0.10	0.73 ± 0.02
7	3	1	2	7.12 ± 0.13	0.60 ± 0.09
8	3	2	3	6.37 ± 0.11	0.64 ± 0.08
9	3	3	1	5.12 ± 0.17	0.59 ± 0.04

A: Extraction time (min); B: Extraction temperature (°C); C: Extraction proportions of sample/solvent. Each value presents mean ± SD ( $n = 3$ ).




**Figure 1.** *A. nipponensis* treatment effects on DPPH scavenging and reducing power. (A) The DPPH scavenging effects of *A. nipponensis* at different concentrations compared with BHA; (B) Reducing power effect of *A. nipponensis* at different concentrations compared with BHA. *A. nipponensis* was extracted by 70% ethanol (sample: extraction solvent (v/v) = 1: 20) at 60 °C for 50 min. BHA was used as the positive control.


power are (3.00 ± 0.09) mg/mL and (2.43 ± 0.04) mg/mL respectively.

### 3.3. Hepatoprotective effects of *A. nipponensis* extracts by UAE in CCl<sub>4</sub>-induced acute liver injury


Figures 2 and 3 represented the results of serum GPT and GOT levels in experimental animals. Consistent with previous evidences, the GPT and GOT levels increased after CCl<sub>4</sub> treatments (Figures 2 and 3). However, the data showed that in CCl<sub>4</sub>-induced model, *A. nipponensis* extracts treatments (500 mg/kg body weight, *p.o.*) significantly reduced the serum GPT and GOT levels compared to the hepatotoxic control groups ( $P < 0.01$  or 0.05), implying its ability to restore the liver injury.



**Figure 2.** GOT values of CCl<sub>4</sub>-induced mice after administration of *A. nipponensis* extracts. Each value presents mean ± SD ( $n = 6$ ).  $^*P < 0.01$ . Statically significant was calculated relative to the hepatotoxic control group.  $^{##}P < 0.01$ . Statically significant was calculated relative to the normal group.



**Figure 3.** GPT values of CCl<sub>4</sub>-induced mice after administration of *A. nipponensis* extracts. Each value presents mean ± SD ( $n = 6$ ).  $^*P < 0.05$ ,  $^{**}P < 0.01$ . Statically significant was calculated relative to the hepatotoxic control group.  $^{##}P < 0.01$ . Statically significant was calculated relative to the normal group.



**Figure 4.** The pathological assessment ( $\times 200$ ). A: Blank; B:  $\text{CCl}_4$ ; C:  $\text{CCl}_4$ +Silymarin; D:  $\text{CCl}_4$ +*A. nipponensis* (250 mg/kg); E:  $\text{CCl}_4$ +*A. nipponensis* (500 mg/kg).

### 3.4. Histological analyses

As shown in Figure 4, hepatic cells of  $\text{CCl}_4$ -induced group were observed vacuole formation, necrosis and hemolysis around the central vein when compared to the control group. This phenomenon was significantly improved by treatment with *A. nipponensis* extracts (500 mg/kg).

## 4. Discussion

UAE functions creating bubbles or cavities close to a solid surface and causes a strong impact to the solid surface from its high-speed jets of liquid, thus resulting in the collapse of cavities and the erosion of the surfaces [16,17]. Moreover UAE has been used in extraction process for the food industry and processing in several researches for essential oil extraction of sage (*Salvia officinalis*); extraction of polyphenols from orange (*Citrus sinensis* L.) peel; extraction of antioxidants from *Rosmarinus officinalis*, rutin from *Sophora japonica*, etc [18–22]. Except flavonoids concerned as vital antioxidants in our initial experiments for the determination of extraction method used for *A. nipponensis* extracts, ecdysterones are unique active compounds contained in *Ajuga* plants. Beyond traditional usages like antifungal, anthelmintic against intestinal disorders and other functions, recent works have confirmed multiple biological activities related to ecdysterones, for example, hepatic function stimulations [23]. Despite lack of hydroxyl groups, the antioxidant activity of ecdysterones has been truly

proved by some studies [9] and it might be contributed to their hepatoprotective abilities. As a result, we both took flavonoids and ecdysterones as active antioxidants for optimization of *A. nipponensis* extracts by UAE. In our previously study, the hypoglycaemic effects of UAE and SFE extracts from five common *Ajuga* species in Taiwan were investigated. The results indicated that UAE was a better extraction technique for getting higher flavonoid and ecdysterone contents than SFE. Furthermore, UAE prepared *A. nipponensis* extracts had  $\alpha$ -glucosidase inhibitory activity and glucose uptake effect *in vitro* and showed the hypoglycaemic effect *in vivo*.

In the antioxidant system, oxidation is accompanied by reduction. Antioxidant index is the measurement of reducing power. Antioxidant capacity can be measured from absorbance values, the higher the absorbance the greater the antioxidant capacity. Figure 1 showed *A. nipponensis* extracts had slightly lowered free radical scavenging effect and reducing power than synthetic BHA, *A. nipponensis* extracts being a natural product and an efficient scavenging power has a higher commercial value.

GPT and GOT, also called alanine aminotransferase and aspartate aminotransferase, are important hepatic enzymes. GPT as well as GOT is important markers of liver injury, and their elevation after  $\text{CCl}_4$ -induction has been clarified [14]. In the hepatoprotective test, pretreatment of *A. nipponensis* significantly decreased GOT and GPT levels. These results demonstrated the hepatoprotective potentials of *A. nipponensis* extracts *in vivo*. On the other hand, histological biopsies

showed the hepatic cells were protected by treatment with *A. nipponensis*. The above-mentioned findings further illustrated that *A. nipponensis* extracts prevented CCl<sub>4</sub>-induced liver injury and maintained hepatocellular membrane structural integrity.

Suitable extraction methods, extraction parameters optimization, and *A. nipponensis* antioxidant as well as hepatoprotective abilities were investigated in this study. This study determined that the UAE was the best extraction process for isolating flavonoids from *A. nipponensis*. Through the orthogonal test, the extraction time, extraction temperature and the ratio of sample to solvent factors for UAE of *A. nipponensis* were optimized. The antioxidant abilities, DPPH scavenging and reducing power abilities of the *A. nipponensis* extract under the defined extraction conditions were also clarified. Finally, this study found that *A. nipponensis* extracts treatment restored the serum GPT and GOT levels significantly and prevented the hepatocellular injury. Thus, the present study demonstrated that *A. nipponensis* extracts by UAE possess excellent antioxidant abilities which might contribute to its hepatoprotective potentials.

### Conflict of interest statement

We declare that we have no conflict of interest.

### Acknowledgment

This work was in part supported by the Forestry Bureau of the Republic of China (Taiwan) (no. 99-06-5-02).

### References

- [1] Domitrovic R, Jakovac H, Vasiljev Marchesi V, Vladimirov Knezevic S, Cvijanovic O, Tadic Z, et al. Differential hepatoprotective mechanisms of rutin and quercetin in CCl<sub>4</sub>-intoxicated BALB/cN mice. *Acta Pharmacol Sin* 2012; **33**(10): 1260-1270.
- [2] Chen KH, Li PC, Lin WH, Chien CT, Low BH. Depression by a green tea extract of alcohol-induced oxidative stress and lipogenesis in rat liver. *Biosci Biotech Biochem* 2011; **75**(9): 1668-1676.
- [3] Valko M, Leibfritz D, Moncol J, Cronin MTD, Mazur M, Telser J. Free radicals and antioxidants in normal physiological functions and human disease. *Int J Biochem Cell Biol* 2007; **39**(1): 44-84.
- [4] Cadot Y, Chevalier M, Barbeau G. Evolution of the localisation and composition of phenolics in grape skin between veraison and maturity in relation to water availability and some climatic conditions. *J Sci Food Agric* 2011; **91**(11): 1963-1976.
- [5] Israili ZH, Lyoussi B. Ethnopharmacology of the plants of genus *Ajuga*. *Pak J Pharm Sci* 2009; **22**(4): 425-462.
- [6] Hsieh CW, Cheng JY, Wang TH, Wang HJ, Ho WJ. Hypoglycaemic effects of *Ajuga* extract *in vitro* and *in vivo*. *J Funct Foods* 2014; **6**(0): 224-230.
- [7] Coll J, Tandrón YA, Zeng X. New phytoecdysteroids from cultured plants of *Ajuga nipponensis* Makino. *Steroids* 2007; **72**(3): 270-277.
- [8] Liu ZQ, Ma LP, Zhou B, Yang L, Liu ZL. Antioxidative effects of green tea polyphenols on free radical initiated and photosensitized peroxidation of human low density lipoprotein. *Chem Phys Lipids* 2000; **106**(1): 53-63.
- [9] Cai YJ, Dai JQ, Fang JG, Ma LP, Hou LF, Yang L, et al. Antioxidative and free radical scavenging effects of ecdysteroids from *Serratula strangulata*. *Can J Physiol Pharm* 2002; **80**(12): 1187-1194.
- [10] Zheng XK, Zhang L, Wang WW, Wu YY, Zhang QB, Feng WS. Anti-diabetic activity and potential mechanism of total flavonoids of *Selaginella tamariscina* (Beauv.) Spring in rats induced by high fat diet and low dose STZ. *J Ethnopharmacol* 2011; **137**(1): 662-668.
- [11] Chen S, Wu BH, Fang JB, Liu YL, Zhang HH, Fang LC, et al. Analysis of flavonoids from lotus (*Nelumbo nucifera*) leaves using high performance liquid chromatography/photodiode array detector tandem electrospray ionization mass spectrometry and an extraction method optimized by orthogonal design. *J Chromatogr A* 2012; **1227**: 145-153.
- [12] Floegel A, Kim DO, Chung SJ, Koo SI, Chun OK. Comparison of ABTS/DPPH assays to measure antioxidant capacity in popular antioxidant-rich US foods. *J Food Compos Anal* 2011; **24**(7): 1043-1048.
- [13] Lin JY, Tang CY. Determination of total phenolic and flavonoid contents in selected fruits and vegetables, as well as their stimulatory effects on mouse splenocyte proliferation. *Food Chem* 2007; **101**(1): 140-147.
- [14] Tseng SH, Chien TY, Tzeng CF, Lin YH, Wu CH, Wang CC. Prevention of hepatic oxidative injury by Xiao-Chen-Chi-Tang in mice. *J Ethnopharmacol* 2007; **111**(2): 232-239.
- [15] Comar KM, Kirby DF. Herbal remedies in gastroenterology. *J Clin Gastroenterol* 2005; **39**(6): 457-468.
- [16] Carvalho RN Jr, Moura LS, Rosa PTV, Meireles MAA. Supercritical fluid extraction from rosemary (*Rosmarinus officinalis*): kinetic data, extract's global yield, composition, and antioxidant activity. *J Supercrit Fluid* 2005; **35**(3): 197-204.
- [17] Cheung YC, Siu KC, Wu JY. Kinetic models for ultrasound-assisted extraction of water-soluble components and polysaccharides from medicinal fungi. *Food Bioprocess Technol* 2013; **6**(10): 2659-2665.
- [18] Horžić D, Jambrak A, Belščak-Cvitanović A, Komes D, Lelas V. Comparison of conventional and ultrasound assisted extraction techniques of yellow tea and bioactive composition of obtained extracts. *Food Bioprocess Technol* 2012; **5**(7): 2858-2870.
- [19] Khan MK, Abert-Vian M, Fabiano-Tixier AS, Dangles O, Chemat F. Ultrasound-assisted extraction of polyphenols (flavonone glycosides) from orange (*Citrus sinensis* L.) peel. *Food Chem* 2010; **119**(2): 851-858.
- [20] Paniwnyk L, Beaufoy E, Lorimer JP, Mason TJ. The extraction of rutin from flower buds of *Sophora japonica*. *Ultrason Sonochem* 2001; **8**(3): 299-301.
- [21] Paniwnyk L, Cai H, Albu S, Mason TJ, Cole R. The enhancement and scale up of the extraction of anti-oxidants from *Rosmarinus officinalis* using ultrasound. *Ultrason Sonochem* 2009; **16**(2): 287-292.
- [22] Sališová M, Toma Š, Mason TJ. Comparison of conventional and ultrasonically assisted extractions of pharmaceutically active compounds from *Salvia officinalis*. *Ultrason Sonochem* 1997; **4**(2): 131-134.
- [23] Lafont R, Dinan L. Practical uses for ecdysteroids in mammals including humans: and update. *J Insect Sci* 2003; **3**: 7.