

# A Geometric Integration Based on Magnus Series Expansion for Human T-Cell Lymphotropic Virus I (HTLV-I) Infection of CD4<sup>+</sup> T-Cells Model

M. Tarık Atay<sup>1</sup>, Musa Başbük<sup>2</sup> and Aytakin Eryılmaz<sup>2\*</sup>

<sup>1</sup>Department of Mechanical Engineering, Abdullah Gül University, 38080, Kayseri/Turkey

<sup>2</sup>Department of Mathematics, Nevşehir Hacı Bektaş Veli University, 50300, Nevşehir/Turkey  
Email: eryilmazaytekin@gmail.com

**Abstract** In this paper, we investigated a numeric integration based on Magnus series expansion namely Magnus Series Expansion Method (NMG) for nonlinear Human T-Cell Lymphotropic Virus I (HTLV-I) infection of CD4<sup>+</sup> T-cells model. Fourth order Magnus series expansion method (NMG4) and explicit Runge-Kutta (RK45) are used to obtain numerical solutions of HTLV-I infection of CD4<sup>+</sup> T-cells model. The results obtained by NMG4 and RK45 are compared.

**Keywords:** Non-linear Differential Equation System, Magnus Series Expansion Method, Geometric Integration, Lie Group Method.

## 1 Introduction

Dynamics of Human T-Cell Lymphotropic Virus I (HTLV-I) infection of CD4<sup>+</sup> T-cells model examined in [1-7] is,

$$\frac{dT}{dt} = \lambda - \mu_T T - k T_A T \quad (1)$$

$$\frac{dT_L}{dt} = k T_A T - (\mu_L + \alpha) T_L \quad (2)$$

$$\frac{dT_A}{dt} = \alpha T_L - (\mu_A + \rho) T_A \quad (3)$$

$$\frac{dT_M}{dt} = \rho T_A + \beta T_M (1 - T_M / T_{max}) - \mu_M T_M \quad (4)$$

with the initial conditions,

$$T(0) = P_1, \quad T_L(0) = P_2, \quad T_A(0) = P_3, \quad T_M(0) = P_4, \quad (5)$$

where  $T(t), T_L(t), T_A(t), T_M(t)$  denote the concentration of healthy CD4<sup>+</sup> T-cells at time  $t$ , the concentration of latently infected CD4<sup>+</sup> T-cells, the concentration of actively infected CD4<sup>+</sup> T-cells and the concentration of leukemic cells at time  $t$  respectively. The parameters  $\lambda, \mu_T$  and  $k$  denote the natural date rate of CD4<sup>+</sup> T-cells, the rate at which uninfected cells are contacted by actively infected cells, the rate of infection of T-cells with virus from actively infected cells respectively.  $\mu_L, \mu_A, \mu_M$  are blanket death terms for latently infected, actively infected and leukemic cells respectively.  $\alpha$  and  $\rho$  denote the rates at which latently infected and actively infected cells become actively infected and leukemic respectively. The rate  $\beta$  determines the the speed at which the saturation level for leukemia cells is reached.  $T_{max}$  is the maximal value that adult T-cell leukemia can reach [7].

The purpose of this paper is to obtain numerical solution of the system (1-4) subject to the initial conditions (5) by using a structure preserving numerical integrator based on Magnus Expansion, namely Magnus Expansion Method.

In 1954, Magnus [8] provided an exponential representation of the solution of a first order linear homogeneous differential equation for a linear operator that was named Magnus Expansion after him.

Since 1960's the Magnus expansion has been successfully applied in various areas of physics and chemistry (see [9] for a list of references). Iserles and Norsett (1997) [10,11] presented a practical recursive algorithm that generated the terms of Magnus expansion. Blanes et al. (1998) [12] considered the approximate solutions of matrix linear differential equations by matrix exponentials and the convergence issue of Magnus and Fer expansions. They obtained the upper bounds for the convergence radius in terms of the norm of the defining matrix of the system. Moan and Niesen (2008) [13] considered the question: When does the series converge? The main result they obtained, established a necessary condition for convergence.

## 2 Magnus Expansion Method

The linear differential equation on a matrix Lie-group is an equation of the form

$$Y' = A(t)Y, \quad t \geq 0, \quad Y(0) = Y_0 \in \mathcal{G}, \tag{6}$$

where  $A : \mathcal{G} \rightarrow \mathcal{R}$  is the matrix function,  $\mathcal{G}$  is the Lie group,  $\mathcal{g}$  is the Lie algebra of the corresponding Lie-group  $\mathcal{G}$ . Magnus expressed the solution of equation (6) as the exponential of a certain function [8],

$$Y(t) = e^{\Omega(t)}, \tag{7}$$

and obtained an infinite recursive series for  $\Omega$  as follows,

$$\Omega_0 \equiv 0, \tag{8}$$

$$\Omega_{n+1} = \int_0^t \text{dexp}_{\Omega_n}^{-1} A(\xi) d\xi = \sum_{k=0}^{\infty} \frac{B_k}{k!} \int_0^t \text{ad}_{\Omega_n}^k A(\xi) d\xi, \quad n = 0, 1, 3, \dots \tag{9}$$

where  $B_k$  are the Bernoulli numbers. Substituting the equation (8) into the equation (9) one can get the  $\Omega_i$  for  $i = 0, 1, 2, 3, \dots$ , respectively

$$\begin{aligned} \Omega_1 &= \int_0^t A(t_1) dt_1 \\ \Omega_2 &= \int_0^t A(t_1) dt_1 - \frac{1}{2} \int_0^t \left[ \int_0^{t_2} A(t_1) dt_1, A(t_2) \right] dt_2 + \dots \\ \Omega_3 &= \int_0^t A(t_1) dt_1 - \frac{1}{2} \int_0^t \left[ \int_0^{t_2} A(t_1) dt_1, A(t_2) \right] dt_2 \\ &\quad + \frac{1}{12} \int_0^t \left[ \int_0^{t_3} A(t_1) dt_1, \left[ \int_0^{t_3} A(t_1) dt_1, A(t_3) \right] \right] dt_3 \\ &\quad + \frac{1}{4} \int_0^t \left[ \int_0^{t_3} \left[ \int_0^{t_2} A(t_1) dt_1, A(t_2) \right] dt_2, A(t_3) \right] dt_3 + \dots \end{aligned} \tag{10}$$

The Magnus series expansion is,

$$\Omega(t) = \sum_{k=0}^{\infty} H_k(t), \tag{11}$$

where each  $H_k$  includes exactly  $k + 1$  integrals and  $k$  commutators [14].

Thus,

$$\begin{aligned} H_1 &= \int_0^t A(t_1) dt_1 \\ H_2 &= -\frac{1}{2} \int_0^t \left[ \int_0^{t_2} A(t_1) dt_1, A(t_2) \right] dt_2 \end{aligned}$$

$$\begin{aligned}
H_3 &= \frac{1}{12} \int_0^t \left[ \int_0^{t_3} A(t_1) dt_1, \left[ \int_0^{t_3} A(t_1) dt_1, A(t_3) \right] \right] dt_3 \\
&\quad + \frac{1}{4} \int_0^t \left[ \int_0^{t_3} \left[ \int_0^{t_2} A(t_1) dt_1, A(t_2) \right] dt_2, A(t_3) \right] dt_3 \\
H_4 &= -\frac{1}{24} \int_0^t \left[ \int_0^{t_4} A(t_1) dt_1, \left[ \int_0^{t_4} \left[ \int_0^{t_2} A(t_1) dt_1, A(t_2) \right] dt_2, A(t_4) \right] \right] dt_4 \\
&\quad - \frac{1}{24} \int_0^t \left[ \int_0^{t_4} \left[ \int_0^{t_2} A(t_1) dt_1, A(t_2) \right] dt_2, \left[ \int_0^{t_4} A(t_1) dt_1, A(t_4) \right] \right] dt_4 \\
&\quad - \frac{1}{24} \int_0^t \left[ \int_0^{t_4} \left[ \int_0^{t_3} A(t_1) dt_1, \left[ \int_0^{t_3} A(t_1) dt_1, A(t_3) \right] \right] dt_3, A(t_4) \right] dt_4 \\
&\quad - \frac{1}{8} \int_0^t \left[ \int_0^{t_4} \left[ \int_0^{t_3} \left[ \int_0^{t_2} A(t_1) dt_1, A(t_2) \right] dt_2, A(t_3) \right] dt_3, A(t_4) \right] dt_4.
\end{aligned}$$

By using multivariate Gaussian quadrature, Casas & Iserles [15] presented algorithms for third and fourth order Magnus Expansion Method (NMG3 and NMG4) for nonlinear Lie type differential equations as follows, Third order Magnus Expansion Method for nonlinear equations (NMG3)

$$\begin{aligned}
u_1 &= 0 \\
k_1 &= hA(0, Y_0) \\
u_2 &= \frac{1}{2}k_1 \\
k_2 &= hA\left(\frac{h}{2}, e^{u_2}Y_0\right) \\
u_3 &= \frac{1}{4}(k_1 + k_2) \\
k_3 &= hA\left(\frac{h}{2}, e^{u_3}Y_0\right) \\
u_4 &= k_2 \\
k_4 &= hA(h, e^{u_4}Y_0) \\
v_3 &= \frac{1}{6}(k_1 + 4k_3 + k_4) - \frac{1}{3}[u_3, k_3] - \frac{1}{12}[u_4, k_4] \\
Y_1(t) &= e^{v_3}Y_0.
\end{aligned} \tag{12}$$

Fourth order Magnus Expansion Method for nonlinear equations (NMG4)

$$\begin{aligned}
u_1 &= 0 \\
k_1 &= hA(t_n, Y_n) \\
Q_1 &= k_1 \\
u_2 &= \frac{1}{2}Q_1 \\
k_2 &= hA\left(t_n + \frac{h}{2}, e^{u_2}Y_0\right) \\
Q_2 &= k_2 - k_1 \\
u_3 &= \frac{1}{2}Q_1 + \frac{1}{4}Q_2 \\
k_3 &= hA\left(t_n + \frac{h}{2}, e^{u_3}Y_0\right) \\
Q_3 &= k_3 - k_2 \\
u_4 &= Q_1 + Q_2
\end{aligned}$$

$$\begin{aligned}
 k_4 &= hA(t_n + h, e^{u_4}Y_0) \\
 Q_4 &= k_4 - 2k_2 + k_1 \\
 u_5 &= \frac{1}{2}Q_1 + \frac{1}{4}Q_2 + \frac{1}{3}Q_3 - \frac{1}{24}Q_4 - \frac{1}{48}[Q_1, Q_2] \\
 k_5 &= hA(t_n + \frac{h}{2}, e^{u_5}Y_0) \\
 Q_5 &= k_5 - k_2 \\
 u_6 &= Q_1 + Q_2 + \frac{2}{3}Q_5 + \frac{1}{6}Q_4 - \frac{1}{6}[Q_1, Q_2] \\
 k_6 &= hA(t_n + h, e^{u_6}Y_0) \\
 Q_6 &= k_6 - 2k_2 + k_1 \\
 v &= Q_1 + Q_2 + \frac{2}{3}Q_5 + \frac{1}{6}Q_6 - \frac{1}{6}[Q_1, Q_2 - Q_3 + Q_5 + \frac{1}{2}Q_6] \\
 Y_{n+1} &= e^v Y_n.
 \end{aligned}
 \tag{13}$$

### 3 Numerical Results

In this section, NMG4 and the fourth order explicit Runge-ÅŠKutta (RK45) are applied to the Human T-Cell Lymphotropic Virus I (HTLV-I) infection of CD4<sup>+</sup> T-cells model by Mathematica 7. Throughout this paper we set  $\mu_T = 0.66mm^3/day$ ,  $\mu_L = 0.06day$ ,  $\mu_A = 0.05day$ ,  $\mu_M = 0.005day$ ,  $k = 0.5$ ,  $\lambda = 0.6$ ,  $\alpha = 0.004day$ ,  $\beta = 0.0003day$ ,  $\rho = 0.00004day$  and  $T_{max} = 2200mm^3$ .

$$\frac{dT}{dt} = \lambda - \mu_T T - kT_A T \tag{14}$$

$$\frac{dT_L}{dt} = kT_A T - (\mu_L + \alpha)T_L \tag{15}$$

$$\frac{dT_A}{dt} = \alpha T_L - (\mu_A + \rho)T_A \tag{16}$$

$$\frac{dT_M}{dt} = \rho T_A + \beta T_M (1 - T_M/T_{max}) - \mu_M T_M \tag{17}$$

with the initial conditions,

$$T(0) = 1000, \quad T_L(0) = 250, \quad T_A(0) = 1.5, \quad T_M(0) = 0, \tag{18}$$

By using the following transformation [16],

$$T = p, \quad T_L = q, \quad T_A = r, \quad T_M = s, \tag{19}$$

the equation system (14-17) yields the Lie-type matrix equation,

$$\begin{bmatrix} p \\ q \\ r \\ s \\ 1 \end{bmatrix}' = \begin{bmatrix} -\mu_T - kr & 0 & 0 & 0 & \lambda \\ kr & -\mu_L - \alpha & 0 & 0 & 0 \\ 0 & \alpha & -\mu_A - \rho & 0 & 0 \\ 0 & 0 & \rho & \beta(1 - \frac{s}{T_{max}}) - \mu_A & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} p \\ q \\ r \\ s \\ 1 \end{bmatrix}, \tag{20}$$

where prime denotes derivative with respect to time  $t$ .

If we call  $\begin{bmatrix} p \\ q \\ r \\ s \\ 1 \end{bmatrix}$  as  $Y$  and  $\begin{bmatrix} -\mu_T - kr & 0 & 0 & 0 & \lambda \\ kr & -\mu_L - \alpha & 0 & 0 & 0 \\ 0 & \alpha & -\mu_A - \rho & 0 & 0 \\ 0 & 0 & \rho & \beta(1 - \frac{s}{T_{max}}) - \mu_A & 0 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$  as  $A$ , then equation system (14-17) takes the form,

$$Y' = AY. \tag{21}$$

By using the algorithm (13) and conducting the iteration for the time interval (0,1000) with the step size  $\frac{1}{10}$ , the following results are obtained. Table 1-4. compare the NMG4 and RK45 solutions for the time interval (0,20) with the step size  $\frac{1}{10}$  for the equation system (14-17). The obtained results for  $T(t)$ ,  $T_L(t)$ ,  $T_A(t)$  and  $T_M(t)$  are given in Fig. 1-13.

**Table 1.** NMG4 and RK45 solutions of  $T(t)$  for the time interval (0,20) with the step size  $\frac{1}{10}$  for the equation system (14-17).

$t$	NMG4 solution	RK45 solution	Absolute Difference
0	1000	1000	0
2	7.13582	7.13582	$4.12047 \times 10^{-6}$
4	0.096573	0.096573	$6.62869 \times 10^{-6}$
6	0.0723808	0.0723808	$6.33054 \times 10^{-7}$
8	0.0610988	0.0610988	$5.42661 \times 10^{-7}$
10	0.0549001	0.0549001	$4.25295 \times 10^{-7}$
12	0.0513063	0.0513063	$3.03046 \times 10^{-7}$
14	0.049261	0.049261	$1.8866 \times 10^{-7}$
16	0.048239	0.048239	$8.85135 \times 10^{-8}$
18	0.047952	0.047952	$4.96236 \times 10^{-9}$
20	0.0482307	0.0482307	$5.93419 \times 10^{-8}$

**Table 2.** NMG4 and RK45 solutions of  $T_L(t)$  for the time interval (0,20) with the step size  $\frac{1}{10}$  for the equation system (14-17).


$t$	NMG4 solution	RK45 solution	Absolute Difference
0	250	250	0
2	796.75	796.75	$7.44233 \times 10^{-5}$
4	707.304	707.304	$6.10276 \times 10^{-5}$
6	623.37	623.37	$5.38087 \times 10^{-5}$
8	549.53	549.53	$4.73637 \times 10^{-5}$
10	484.566	484.566	$4.16416 \times 10^{-5}$
12	427.411	427.411	$3.6581 \times 10^{-5}$
14	377.125	377.125	$3.21183 \times 10^{-5}$
16	332.882	332.882	$2.81897 \times 10^{-5}$
18	293.955	293.955	$2.47478 \times 10^{-5}$
20	259.704	259.703	$2.17017 \times 10^{-5}$

**Table 3.** NMG4 and RK45 solutions of  $T_A(t)$  for the time interval (0,20) with the step size  $\frac{1}{10}$  for the equation system (14-17).


$t$	NMG4 solution	RK45 solution	Absolute Difference
0	1.5	1.5	
2	6.39744	6.39744	$1.79348 \times 10^{-7}$
4	11.5167	11.5167	$6.37668 \times 10^{-7}$
6	15.4728	15.4728	$1.00891 \times 10^{-7}$
8	18.453	18.453	$1.29307 \times 10^{-6}$
10	20.6224	20.6224	$1.50456 \times 10^{-6}$
12	22.1215	22.1215	$1.65579 \times 10^{-6}$
14	23.0698	23.0698	$1.75723 \times 10^{-6}$
16	23.5689	23.5689	$1.81803 \times 10^{-6}$
18	23.7046	23.7046	$1.84408 \times 10^{-6}$
20	23.5496	23.5496	$1.84767 \times 10^{-6}$

**Table 4.** NMG4 and RK45 solutions of  $T_M(t)$  for the time interval (0,20) with the step size  $\frac{1}{10}$  for the equation system (14-17).


$t$	NMG4 solution	RK45 solution	Absolute Difference
0	0	0	0
2	0.000290763	0.000290763	$5.77703 \times 10^{-11}$
4	0.00100986	0.00100986	$2.48410 \times 10^{-11}$
6	0.00208226	0.00208226	$4.16242 \times 10^{-11}$
8	0.00341953	0.00341953	$1.33443 \times 10^{-10}$
10	0.00494826	0.00494826	$2.44045 \times 10^{-10}$
12	0.00660783	0.00660783	$3.67955 \times 10^{-10}$
14	0.00834854	0.00834854	$5.00714 \times 10^{-10}$
16	0.01013	0.01013	$6.38569 \times 10^{-10}$
18	0.0119194	0.0119194	$7.79346 \times 10^{-10}$
20	0.0136909	0.0136909	$9.17819 \times 10^{-10}$




**Figure 1.** NMG4 solutions of  $T(t)$  for the time interval (0,20) with the step size  $\frac{1}{10}$  for the equation system (14-17)




**Figure 2.** NMG4 solutions of  $T_L(t)$  for the time interval (0,20) with the step size  $\frac{1}{10}$  for the equation system (14-17)




**Figure 3.** NMG4 solutions of  $T_A(t)$  for the time interval (0,20) with the step size  $\frac{1}{10}$  for the equation system (14-17)




**Figure 4.** NMG4 solutions of  $T_M(t)$  for the time interval (0,20) with the step size  $\frac{1}{10}$  for the equation system (14-17)




**Figure 5.** NMG4 solutions of  $T(t)$  for the time interval  $(0,100)$  with the step size  $\frac{1}{10}$  for the equation system (14-17) **Figure 6.** NMG4 solutions of  $T_L(t)$  for the time interval  $(0,100)$  with the step size  $\frac{1}{10}$  for the equation system (14-17)




**Figure 7.** NMG4 solutions of  $T_A(t)$  for the time interval  $(0,100)$  with the step size  $\frac{1}{10}$  for the equation system (14-17) **Figure 8.** NMG4 solutions of  $T_M(t)$  for the time interval  $(0,100)$  with the step size  $\frac{1}{10}$  for the equation system (14-17)



**Figure 9.** NMG4 solutions of  $T_A(t)$  for the time interval  $(0,1000)$  with the step size  $\frac{1}{10}$  for the equation system (14-17) **Figure 10.** NMG4 solutions of  $T_M(t)$  for the time interval  $(0,1000)$  with the step size  $\frac{1}{10}$  for the equation system (14-17)



**Figure 11.** NMG4 solutions of  $T(t)$  for the time interval  $(0,1000)$  with the step size  $\frac{1}{10}$  for the equation system (14-17) **Figure 12.** NMG4 solutions of  $T_L(t)$  for the time interval  $(0,1000)$  with the step size  $\frac{1}{10}$  for the equation system (14-17)



**Figure 13.** NMG4 solutions of  $T(t), T_L(t), T_A(t)$  and  $T_M(t)$  for the time interval  $(0,1000)$  with the step size  $\frac{1}{10}$  for the equation system (14-17)

### 4 Conclusion

In this study, a reliable, efficient and structure preserving [17] numerical algorithm based on the Magnus series expansion is applied to solve the nonlinear differential equation system (14-17) which occurs in Human T-Cell Lymphotropic Virus I (HTLV-I) infection of  $CD4^+$  T-cells model. The obtained results by NMG4 and the fourth order explicit Runge-Kutta (RK45) are compared in Table 1-4. The obtained results are shown in Fig. 1-13. The numerical solutions of the NMG4 are in excellent agreement with respect to the RK45 solutions. As a result Magnus Expansion Method is an efficient and accurate tool for Human T-Cell Lymphotropic Virus I (HTLV-I) infection of  $CD4^+$  T-cells model.



## References

1. Gómez-Acevedo, H., Li, M.Y., *Backward bifurcation in a model for HTLV-I infection of CD4<sup>+</sup> T cells*, Bulletin of Mathematical Biology, Volume 67: 101-114, (2005).
2. Song, X., Li, Y., *Global stability and periodic solution of a model for HTLV-I infection and ATL progression*, Applied Mathematics and Computation, Volume 180: 401-410, (2006).
3. Stilianakis, N., Seydel, J., *Modeling the T-cell Dynamics and Pathogenesis of HTLV-I Infection*, Bulletin of Mathematical Biology, Volume 61: 935-947, (1999).
4. Eshima, N., Tabata, M., Okada, T., Karukaya, S., *Population dynamics of HTLV-I infection: a discrete-time mathematical epidemic model approach*, Mathematical Medicine and Biology 20(1):29-45, (2003).
5. Seydel, J., Stilianakis, N., *HTLV-I Dynamics: A Mathematical Model*, Sexually Transmitted Diseases 27(10): 652-653, (2000).
6. Wang, L., Li, M.Y. and Kirschner, D., *Mathematical analysis of the global dynamics of a model for HTLV-I infection and ATL progression*, Mathematical Biosciences 179: 207-217, (2002).
7. Gokdogan, A. and Merdan, M., *A multistage homotopy perturbation method for solving Human T-Cell Lymphotropic Virus I (HTLV-I) infection of T-cells model*, Middle-East Journal of Scientific Research, 9(4): 503-509, (2011).
8. Magnus, W., *On the exponential solution of differential equations for a linear operator*, Comm. Pure and Appl. Math. 7: 639-673, (1954).
9. Blanes S. et al., *The Magnus expansion and some of its applications*, Physics Reports 470: 151-238, (2009).
10. Iserles, A. and Norsett, S.P., *On the solution of linear differential equations in Lie groups*, Technical Report 1997/NA3, Department of Applied Mathematics and Theoretical Physics, University of Cambridge, England, (1997).
11. Iserles, A. and Norsett, S.P., *Linear ODEs in Lie groups*, DAMTP tech. report 1997/NA9, University of Cambridge, (1997).
12. Blanes, S. et al., *Magnus and Fer expansions for matrix differential equations: the convergence problem*, J. Phys. A 31: 259-268, (1998).
13. Moan, P.C. and Niesen, J., *Convergence of the Magnus Series*, Foundations of Computational Mathematics 8(3): 291-301, (2008).
14. Iserles, A. et al., *Lie-group Methods*, Acta Numerica 9: 215-365, (2000).
15. Casas, F. and Iserles, A., *Explicit Magnus expansions for nonlinear equations*, Journal of Physics A: Mathematical and General 39: 5445-5461, (2000).
16. Blanes, S. et al., *Time-averaging and exponential integrators for non-homogeneous linear IVPs and BVPs*, Applied Numerical Mathematics 62: 875-894, (2014).
17. Castellano, A. et al., *Geometric numerical integrators based on the Magnus expansion in bifurcation problems for non-linear elastic solids*, Frattura ed Integrità Strutturale 29: 128-138, (2014).