

ECONOMIC ANALYSIS OF POST HARVEST LOSSES IN ONION IN JODHPUR DISTRICT OF RAJASTHAN

SWATI SHARMA

Assistant Professor, ASPEE Agribusiness Management Institute, Navsari Agricultural University, Navsari, Gujarat, India

ABSTRACT

A study was undertaken to examine the nature and extent of post-harvest losses in onion supply chain in the Jodhpur district which is major onion district of Rajasthan. A total sample size of 75 onion growers, 20 wholesalers and 25 retailers was taken from Jodhpur district. Maximum aggregate post-harvest losses (24.47 kg/q) have been found at producer level due to faulty storage, lack of adequate transportation, drying, improper handling of the produce at the time of marketing, rotted bulbs, doubles, bolters, poor packing facilities, injury at the time of harvesting and de-topping. Total losses in the supply chain were estimated to be 30.87 kg/q (79.27 %) losses were observed at farm level and rest were contributed at wholesale and retail level. The farm level post harvest losses excluding the losses at farm level storage for Jodhpur district was estimated to be 119013.58q for the year 2009-10

KEYWORDS: Post-Harvest Losses, Onion, Supply Chain, Farm Level

INTRODUCTION

An onion, today being compared with diamonds indicates its value for a normal household budget. Global review states that China is the first in area and production of onion while India occupies second position in the production and exports to Dubai, Kuwait, Saudi Arabia, Middle East, Malaysia, Singapore, Bangladesh, Sri Lanka etc. Onion is an important commercial vegetable crop. About 82.02 million tonnes onion is produced in the world from 8217 thousand hectares of area. India is one of the major onions producing country with a production of 14.84 million tonnes from an area of 1.01 million hectares.

Onion is one of the most important commercial vegetable crops grown in Rajasthan. It occupies about 25 -30% area of the total vegetable crops in the state. It is predominantly a rabi season crop but in kharif season it accounts for about 10 -15% of the total production. Rajasthan has a comparative advantage in onion production. In the total area and production in the country, Rajasthan stands 7th position in area and production and productivity in India and contributes about 57.46('000 ha) in area and 704.96 (in '000 MT) in production (NHB, 2013-14).

In India post harvest losses has been accounted as one of the major problem in most of the vegetables including onion. Verma and Singh (2004) reported overall losses in vegetables up to 25 per cent of total production. Severe loses occur because of poor transportation facilities, lack of know-how, poor management and improper market facilities or due to careless handling of the produce by farmers, market intermediaries and consumers (Gauraha and Thakur, 2008; Singh et al., 2008). The study by Karim and Wee (1996) had revealed that well managed post-harvest activities for vegetables led to higher yields and profits to producers. It is therefore, important that the post-harvest practices be given as much attention as production practices.

Therefore, a study on post-harvest losses of onion was undertaken. The study aimed at assessing the extent of losses, which in turn will facilitate development of proper measures to reduce post-harvest losses at farm and trade level.

MATERIAL AND METHODS

The study was carried out in the Jodhpur district of Rajasthan. The methodology for collection of primary data involved structured interview schedule using personal interview method. A structured schedule was prepared for collection of data from 750nion farmers from district for the fulfillment of objectives.

Multistage stage sampling was adopted: At first stage, only highest onion producing 3 tehsils were selected in district. At second stage 3-4 villages were randomly selected for the purpose of primary data collection in district. At third stage the list of the onion growers along with their operational holdings in each of the randomly selected village was prepared with the help of villagers. From this prepared list of onion growers, 7-8 onion growers were randomly selected from each village for the present study. A total sample of seventy five onion growers from ten villages was drawn from district. Also a sample of 20 wholesalers and, 25 retailers dealing in onion were selected randomly for obtaining the information pertaining to the postharvest losses. Data obtained from the survey was analyzed through tabular analysis including appropriate statistical tools.

RESULTS AND DISCUSSIONS

The post-harvest losses were estimated at producer level to trader level. Yet the losses at producer level have been estimated at different stages like; harvesting, grading and packing, handling and transportation and marketing; whereas the losses at trader level have been estimated at loading-unloading, transportation, grading and selling stages. The findings of whole post – harvest losses of onion were analyzed at farm level first and then it was worked out on per hectare basis and finally it was estimated on per quintal of output produce and the findings are depicted in the Tables 1 to 5.

Analysis of Post Harvest Losses in Jodhpur District

The post-harvest losses have been assessed at different stages of supply chain of onion from produce to consumer *viz.*, at the farm level, during storage, wholesale marketing level and retailing level.

Total onion bulbs produce by all the selected onion growers were 12736.43q from the 43.61 ha area. Total marketable yield was recorded 11598.70q and unmarketable bulbs at field level was recorded 1137.73q at the time of harvesting due to various losses at field levels like doubles, bolters, rotted bulbs, drying, bulbs injuries, de-topping, packing, transportations, marketing etc. Out of the total marketable bulb yield (11598.70) 557.80 q of onion bulbs kept by the sample onion growers for own used and for onion seed production programme in the next crop season at own farms and remaining 11040.90q bulbs was available for marketing of onion (Table 1). Results further showed that out of total available marketable produce (11040.90q), 3386.24q was sold and 7654.66q was stored by 30.67 per cent and 69.33 per cent of the onion growers, respectively. It was also evident from the results that among the total farmers involved for sold of onion after harvesting of crop, 43.48 per cent sold their produce immediately in the market within seven days and 56.52 per cent sold their produce in the market within one month period. It could be inferred from the Table 1 that out of the total selected farmers in the sample size group's farm, 69.33 per cent farmers were stored onion (7654.66q) at farm level in Jodhpur. The analysis of stored onion further revealed that 15.54 per cent (1036.38q) post harvest losses was occurred in the storage during six months storage period (June-November).

121

Total Onion Producti on by Selected Farmers (Q)	Losses In Total Productio n at Farm Level (Q)	Total Market able Bulbs (Q)	Onion Kept for Own Used (Q)	Total Marketed Surplus (Q)	Quantity of Produce Sold Within One Month (Q)	Share of Farmer's Sold Produce with in One Month Period (%)	Total Quantity Stored for Storage (Q)	Share of Farmer's Stored Produce in Onion Storage (%)
12736.43	1137.73	11598.7 0	557.80	11040.90	3386.24	30.67	7654.66	69.33

 Table 1: Overall Average Quantity of Onion Bulbs Produce, Marketable Quantity, Marketed

 Surplus and Stored Quantity of Onion in the Jodhpur District

*Total number of respondents= 75 farmers, 20 wholesalers, 25 retailers

Post Harvest Losses at Farm Level

The post harvest losses in onion at the field level in Jodhpur district were estimated at 8.93 kg/q of produce (Table 2). The resultant loss at farm level were due to injury at the time of harvesting, de-topping, doubles, bolters, rotted bulbs, drying ,under sized unmarketable bulbs, faulty storage and transportation and improper handling of the produce at the time of marketing. Among these, loss to faulty storage was the highest (15.54 kg/q) followed by improper transportation, which resulted in a loss of 1.71 kg/q. in the produce The loss due to faulty storage appeared to be rather high because most of the respondents stored the produce by adopting traditional method of storage. The drying loss was 1.45 kg/q. The loss of output due to faulty de-topping in onion resulted in a loss of 0.59 kg/q because of improper cutting of the top. The losses due to injuries at the time of harvest in onion resulted in a loss of 0.80 kg/q. Improper packaging and rough handling of the produce during marketing resulted in post harvest losses and these losses were estimated to be respectively 0.90 kg/q and 1.33 kg/q for onion (Table 2).

Post Harvest Losses at Wholesaler and Retailer Level

The total post harvest losses at wholesale level were estimated at 3.22 kg/q of produce. The storage losses in onion at the wholesale level were 1.15 kg/q, and loss in transit have been worked out to be 1.97 kg/q. Transportation loss in onion crop was higher because of the use of unsuitable transport means, negligent driving and rough roads. The post harvest loss at the retail level was 3.09 kg/q for onion. The transit and storage loss was 1.18 kg/q in the marketed produce. The loss due to spoilage and multiple handling of produce during retailing was 0.81 kg/q. The post harvest loss at the retailer level due to bad weather and foreign matter content was 1.10 kg/q (Table 2).

S. No	Different Stages	Loss (Kg/Q)	Per Cent Loss	
Ia	Farm	Level Losses I	Due to	
1.	Harvesting injuries	0.80	2.59	
2.	De-topping	0.59	1.91	
3.	Drying	1.45	5.93	
4.	Doubles and bolters and rotted	1.05	4.31	
5.	Rotted and undersized bulbs	1.10	3.56	
6.	Packing	0.90	2.92	

Table 2: Post Harvest Losses in Onion at Different Stages in the Jodhpur District

7.	Transportation	1.71	5.54		
8.	Marketing	1.33	4.31		
	Total Losses at farm Level	8.93	28.93		
Ib	Losses during Storage	15.54	50.34		
	Overall total losses at farm level	24.47	79.27		
II	Wholesaler Level	Losses Due to			
1.	Storage	1.15	3.73		
2.	Transit	1.97	6.38		
	Total loss at wholesale level	3.22	10.43		
III	Retailer Level Losses Due to				
1.	Transit and storage	1.18	3.82		
2.	Bad weather and foreign matter content	1.10	3.56		
3.	spoilage and multiple handling loss	0.81	2.62		
	Total Loss at Retailer Level	3.09	10.01		
	Total Logg	20.97	100.00		

*Total number of respondents= 75 farmers, 20 wholesalers, 25 retailers

Total Post Harvest Loss

The post harvest loss occurring at field and market was added up to 30.87 kg/q. Maximum post harvest loss was observed at the farm level (24.47 kg/q) accounting for 79.27% of the total post harvest loss (Table 2). This loss was observed due to the tender texture and high moisture content of onion led to deterioration of quality in onion and in turn the quantity loss occurred at different post harvest stages like drying, storage, packing and transportations at field level. Further 3.22 kg/q of the output losses were observed at the wholesale level, accounting for 10.43%. The loss at retail level was to the tune of 3.09 kg/q (10.01%). Result of the study further revealed that post harvest loss at the wholesale level was observed relatively more as compared to that at the retailer level during marketing of onion in Jodhpur market (2011).

Table 3: Average per Hectare Post Harvest Loss at Farm Level in the Jodhpur District

Produce Quantity of Onion Bulbs by the Total Sample Farmers (Q)	Total Average Area Under Onion Cultivation with the Total Sample Farmers (Ha)	Average Per Ha Onion Yield Produce by the Each Sample Farmers (Q/Ha)	Post Harvest Losses at Farm Level (Kg/Q)	Per Ha Post Harvest Losses (Q/Ha)
12736.43	43.61	292.05	8.93	26.08

*Total number of respondents= 75 farmers, 20 wholesalers, 25 retailers

The per hectare post harvest loss at farm level was estimated to be 26.08q (Table 3). The average yield for the sample farmers was 292.05 /ha for onion. This means that farmers in the process of post harvest operations lost about 8.93 per cent of onion output produce by the farmers. The post harvest losses in Jodhpur district in onion crop would be much higher (15.33%), if the post harvest at the market level were also added to the above values.

Produce Quantity of Onion by the Total Sample Farmers (Q)	Total Number of Sample Farmers in Each District	Average Per Farm Onion Bulb Yield Production by the Each Sample Farmer (Q/Farm)	Post Harvest Losses at farm Level (Kg/Q)	Per Farm Post Harvest Losses (Q/Farm)
12736.43	75	169.82	8.93	15.16

 Table 4: Average per farm Post Harvest Loss in the Jodhpur District

*Total number of respondents= 75 farmers, 20 wholesalers, 25 retailers

The average per farm onion output was 169.82q. The per farm post harvest loss was estimated to be 15.16q in onion (Table 4).

Total Onion Cultivated Area in Each District (Ha)	Total Onion Production in Each District (Q)	Average Onion Productivity in Each District (Q/Ha)	Post Harvest Losses at Farm Level in District (Kg/Q)	Per Hectare Post Harvest Losses in Each District (Q /Ha)	Quantity of Spoilage Produce Due to Post Harvest Losses (Q)
15701	1332340	84.86	8.93	7.58	119013.58

Table 5: Post Harvest Loss at Farm Level (Per Hectare) in the Jodhpur District

*Total number of respondents= 75 farmers, 20 wholesalers, 25 retailers

The post harvest loss at farm level (7.58q/ha) for the Jodhpur district works out to be 119013.58q during 2009-10 (Table 5).

CONCLUSIONS

The study has estimated post-harvest losses in onion in Jodhpur district of Rajasthan. At producer level, the postharvest losses have been found maximum (24.47 kg/q). The total post-harvest losses in onion at wholesale level were found to be 3.22 kg/q and at retailer level it was 3.09 kg/q. And overall loss was reported as 30.87 kg/q. A large amount of losses (15.54 kg/q) also takes place during storage at farm. Across different stages, the losses have been found maximum at the grower level in onion. The spoilage/loss of onion at the grower level results from lack of his knowledge about proper post-harvest management loss at farm level were due to injury at the time of harvesting, de-topping, doubles, bolters, rotted bulbs, drying, under sized unmarketable bulbs, faulty storage and transportation and improper handling of the produce at the time of marketing contributes more to the problem. This results from farmer's lack of knowledge about post harvest management. Therefore, there is an urgent need of training the vegetable growers on scientific post-harvest techniques, if the vegetable production is to be sustained on a profitable basis in the region. Appropriate farm level storage also needs to be given due attention for reducing post harvest losses.

REFERENCES

- 1. Gajanana, T.M., Murthy, D. Sreenivasa, Sudha, M. and Dakshinamoorty, V. (2006) Marketing and estimation of post-harvest losses of tomato crop in Karnataka, Indian Journal of Agricultural Marketing, 20 (1): 1-7.
- 2. Gauraha, A.K. and Thakur, B.S. (2008) Comparative economic analysis of post-harvest losses in vegetables and foodgrains crops in Chhattisgarh, Indian Journal of Agricultural Economics, 63(3): 376.

- Hazarika, C. (2006) Post-harvest loss and food security A study on fruits and vegetables in Assam, Indian Journal of Agricultural Economics, 61(3): 418-419.
- Hazarika, C. (2008) Extent of post-harvest losses of ginger in Assam A micro level analysis, Indian Journal of Agricultural Economics, 63(3): 370-371.
- 5. Karim, A. and Wee, M.M.B. (1996) Reducing post harvest losses in vegetables. In: Proceedings of Workshop on Vegetable Crops Agribusiness, held at BARC, Farm Gate, Dhaka. 2-4 May.
- 6. Kumar, D. Kishor, Basavaraja, H. and Mahajanshetti, S.B. (2006) An economic analysis of post-harvest losses in vegetables in Karnataka, Indian Journal of Agricultural Economics, 61(1): 134-146.
- 7. National Horticulture Board database- 2013-14
- 8. Singh, R.B., Kushwaha, R.K. and Verma, Sunil Kumar (2008) An economic appraisal of post-harvest losses in vegetable in Uttar Pradesh, Indian Journal of Agricultural Economics, 63(3): 378.
- 9. Verma, Ajay and Singh, K.P. (2004) An economic analysis of post-harvest losses in fresh vegetables, Indian Journal of Agricultural Marketing, 18(1): 136-139.