Journal of Coastal Life Medicine

journal homepage: www.jclmm.com

Original article https://doi.org/10.12980/jclm.5.2017J7-48

©2017 by the Journal of Coastal Life Medicine. All rights reserved.

Unusual presence of *Coryphaena hippurus* Linnaeus, 1758 (Perciformes: Coryphaenidae) under an offshore oil platform in Southern Brazil

Mauro Sinopoli¹, Pietro Battaglia², João Pedro Barreiros^{3,4*}

¹Institute for Environmental Protection and Research (ISPRA), S.T.S of Palermo, Via Salvatore Puglisi 9, Palermo 90143, Italia

²Institute for Environmental Protection and Research (ISPRA), Laboratory of Milazzo, Via dei Mille 46, Milazzo (Me) 98057, Italia

³CE3C/ABG – Centre for Ecology, Evolution and Environmental Changes/Azorean Biodiversity Group, Angra do Heroísmo 9700-042, Portugal

⁴Faculty of Agrarian and Environmental Sciences, University of the Azores, Angra do Heroísmo 9700-042, Portugal

ARTICLE INFO

Article history: Received 10 Mar 2017 Received in revised form 13 May 2017 Accepted 30 May 2017 Available online 14 Jun 2017

Keywords: Food habits Fish aggregating device Pelagic trophic chains Artificial reef

ABSTRACT

Objective: To study the food habits of a sample of the dolphinfish (*Coryphaena hippurus*) around the Petrobrás PXIV oil platform located off Southern Brazil.

Methods: We examined the stomach contents of 28 adult dolphinfish caught around this platform between 2000 and 2001.

Results: Results highlighted a diet consisting mainly of pelagic fish, although cephalopods and crustaceans are also present. A large amount of clupeids found in the stomachs could be due to baits used by fishermen that do fish around the platform.

Conclusions: In this research we discuss the possibility of this occurrence being related to trophic opportunistic behavior. This could be a reason why adult dolphinfish are attracted by this platform since the whole structure does act as a fish aggregating device.

1. Introduction

Offshore oil platforms are known as effective fish aggregating devices. Much like artificial reefs, platforms constitute an artificial substrate which may be quite relevant in pelagic environments. These are rapidly colonized by a number of encrusting organisms and act as direct/indirect food sources for a vast number of fish and other taxa. They also play an important role as sheltering places and spots for reproduction, especially for demersal fish that show parental care, such as adhesive egg clusters. Offshore platforms also act as visually attractive sites for pelagic fish species[1,2].

After a few field trips to this particular platform (Petrobrás,

Tel: +351-965691616

E-mail: joao.ps.barreiros@uac.pt

Foundation Project: Supported by Universidade do Vale do Itajaí's Center for Marine Sciences, Fernando L. Diehl. Petrobrás – Petróleos do Brasil, S.A. (PB/2000/01/10).

The journal implements double-blind peer review practiced by specially invited international editorial board members.

PXIV), we noticed that it attracted a variety of species, including economically valuable ones such as the dolphinfish [*Coryphaena hippurus* (*C. hippurus*)] and many scombrids especially the skipjack tuna *Katsuwonus pelamis* (Linnaeus, 1758). Dolphinfish are in fact widely known to form schools under floating objects^[3]. Most dolphinfish found were adults, while juveniles are more often reported to aggregate in similar circumstances^[4]. The specific purpose of this study is to understand whether the unusual high presence of adult dolphinfish rather than juvenile associated with the platform can be due or linked to trophic reasons. For this reason a study of stomach contents was carried out. This study also has the general objective to collect information on the diet of this species in the SW Atlantic from where no previous data are known.

2. Materials and methods

Overall 28 dolphinfish were collected between 2000 and 2001

^{*}Corresponding author: João Pedro Barreiros, Faculty of Agrarian and Environmental Sciences, University of the Azores, Angra do Heroísmo 9700-042, Portugal.

around Petrobrás oil platform PXIV located ca. 100 nautical miles off the northern coast of Santa Catarina State (26°46'2.2" S; 46°47'2.15" W) over a depth of 200 m. Specimens were collected by hook and line using artificial lures and natural baits (squid, tuna and beef).

After collection, all fish were sexed, weighed to the nearest g, measured to the nearest cm (total length, standard length) and their stomachs were removed for weighing (nearest g) and preserved in a formaldehyde solution (10%). In the laboratory, stomachs were opened by longitudinal dissection for better removal of contents.

The preys were dried with blotting paper, weighed and counted. Taxa identification was carried out to the lowest possible taxonomic level. The dietary importance of each food item was assessed by calculating the following indexesas used in[5]:

Percentage frequency of occurrence where,

% F = (number of stomachs containing prey item i/total number of non-empty stomachs \times 100)

Percent abundance where,

% N = (number of individuals of prey item i/total number of all prey items \times 100)

Percent weight where,

% W = (weight of prey item i/total weight of all prey items \times 100)

Index of relative importance (IRI) = (% N + % W) × % F expressed as % IRI = (IRI/ Σ IRI) × 100

In order to compare the diet of males and females, the % F index was then calculated after grouping prey as follows: C – cephalopods; Cr – crustaceans; F – fish and E – empty stomach.

3. Results

Twenty-eight specimens of *C. hippurus* (13 males and 15 females) ranging between 104 and 161 cm total length were collected and studied.

Only 7 of the 28 stomachs analyzed were empty. Dolphinfish associated with platforms showed a marked preference for fish that make upto more than 90% in terms of % IRI in the stomachs. Among fish prey, coastal pelagic clupeids were the main prey item with 39.3% of IRI, while oceanic pelagic fish belonging to Exocoetidae, demersal fish (*i.e.* Monacanthidae) and reef-associated fish (*i.e.* Carangidae) each contributed with less than 5% to *C. hippurus* diet in terms of % IRI (Table 1). Pelagic cephalopods (Loliginidae) (% IRI = 7.1) and amphipod crustaceans (% IRI = 0.1) were also present (Table 1).

Male and female C. hippurus feed on comparable proportions

of fish, cephalopods, crustaceans, while a greater proportion of empty stomachs was found in males than in females (Figure 1).



Figure 1. Frequency of occurrence calculated separately for male (13 stomachs) and female (15 stomachs).

F: Fish; C: Cephalopods; Cr: Crustaceans; E: Empty stomachs. Sample of *C. hippurus* was from platform Petrobrás PXIV, South Brazil.

Table 1

Food items found in the stomachs of adult *C. hippurus* from the offshore oil platform Petrobrás (PXIV, S Brazil) and dietary indexes calculated for each prey.

Group	ITEM	Species	% N	% F	% W	% IRI
Fish	Belonidae	Tylosurus acus	4.3	9.5	14.3	4.0
	Carangidae	unidentified	3.8	9.5	2.6	1.4
	Clupeidae	unidentified	46.5	23.8	26.0	39.3
	Exocoetidae	unidentified	3.2	9.5	8.6	2.6
	Monacanthidae	Aluterus monoceros	0.5	4.8	34.3	3.8
	Unidentified fish		27.0	47.6	11.4	41.7
Molluscs	Loliginidae	unidentified	13.5	19.0	2.8	7.1
Crustaceans	Amphipoda	unidentified	1.1	4.8	0.1	0.1

N: Abundance percentage; W: Weight percentage; F: Frequency of occurrence; IRI: Index of relative abundance percentage.

4. Discussion

This study focuses on the diet of adult dolphinfish around an oil platform in the Western Atlantic Ocean. Although the sample size is low, the information collected adds important data about a poorly-known life stage of this species. Indeed, most existing studies are based on youngof-the-year dolphinfish[6], as the adults are erratic making it difficult to track their movements which are still poorly understood. Our results do show that adult dolphinfish are mainly piscivorous predators and the main prey observed in this study are comparable with food items identified in sampled C. hippurus from the Eastern Pacific[7], the Western Pacific from Colombia to Panama, the Caribbean^[3] and in the Western Mediterranean[4,8]. These studies also reported that Exocoetidae were the main fish predated by dolphinfish. In our sample, although the preys mentioned above are certainly present, the dominant prey item detected was mainly clupeids. Presence of other food items such as mollusks, crustaceans and even algae led us to confirm an opportunistic feeding behaviour also referred by other authors[4,6,7].

Juvenile dolphinfish are known to use floating devices

or objects as refuge sites, nursery grounds and recruitment areas[9] and this may well be a direct cause for the aggregation of *C. hippurus* in our study area (Petrobrás oil platform PXIV). The condition in which adults represent the most abundant portion of the dolphinfish population associated with a platform is an unusual situation that could be due to the period of study. Indeed, juveniles (from 20 to 60 cm total length) are very abundant under floating structures in their early months and this study was carried out far from the period in which juveniles are associated with platforms.

Petrobrás Platform PXIV as well as other similar structures act as attractive devices towards a guild of pelagic predators ranging from large Scombridae (*e.g. Katsuwonus pelamis* and *Thunnus albacares* (Bonnaterre, 1788) to sharks and cetaceans. All this concentrated biomass is known by fishermen who often make impressive catches under these favorable circumstances^[9,10]. The massive presence of fishing vessels that use live bait, mainly small Clupeids, justifies the high number of these fishes in the stomachs. This could be a reason for adult dolphinfish to be attracted by this platform. The hypothetical role of the fishermen's activity in attracting dolphinfish near to the platforms is also strengthened by the fact that clupeids are not species normally attracted to these structures^[10].

This study gives a contribution to our knowledge of this species' feeding ecology, confirming that adult *C. hippurus* have similar feeding habits to those from other areas in terms of typical prey. Data from the stomach contents give an important indication regarding the considerable abundance of adults of this species, rather than juveniles, near the platform. Anyway, the assumption that it is the bait used by fishermen that attracts dolphinfish needs more data to be verified.

Conflict of interest statement

We declare that we have no conflict of interest.

Acknowledgments

The Authors wish to thank Maurício Hostim-Silva for his co-coordination of Project PXIV and Fellippe Daros for his tremendous effort in collections of specimens. Thanks also goes to all of the vast staff that worked in this project and to the former Director of Universidade do Vale do Itajaí's Center for Marine Sciences, Fernando L. Diehl. Petrobrás – Petróleos do Brasil, S.A. for funding (PB/2000/01/10) and the personnel aboard PXIV platform for their constant help and enthusiasm during our many journeys and stays with them. Finally, the authors are grateful to Anthony Green, who kindly improved the syntax of the manuscript.

References

- Claisse JT, Pondella DJ, Love M, Bull AS. Biological productivity of fish associated with offshore oil and gas structures on the Pacific OCS. Vantuna Research Group, Occidental College, Los Angeles, California. BOEM Cooperative Agreement No. M12AC00003. 2014. [Online] Available from: https://www.boem. gov/2014-030/ [Accessed on 9th June, 2017]
- [2] Cervigón F, Cipriani R, Fisher W, Garibaldi L, Hendrickx M, Lemus AJ, et al. [FAO species identification cards for fishing purposes. A field guide to commercial marine and brackish water species on the northern coast of South America]. Commission of European Communities and NORAD. Roma: FAO; 1992. Spanish.
- [3] Oxenford HA, Hunte W. Feeding habits of the dolphinfish (Coryphaena hippurus) in the Eastern Caribbean. Sci Mar 1999;
 63(3-4): 303-15.
- [4] Castriota L, Pipitone C, Campagnuolo S, Romanelli M, Potoschi A, Andaloro F. Diet of *Coryphaena hippurus* (Coryphaenidae) associated with FADs in the Ionian and Southern Tyrrhenian Seas. *Cybium* 2007; **31**(4): 435-41.
- [5] Galván-Magaña F, Olson RJ. Food habits and consumption rates of dolphinfish (*Coryphaena hippurus*) in the Eastern Pacific Ocean. Ocean Fishery Bull 2002; 100(2): 279-98.
- [6] Froese R, Pauly D. FishBase. Coryphaena hippurus Linnaeus, 1758. Common dolphinfish. [Online] Available from: http://www. fishbase.org/summary/Coryphaena-hippurus.html [Accessed on 9th June, 2017]
- [7] Varela JL, Lucas-Pilozo CR, Gonzalez-Duarte MM. Diet of common dolphinfish (*Coryphaena hippurus*) in the Pacific coast of Ecuador. J Mar Biol Assoc UK 2016; 97(1): 207-13.
- [8] Riera F, Grau A, Grau AM, Pastor E, Quetglas A, Pou S. Ichthyofauna associated with drifting floating objects in the Balearic Islands (western Mediterranean). *Sci Mar* 1999; 63(3-4): 229-35.
- [9] Hostim-Silva M, Fontes J, Afonso P, Serpa N, Sazima C, Barreiros JP, et al. [Oil platforms Deep sea fish meet]. *Ciência Hoje* 2002; 31(183): 20-6. Portuguese.
- [10] Ajemiana MJ, Wetz JJ, Shipley-Lozano B, Stunz GW. Rapid assessment of fish communities on submerged oil and gas platform reefs using remotely operated vehicles. *Fish Res* 2015; 167: 143-55.