

A Bed for Burn Wound Care: A Stainless Steel Grating-Top Table as a Moveable Sterilized Burn Wound Cleansing and Operating Table

Kriangsak Sirirak, M.D.

Department of Surgery, Prapokklao hospital, Muang, Chantaburi 20000, Thailand.

ABSTRACT

A bed for burn wound care: a stainless steel grating-top table is used as a moveable sterilized table for burn wound cleansing or an operating table. Its top can be sterilized by autoclaving and after sterilization; burned patients can lie on this sterilized area for burn wound cleansing or surgical manipulation. The grating-top table can be moved room to room like a trolley by the wheel and its brake at each leg. In Prapokklao hospital, the grating-top table has been used for burn wound cleansing (using sterile solutions: warm saline, sterile water, and chlorhexidine scrub), and as an operating table when there is an indication for surgical manipulation. This study covered a total of 88 inpatients (during October 2009 - August 2012), and there were 213 times of wound cleansing (bathing and dressing) and 141 times of surgical manipulation (excision debridement and skin coverage operations). Follow with standard burn wound care principles: the grating-top table has been shown not only to be easy to use and comfortable for patients, but also results in less infection rate with good healing resulting in 86 success healing cases (97.73%) and 2 death cases (2.27%). The bed provides very cheap cost in burn wound care with appropriate outcomes.

Keywords: Burn wound care, burn wound care device, burn wound cleansing, and burn wound dressing

Siriraj Med J 2014;66:97-102

E-journal: <http://www.sirirajmedj.com>

INTRODUCTION

Burn wound care has undergone a change in the past few decades; resulting in increased incidence of patient survival.¹⁻⁴ Greater understanding of burn injury introduces better aspects of burn care that have substantial advantages in a hospital burn clinic. The parti-

cular aspects in the hospital burn clinic cover: to improve acute care (such as resuscitation),⁵ early burn wound excision, early evolution of effective skin bank, infection control, and alternative wound-closure materials and strategies.³ Strict infection control practices⁶⁻⁸ (isolation, use gloves and gown during patient contact) are necessary to prevent incidence of infection. Early excision of burn tissue is needed mainly to control wound-related infection in burns.^{9,10} Prapokklao hospital, a local public hospital in eastern Thailand, has no burn unit, so burn patients with stable overall conditions are

Correspondence to: Kriangsak Sirirak
E-mail: kriangsak.sirirak@gmail.com
Received 1 February 2013
Revised 20 September 2013
Accepted 24 September 2013

admitted in regular surgical wards and burn patients with unstable overall conditions are admitted in a critical care unit. There are few types of equipment for severe burn care, so many types of equipment have been adapted for serving this problem. For burn wound care, a sterilized moveable stainless steel-grating table has been adapted for burn wound cleansing, dressing and as a sterilized operating table since 1990.

MATERIALS AND METHODS

Burn wound care makes a significant role in improvement of total burn care, in a hospital burn clinic. In Prapokklao hospital, there were all kinds of burn patients the same as hospitals with burn units have. Before burn wounds were cleansed and dressed on a bed where it was extremely difficult to control burn wound contamination and infection. Since 1990, in a ward, an author has used a 60 by 70 centimeters stainless steel grating-top with sink below [Fig 1] for cleansing the burn wound on extremities, but this method was limited for severe burns or burn wounds at trunk. The author has consequently made a larger table [Fig 2] for taking care of these burn patients, of all sizes and in all locations. The new grating-top table size is 70 by 200 centimeters with an upside down pyramid stainless sink below for waste solution drainage and wheel with lock at all four legs.




Fig 1. Small sterilized grating top table.

The author designed the stainless grating-top into three separated gratings, sized 66 by 69 centimeters. The autoclaved grating-top was changed for each manipulation time. A burn patient was gently placed on this sterilized grating table and started cleansing with warm saline and chlorhexidine scrub, beginning from their surrounding burned area. Using a large amount of warm saline a burn wound was washed until clean, shown by clear waste solution, and then necrotic tissue and debris was removed. If there was an indication for surgical manipulation; it could immediately be performed on the table. At the end of the manipulation, each patient was dressed in nanocrystalline silver dressing,^{12,13} which was changed twice a week. Patients were allowed to shower at the ward when the patient could tolerate and was able to shower by themselves. Cleaning of the stainless grating-top table and sink used detergents and antiseptic solutions after manipulation was finished. The stainless steel medical instruments and grating top were sterilized by autoclaving technique. Culture specimens from equipment were regularly obtained as usual according to standard infection control protocol. Retrospective open trial study was conducted in surgical wards at Prapokklao hospital and was approved by Prapokklao's Institutional Ethics Committee (No.18/December 24, 2012). During October, 2009 - August, 2012, there were 91 hospitalized



Fig 2. Burn Wound Care Device: A Movable sterilized stainless steel grating top table.

burn patients in Prapokklao hospital. The rule in this department was to include burn patients who needed admission for definite wound care by dressing and/or surgical manipulation, in an easily cleaned infection control room, or operating room. All burn patients were treated with these techniques, on the stainless grating-top table (Fig 3, Fig 4, and Fig 5). The wound inspection and clinical monitoring were regularly done following the standard guideline (CDC criteria) for wound infection with swab culture, tissue culture and hemoculture.

RESULTS

There were 88 burn patients who met the inclusion criteria (Fig 6), 61 males and 27 females, with average age 28.30612 ± 22.79721 years (8 months-86 years), and most of them (27 patients) were young with ages below 10 years of age, average %TBSA burn 26.04545 ± 19.3936 (10-90%) varied in all degrees of burn depth. Dressing meant as scrubbing with chlorhexidine solution, swabbing, low-pressure irrigation, cleansing with sterile solutions (normal saline and sterile water), and followed by nanocrystalline silver dressing. Surgical manipulations included debridement (surgical blade, free hand dermatome, and hydrosurgery device), negative pressure dressing, amputation, and skin coverage with flap and/or skin graft. Of the 88 patients, there were 354 times of manipulations, wound cleansing with dressing was 213 times and surgical manipulation was 141 times. There were 10 samples (9 patients, 10.23%) from 96 samples who demonstrated positive colony growth of organisms: Methicilin Resistant *Staphylococcus Aureus* (MRSA) (4 samples), Methicilin-Susceptible *Staphylococcus Aureus* (MSSA) (one sample), *Pseudomonas* (3 samples), *Acinetobacter* (3 samples), *Klebsiella* (2 samples), *Streptococcus* (one sample), and *Enterococcus* (one sample). One hemoculture was positive for MSSA. There was one severe burn wound sepsis, but few burn

Abbreviation

%TBSA= Total Body Surface Area



Fig 3. Demonstration of burn wound dressing.




Fig 4. Demonstration of excisional debridement on the table.




Fig 5. Demonstration of excisional debridement of burn wound.

wound infections occurred and they responded well to systemic antibiotic treatment. There were two deaths (2.7%); the first an 8-month-old child with 25% TBSA with third degree burn which involved the scalp who died from accidental massive hemorrhage by arterial rupture,




Fig 6. Age distribution of burn patients.

Age in years = a

and the second a 17-year-old man with 70% TBSA concomitant with inhalation injury who died from severe sepsis within a few days after admission. There were three cases who needed to be referred to the Burn Unit at Siriraj Hospital due to critical medical conditions with critical sepsis and after their general condition was stabilized. There was a 3-year-old boy with 45% TBSA that included scalp, chest wall, both upper extremities, and lower abdomen (third degree burn at the scalp and both upper extremities). Another was a 68-year-old man with 80% TBSA. The third was a 37-year-old man with 90% TBSA. There was a total recovery of a severely burned patient, who was a 13-year-old boy with 90% TBSA with toxic exfoliative necrosis. The average length of hospital stay was 26.66 ± 29.84 days, and the maximum with 60% second to third degree acid burns was 227 days (Fig 6).

DISCUSSION

Burn wound care is one of the factors that improve mortality and outcome of patients with severe burn injury.^{3,14} Wound cleansing is a vital role in wound care to control burn wound infection and promotion of the healing process. Standard techniques used in burn wound cleansing include high-pressure irrigation, swabbing, and low-pressure irrigation, showering, bathing and washing the affected area under running liquid or immersion in a whirlpool bath. These techniques need a specialized team and equipment for serious burn patients that some hospitals do not have. To improve the quality of burn wound care, a sterile area, strict infection control, and new techniques in wound management help to improve outcomes. To create a condition like that in Prapokklao hospital, the author has adapted a sterilized stainless steel-




Fig 7. Frequency of length of hospital stay in all intervals of admission.

grating table for burn wound cleansing (Fig 3) and surgical manipulation (Fig 4, Fig 5) in a sterile area in a clean room or operating room.³ All methods of burned wound cleansing were done on the table except immersion and the main advantage of the table was that it could be used as a sterilized operating table for surgical manipulations. The author made an early burn wound excision, within a few days post burn, after cleansing to improve control of wound-related infection. Besides strict infection control practice (isolation, use of gloves and gown during patient contact), additive manipulation of burn wound(s) in a clean area took place and effective cleansing of the contaminated area on the sterilized table was used for reducing infection. Cleansing in this study used warm saline and chlorhexidine scrub, but it caused more pain for some patients, especially in this study, the young or unstable mood patients, so it may need anesthesia and/or mild sedation. There were no other pain control interventions. The duration of the procedure varied considerably from patient to patient, but this would not cause harm by using warm sterile saline solution keeping body regions warm by sterilized clothes to prevent heat loss. There was no limitation in using the table despite septicemia and/or unstable hemodynamic conditions. This moveable table was moved from room to room in Prapokklao hospital including the intensive care unit, operating theater, and wards. These manipulations were done several times a day, changing the sterilized stainless steel-grating top case by case, and cleaning its sink and legs like an operating table. During this study, there were dramatic reductions of burn wound related infections in Prapokklao hospital (one case with sepsis who survived) and acceptable mortality rate 15 (2 deaths), 1-hemorrhage shock, and 1-pulmonary complication with sepsis (2.27%, acute phase outcome). The study had to exclude 3 serious cases because we referred these cases to the Burn Unit in Siriraj Hospital. The reason was because of their needs for critical medical advanced burn care. All of them were sent back to Prapokklao hospital after their condi-

tions were stable. This bed costs about \$ 700 and each autoclaving about \$ 15 in Prapokklao hospital.

CONCLUSION

A bed for burn wound care which is a moveable sterilized stainless steel grating-top table is an extremely useful and inexpensive device that can improve burn wound care and promote wound healing process with most cleansing techniques and surgical manipulations. The bed can be widely used for basic to severe burn patients, and the main advantages of the table are that it is like a sterilized bed for all purposes in burn wound care with very cheap device. The bed can help in a large part of burn wound care, but the best systemic outcomes in burn care still need specialist burn center facilities.

ACKNOWLEDGMENTS

The author wishes to thank: Clinical Professor Apirag Chuangsuwanich for his advice, intensive care teams of Pediatric and Surgical Divisions in Prapokklao hospital, and all surgical wards teams, Siriraj Burn Unit, Mrs. Pongsri Supanpayup, Miss. Chatrudee Rasboriharn, Miss. Jintawee Buambut, and Plastic Surgery Residents Year 4 from Siriraj Hospital for their helps in good patient cares.

REFERENCES

1. Herndon DN, Spies M. Modern burn care. *Semin Pediatr Surg.* 2001 Feb(1);10:28-31.
2. Atiyeh BS, Gunn SW, Hayek SN. State of the art in burn treatment. *World J Surg.* 2005 Feb;29(2):131-148.
3. Church D, Elsayed S, Reid O, Winston B, Lindsay R. Burn Wound Infections. *Clin Microbiol Rev.* 2006 Apr;19(2):403-434.
4. Light TD, Latenser BA, Kealey GP, Wibbenmeyer LA, Rosenthal GE, Sarrazin MV. The Effect of Burn Center and Burn Center Volume on Mortality of Burned Adult-an Analysis of Data in the National Burn Respository. *J Burn Care Res.* 2009 Sep-Oct;30(5):776-782.

5. Latenser BA. Critical care of the burn patient. *Crit Care Med*. 2009 Oct;37(10):2819-26.
6. Weber J, McManus A. Infection control in burn patients. *Burns*. 2004 Dec;30(8):A16-24.
7. Church D, Elsayed S, Reid O, Winston B, Lindsay R. Burn Wound Infection. *Clin Microbiol Rev*. 2006 Apr;19(2):403-34.
8. Coban YK. Infection control in severely burned patients. *World J Crit Care Med*. 2012 Aug 4;1(4):94-101.
9. Pruitt BA, Goodwin CW, Mason D Jr. Epidemiological, dermographic, and outcome characteristics of burn injury. In: Herndon DN, ed. *Total burn care* Saunders, London England; 2002.p. 16-30.
10. Barrett JP and Herndon DN. Modulation in inflammatory and catabolic response in severe burned children by early burn wound excision in the first 24 hours. *Arch Surg*. 2003 Feb;138(2):127-132.
11. Atiyeh BS, Gunn SW, Hayek SN. State of Art in burn treatment. *World J. Surg*. 2005 Feb;29:131-148.
12. Chuntrasakul C, Muangman P, Benjathanung R, Suvanchote S, Boonpamee S, Jantarapakdee S, Kittidacha S. Clinical Experience of Acticoat™ Treatment in Extensive Burn Wounds. *Siriraj Med J*. 2007 Feb;59(2):47-51.
13. Khundkar R, Malic C, Burg T. Use of Acticoat dressing in Burns: what is the evidence? *Burns*. 2010 Sep; 36(6):751-8.
14. Lloyd JR, Hight DW. Early laminar excision: improved of control burn wound sepsis by partial dermatome debridement. *J Pediatr Surg*. 1978 Dec;13(6D):698-706.
15. Brusselsaers N, Hoste EA, Monstrey S, Colpaert KE, De Welle JJ, Vandewoude KH, Blot SI. Outcome and change over time in survival following severe burn from 1985 to 2004. *Intensive care Med*. 2005 Dec;31(12):1648-53.