


## IF: 1.634

## Asian Pacific Journal of Tropical Medicine





doi: 10.4103/1995-7645.243109

©2018 by the Asian Pacific Journal of Tropical Medicine. All rights reserved.

## Viral metagenomics analysis of poultry faeces in live poultry market, Haikou, China

Fei-fei Yin<sup>1,2,3,#</sup>, You Zhang<sup>1,2,3,#</sup>, Jasper Fuk-Woo Chan<sup>2,4,5,#</sup>, Xiu-ji Cui<sup>1</sup>, Jing-long Chen<sup>1</sup>, Yi-ji Li<sup>1</sup>, Lang-yu Rao<sup>1</sup>, Li-hua Li<sup>1</sup>, Shan-shan Wang<sup>1,2,3</sup>, Yue Wu<sup>1,2,3</sup>, Huan-huan Zhou<sup>1,2,3</sup>, Wen-qi Wang<sup>1,2,3</sup>, Kwok-Yung Yuen<sup>2,4,5</sup>, Gang Lu<sup>1,2,3,⊠</sup>

**ABSTRACT** Objective: To conduct in-depth study of the distribution and diversity of viruses in poultry is of great importance in monitoring the emergence of interspecies transmission of novel viruses that may cause epidemics with public health significance. Poultry is an economically important source of meat, eggs and feathers which plays an important role as natural reservoirs of many pathogenic viruses. Compared with wild animals, poultry have more frequent interactions and therefore opportunities to transmit their viruses to human. Methods: To study the viromes of different types of poultry in Hainan, China, we used metagenomics for deep viral nucleic acid sequencing of the faecal samples collected from chickens, ducks and pigeons from a live poultry market in Haikou. Result: The poultry viromes were identified by sequence similarity comparisons of viral reads (BLASTxE score, <5) against viral reference database. A total of 15 309 viral reads were obtained, approximately 13 063, 1 370 and 876 viral reads were generated from the chicken, duck, pigeon faecal samples, respectively. The majority of the sequences were homologous to the animal virus of Adenoviridae, Herpeaviridae, Picobirnaviridae, Reoviridae, Retroviridae, Circoviridae, Paramyxoviridae, Astroviridae, Caliciviridae, Coronaviridae, Picornaviridae, and Orthomyxoviridae. The VP4 and VP7 segments of a pigeon rotavirus, similar to fox rotavirus in group A, were sequenced and phylogenetically analyzed. The near full genome of a pigeon circovirus was also analyzed. Conclusion: The major types of poultry in a Haikou harbor many different families of viruses in their feces which may have the potential for interspecies transmissions. Further studies should be conducted to identify the most prevalent and important viruses among a larger number of poultry in Haikou and other areas in Hainan.

Keywords: Viral metagenomics; Poultry; Rotavirus; Circovirus

Article history:

Received 21 September 2018 Re Accepted 26 September 2018 Av

Received in revised form 23 September 2018 Available online 15 October 2018

E-mail: luganghn@aliyun.com Tel: +86-8980-66893745 This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-Share Alike 3.0 License, which allows others to remix, tweak and buid upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

©2018 Asian Pacific Journal of Tropical Medicine Produced by Wolters Kluwer- Medknow

How to cite this article: Yin FF, Zhang Y, Chan JF, Cui XJ, Chen JL, Li YJ, et al. Viral metagenomics analysis of poultry faeces in live poultry market, Haikou, China. Asian Pac J Trop Med 2018; 11(10 suppl):41.

<sup>&</sup>lt;sup>1</sup>Department of Pathogen Biology, Hainan Medical University, Haikou, Hainan, 571101, China

<sup>&</sup>lt;sup>2</sup>Hainan Medical University—The University of Hong Kong Joint Laboratory of Tropical Infectious Diseases, Hainan Medical University, Haikou, Hainan, 571101, China

<sup>&</sup>lt;sup>3</sup>Key Laboratory of Translation Medicine Tropical Diseases, Hainan Medical University, Haikou, Hainan, 571101, China

<sup>&</sup>lt;sup>4</sup>State Key Laboratory of Emerging Infectious Diseases, Li Ka Shing Faculty of Medicine, The University of Hong Kong, Pokfulam, Hong Kong

<sup>&</sup>lt;sup>5</sup>Department of Microbiology, Li Ka Shing Faculty of Medicine, The University of Hong Kong, Pokfulam, Hong Kong Special Administrative Region

<sup>#</sup>These authors contributed equally as co-first authors.

First author: Fei-fei Yin, Department of Pathogen Biology, Hainan Medical University, Haikou, Hainan, 571101, China.

Corresponding author: Gang Lu, Xueyuan Road 3#, Hainan Medical University, Haikou, Hainan, 571101, China.