



Evaluation of microbial contamination of ready-to-eat foods (pizza, frankfurters, sausages) in the city of Ilam

Akbar Eslami¹, Zeinab Gholami², Shokofeh Nargesi³, Bahareh Rostami³, Moayed Avazpour^{4,5*}

¹Associate Professor of Environmental Health Engineering, Environmental and Occupational Hazards Control Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran

²MSc of Environmental Health Engineering, Student Research Committee, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

³BSc in Environmental Health, Ilam University of Medical Sciences, Ilam, Iran

⁴PhD Student Research Office of Environmental Health, Department of Environmental Health Engineering, School of Public Health, Shahid Beheshti University of Medical Sciences, Tehran, Iran

⁵Instructor of Environmental Health, Department of Environmental Health Engineering, Ilam University of Medical Science, Ilam, Iran

Abstract

Background: Today in the world, disease resulting from food is considered one of the most important problems in public health. This study aimed to determine the bacterial contamination of ready-to-eat foods, i.e. fast food, in Ilam city.

Methods: In this cross-sectional, analytical study, 270 samples of ready-to-eat food, including pizza, frankfurters, and sausages, were randomly collected and tested for contamination with *Staphylococcus aureus*, *Escherichia coli*, *Shigella sonnei*, *Salmonella arizonae*, and *Enterococcus faecalis*. After examination, the collected data was analyzed using SPSS 20 software and logistic regression.

Results: From a total of 270 samples of ready-to-eat food, 27.77% was contaminated with *E. coli*, 21.48% with *S. aureus*, 13.33% with *S. sonnei*, 14.44% with *S. arizonae*, and 5.9% with *E. faecalis*. The results showed higher rates of *E. coli* and *S. aureus* contamination in pizza, frankfurters, and sausages. Also, a higher percentage of frankfurters were contaminated with microbial species than pizza or sausages. There were significant differences in microbial contamination rates ($P < 0.05$) among the three groups of food. In addition, factors such as indicators (health, sanitation, and lack of hygiene), age, gender, and education level of the operating staff had no effect on the results.

Conclusion: Based on the results, it can be concluded that bacterial contamination of ready-to-eat foods is significantly high in the city of Ilam; therefore, it is suggested that the examination of food in various stages of production and distribution can help reduce bacterial contamination, and training for the operators of shopping centers' ready-to-eat food shops and controlling pathogens are essential.

Keywords: Microbial contamination, Food, Pizza, Frankfurters, Sausages

Citation: Eslami A, Gholami Z, Nargesi S, Rostami B, Avazpour M. Evaluation of microbial contamination of ready-to-eat foods (pizza, frankfurters, sausages) in the city of Ilam. Environmental Health Engineering and Management Journal 2017; 4(2): 117–122. doi: 10.15171/EHEM.2017.16.

Article History:

Received: 29 November 2016

Accepted: 25 February 2017

ePublished: 14 March 2017

*Correspondence to:

Moayed Avazpour

Email: m_f_1859@yahoo.com

Introduction

Foodborne diseases are a major problem affecting people in poor societies (1). Microbial food contamination affects the life of people in developing countries and has a high mortality rate (2). A large variety of microorganisms or their toxins with different mechanisms are involved in causing foodborne illness (3,4). It is estimated that 30% of people in industrialized countries suffer from diseases caused by food at least once a year (5). National health agencies have reported that the average prevalence of foodborne diseases in the EU and third world countries is 3.38 and 8.915 cases per 100 000 people respectively (6-8). The annual cost of foodborne

diseases, including direct medical costs and reduced efficiency, is \$5-6 billion (9). Most cases of disease are caused by bacteria, viruses, or parasites (10). The main foodborne pathogens that can contaminate food are *Salmonella* spp., *S. aureus*, *Clostridium botulinum*, *Bacillus* spp., *Acinetobacter* spp., *Escherichia coli*, *Pseudomonas* spp., and the hepatitis A virus. Intestinal parasites include *Giardia lamblia*, *Entamoeba histolytica*, *Ascaris lumbricoides*, and *Hookworm* (11,12). From 1990 to date, several major groups of foodborne bacteria (*E. coli*, *S. aureus*, *S. arizonae*, and *E. faecalis*), have been considered in a lot of research and have attracted the attention of the food industry (13). The incidence of food-based diseases



in developed countries has had a growing trend. In developing countries such as Iran, although there are no statistics on the incidence of infections and food poisoning, infection and food poisoning occurs far more often than in developed countries because of poor conditions of production, storage, and distribution and due to the low level of health education (14). In a microbiological study, Christison et al examined 4 food distribution centers in South Africa for contamination and reported that 16% and 4% were contaminated with *Salmonella* spp. and *Listeria*, respectively (15). In a 2006 study of bacterial contamination in foods conducted in centers affiliated with Baqiyatallah University, contamination rates of *E. coli* and *Staph* were reported as 12.5% and 13.8%, respectively (16). In a 2000 study of contamination in vegetables, dairy products, and ice cream conducted in Kashan, *E. coli* infection was reported in 15% of samples (17). In a survey conducted on the bacterial contamination of raw milk in Shahre Kord, 70% of samples were contaminated with *E. coli* and 80% with coliform (18). Generally, various studies have been conducted in the field of bacterial contamination of food. In most of these studies, a specific type of bacteria in a food group was studied; therefore, a lot of information on a range of food is not available. The results of studies conducted in Iran show that the quality of ready-to-eat foods is not desirable, and the microbial contamination of these foods is higher than the standard (19,20). A lack of attention to health issues at all stages of pizza and sandwich production decreases the quality of food products. Therefore, to protect the health of consumers, this study aimed to determine the microbial quality of pizza, frankfurters, and sausages in the distribution centers of these products in the city of Ilam.

Materials and Methods

Field study

In this cross-sectional study, a wide range of bacteria (*S. aureus*, *S. sonnei*, *S. arizonae*, *E. coli* and *E. faecalis*) was evaluated in three groups of foods, pizza (90 samples), frankfurters (90 samples), and sausages (90 samples), in the city of Ilam. Different diagnostic tests were done on each sample. A total of 270 food samples were randomly collected by health experts from fast food within 3 months during 2015 in the city of Ilam. Under sterile conditions, the samples were sent to the microbiology laboratory of the Environmental Health Department where they were refrigerated. Experiments were performed less than 24 hours later. The experiments, designed to determine contamination in food products and identify and confirm the presence of bacteria, were performed according to the Iranian national standard 2461-1 method (21).

Bacterial identification

To identify and enumerate *S. aureus*, the surface culture method for *Staphylococcus* in a specific medium of Baird Parker agar was used. Black colonies that were colorless and encircled with a clear halo were counted using the

coagulase test (22). To identify *E. coli*, the Iranian national standard 2946 method was used. In this technique, after preparation, the sample was cultured in brilliant green broth and peptone water. After 24 hours of incubation, a few drops of Kovacs indole reagent were added to the tube with a peptone water medium. If a red circle formed on the top of the tube, the sample was considered positive for contamination with *E. coli*. Each diagnosis was confirmed by IMVIC tests (23). To detect *Salmonella*, after enrichment in non-selective and selective media such as lactose broth and Tetrathionate broth, the samples were cultured linearly in a selective solid medium of Brilliant Green agar (24). Then, the suspect colonies were transferred to a Triple Sugar Iron agar (TSI) differential medium and investigated for the presence or absence of *Salmonella* (24). To detect and isolate *Shigella*, after enrichment, the samples in the enrichment medium were used for Xylose lysine deoxycholate agar (XLD) differential culture media (2627) (25). To isolate the bacteria *Enterococcus* the selective medium of Kanamycin aesculin azide (KAA) Agar (Merck, Germany) was used. In this medium kanamycin was the selective agent. *Enterococcus* was determined using esculin hydrolyze, the fermentation of glucose and hydrolysis of esculin, which, in the presence of ferric citrate creates a brown color. Then, from the colonies grown, a few colonies were selected and cultured again on the same specific medium (26). To evaluate the contamination of the three studied groups of food, other factors such as indicators (health, sanitation, lack of hygiene), age, gender, and education level of operators in food supply centers were evaluated using a questionnaire.

Statistical analyses

The results of tests on various food groups were analyzed using statistical software SPSS 20 and regression test, and a level of 5% ($P < 0.05$) was considered significant. To comply with ethical issues in research, the names of the food supply centers sampled in this study have been withheld.

Results

In this study, contamination with *E. coli* and *S. aureus* was greater in the three studied food groups (pizza, frankfurters, sausages) than other microbial species. From the 270 food samples tested, 75 samples (27.77%), were contaminated with *E. coli*, 58 samples (21.48%) with *S. aureus*, 39 samples (14.44%) with *S. arizonae*, 36 samples (13.33%) with *S. sonnei*, and 16 samples (5.9%) with *E. faecalis*. From total samples of pizza, 12.22% were infected with *S. aureus*, 5.55% to *S. sonnei* 8.88% to *S. arizonae* 23.33% to *E. coli* and 1.11% to *E. faecalis* (Table1).

The number and percentage of frankfurters contaminated with the microbial species are shown in Table 2.

From all frankfurter samples, 27.77% were contaminated with *S. aureus*, 17.77% with *S. sonnei*, 24.44% with *S. arizonae*, 31.11% with *E. coli*, and 8.88% with *E. faecalis*. The number and percentage of sausages contaminated

Table 1. The frequency of bacteria contamination in pizza

Bacteria	Positive cases		Negative cases	
	Number	Percent	Number	Percent
<i>S. aureus</i>	11	12.22	79	87.78
<i>S. sonnei</i>	5	5.55	85	94.45
<i>S. arizonae</i>	8	8.88	82	91.12
<i>E. coli</i>	21	23.33	69	76.67
<i>E. faecalis</i>	1	1.11	89	98.89

Table 2. The frequency of bacteria contamination in frankfurter

Bacteria	Positive cases		Negative cases	
	Number	Percent	Number	Percent
<i>S. aureus</i>	25	27.77	65	72.23
<i>S. sonnei</i>	16	17.77	74	82.23
<i>S. arizonae</i>	22	24.44	68	75.56
<i>E. coli</i>	28	31.11	62	68.89
<i>E. faecalis</i>	8	8.88	82	91.12

Table 3. The frequency of bacteria contamination in sausages

Bacteria	Positive cases		Negative cases	
	Number	Percent	Number	Percent
<i>S. aureus</i>	22	24.44	68	75.56
<i>S. sonnei</i>	15	16.66	75	83.34
<i>S. arizonae</i>	19	21.11	71	78.89
<i>E. coli</i>	26	28.88	64	71.12
<i>E. faecalis</i>	7	7.77	83	92.23

with the microbial species are shown in Table 3. From the total samples of sausage, 24.44% was contaminated with *S. aureus*, 16.66% to *S. sonnei*, 21.11% to *S. arizonae*, 28.88% to *E. coli* and 7.77% to *E. faecalis*. The results showed that the percentage of contamination with microbial species was higher in frankfurters than in pizza or sausages. The high contamination of frankfurters (greater than pizza or sausages) is reasonable. In relation to the contamination in the three groups of food with *S. sonnei*, and *S. arizonae*, there is a significant difference; however, the *E. coli* and *E. faecalis* contamination rates were not significant ($P > 0.05$). The chance of becoming contaminated with *S. aureus*, *S. sonnei*, *S. arizonae*, *E. coli*, and *E. faecalis* was higher for frankfurters and sausages than for pizza (Table 4).

Also, the chances of contamination in three types of ready-for-consumption food were adjusted for the impact of factors such as location, age, gender, and education level, and it was found that these factors did not influence the outcome (Table 5).

Discussion

The food industry and available technologies and the preparation, distribution, and sale of food (including raw materials or ready-to-eat foods) in health centers are both important issues and require special attention (18). Because of people's interest in using ready-to-eat foods

Table 4. Univariate logistic regression analysis of Fast food contamination by a variety of bacteria in llam

Bacteria	Fast food	Univariable		
		P	Odds ratio	95% CI for OR
<i>S. aureus</i>	Pizza	0.033	-	-
	Frankfurter	0.011	2.762	1.264–6.024
	Sausages	0.037	2.325	1.051–5.128
<i>S. sonnei</i>	Pizza	0.041	-	-
	Frankfurter	0.015	3.676	1.283–10.526
	Sausages	0.023	3.40	1.179–9.803
<i>S. arizonae</i>	Pizza	0.023	-	-
	Frankfurter	0.007	3.311	1.388–7.936
	Sausages	0.025	2.739	1.200–6.666
<i>E. coli</i>	Pizza	0.487	-	-
	Frankfurter	0.272	1.483	0.765–2.873
	Sausages	0.397	1.333	0.684–2.604
<i>E. faecalis</i>	Pizza	0.128	-	-
	Frankfurter	0.044	8.695	1.062–71.428
	Sausages	0.062	7.518	0.904–62.5

Table 5. Multivariable logistic regression analysis of Fast food contamination by a variety of bacteria in llam

Bacteria	Fast food	Multivariable		
		P	Odds ratio	95% CI for OR
<i>S. aureus</i>	Pizza	0.029	-	-
	Frankfurter	0.009	2.873	1.295–6.369
	Sausages	0.034	2.398	0.186–1.069
<i>S. sonnei</i>	Pizza	0.037	-	-
	Frankfurter	0.014	3.787	1.310–10.989
	Sausages	0.022	3.496	1.200–10.204
<i>S. arizonae</i>	Pizza	0.021	-	-
	Frankfurter	0.007	3.378	1.406–8.130
	Sausages	0.024	2.785	1.141–6.802
<i>E. coli</i>	Pizza	0.464	-	-
	Frankfurter	0.226	1.526	0.769–3.030
	Sausages	0.381	1.362	0.682–2.659
<i>E. faecalis</i>	Pizza	0.114	-	-
	Frankfurter	0.038	9.433	1.127–76.923
	Sausages	0.056	8.064	0.948–66.666

and fast food such as pizza and meat products (sausages, frankfurters) to resolve a lack of protein and since such meat products are cheaper than pure meat, their use in different countries has increased several times in the past decade (14). Due to the importance of contamination of ready-to-eat foods and fast food, pathogenic microbes, and their role in causing human infection, this study examined 270 samples from 3 groups of food (pizza, frankfurters, and sausages). The results showed that among the three groups of foods tested, frankfurters and pizza, respectively, have the highest and lowest bacterial contamination. Contamination was greater with *E. coli* and *S. aureus* in the three food studied groups than with the other microbial species. Also, the percentages of contamination with *E. faecalis* in pizza, frankfurters, and sausages are low. It is noteworthy that among the three

groups of foods a significant difference was observed in contamination with the microbial species. In addition, as shown in Table 4, the contamination of frankfurters and sausages by the microbial species is greater than that of pizza. It is noteworthy that factors such as indicators (health, sanitation, lack of hygiene), age, gender, and education level did not impact the outcome. In studies conducted in Iran and other countries, hygiene and food contamination by pathogenic bacteria has been evaluated. In a study conducted by Mirzabeygi et al in 2006 in western Tehran, the levels of contamination with *S. aureus* in dairy products and sweetmeats were reported as 16% (27). In a study by Sultan Dallal et al in 2007 surveying the microbiological quality of chicken and red meat, 47.8% of chicken samples and 28.8% of meat samples were contaminated with *Salmonella*, which is less than *S. arizonae* contamination in the present study (28). A study by Tavakoli et al was carried out in 4 restaurants affiliated with a military facility in Tehran. A total of 288 samples of ready-to-eat food and salad were examined. In 5 samples (1.73%), *E. coli* was found. The samples did not confirm *Salmonella* contamination. Also, the mean number of coliforms and *S. aureus* in some foods was found to be more than the standard. However, in this study, the rates of contamination with *E. coli* and *S. arizonae* were higher (29). In a study by Faramarzi et al, 642 food samples (58.33% salads and 9.84% dairy products) were found to be contaminated with *E. coli*, and 4.8% of sweetmeats were contaminated with *S. aureus*, which is consistent with the current study (30). Aycicek et al examined the rate of *S. aureus* contamination of ready-to-eat salads and foods in restaurants at military centers in Ankara, Turkey. They tested 512 samples of salads, pizzas, and a variety of meat foods. Their results indicated that 48 samples were contaminated with *S. aureus*, and in meat foods and salads, *S. aureus* contamination was significantly higher than in the other samples, which is consistent with the current study (31). In a study performed by Meldrum et al conducted in the United Kingdom on 1213 salad and 1208 sauce samples, results showed that 4.7% of salad samples and 5% of sauce samples were of lower than acceptable quality, and the presence of microbial *E. coli* and *S. aureus* were confirmed in them (32). In the current study, 27.77% of food samples were contaminated with *E. coli*. In the study of Sagoo et al, 3% of salad samples had confirmed contamination with *E. coli* (33). The presence of bacterial species (*S. aureus*, *S. sonnei*, *S. arizonae*, *E. coli*, and *E. faecalis*) in the three food groups studied indicates the lack of sufficient attention paid to food hygiene standards in their procurement centers. This may be due to the staff's lack of personal hygiene, contamination of raw materials, or secondary contamination. *E. coli* and *S. aureus* contamination in the studied samples is a serious warning to health officials. *S. aureus* in raw food is not a good competitor for other bacteria, but in cooked foods in which other microorganisms are destroyed, it is easily grown and creates contamination (34). The tendency of a lot of people to use ready-to-eat foods, the lack of proper

observance of environmental health and food processing equipment, the improper washing of hands and the long-term contact of these hands with food are the most important causes of food contamination and disease. The results of this study can be used for operators of fast food centers and also for healthcare providers in health centers. Making the staff of fast food centers aware of the dangers of these bacteria can allow them to act to reduce the bacterial contamination of ready-to-eat foods. The implementation of an HACCP system, another standard certification, and the training of operators at fast food centers can be very effective in preventing or reducing bacterial contamination. Also, a complementary study of bacterial contamination in food preparation, dining room, utensils and equipment used in processing and cooking and personal hygiene operators can be useful.

Conclusion

Based on the results of the current study, the prevalence of bacterial contamination of ready-to-eat foods can have an important role in infection and food poisoning. The spread of foodborne diseases, problems in the field of food hygiene and its importance in health and economics, a lack of awareness, and a lack of respect for the basic principles of health could be the main reasons for contamination and could endanger public health. A lot of people tend to consume fast food, and the bacterial contamination of these foods is a serious problem that needs further investigation. Training for personnel working in fast food preparation centers on the proper observance of health issues, and the overseeing of the preparation, transportation, storage, and supply of ready-to-eat foods seems necessary to prevent the transmission of microbial contamination. The amount of contamination in food must be reduced by taking appropriate actions to ensure the health of the community.

Acknowledgments

This article is the result of the research project number 22/52/93/1409, contract dated 29/11/93, approved by the Ilam University of Medical Sciences and Health Services. The authors are grateful for the financial support provided by the Deputy of Research and Technology of Ilam University of Medical Sciences.

Ethical issues

The authors certify that all data collected during the study is presented in this manuscript, and no data from the study has been or will be published separately.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

MA conceived and designed the study. MTY, MD, MM, FS, and ZG performed the literature search and wrote the manuscript. All authors participated in data acquisition, analysis, and interpretation. All authors critically

reviewed, refined, and approved the manuscript.

References

- Razavilar V. Pathogenic Microorganisms in Foods and Epidemiology of Food Borne Intoxications. 3rd ed. Tehran Tehran University Publications; 2010. [In Persian].
- Isara AR, Isah EC, Lofor PV, Ojide CK. Food contamination in fast food restaurants in Benin City, Edo State, Nigeria: Implications for food hygiene and safety. *Public Health* 2010; 124(8): 467-71. doi: 10.1016/j.puhe.2010.03.028.
- World Health Organization (WH). Food Safety and Food Borne Illness. Geneva: WHO; 2007.
- Newell DG, Koopmans M, Verhoef L, Duizer E, Aidara-Kane A, Sprong H, et al. Food-borne diseases-the challenges of 20 years ago still persist while new ones continue to emerge. *Int J Food Microbiol* 2010; 139 Suppl 1: S3-15. doi: 10.1016/j.ijfoodmicro.2010.01
- Abdollahzadeh E, Rezaei M, Hosseini H, Safari R. Effects of Nisin and thyme essential oil, individually and in combination, on inoculated populations of *Listeria monocytogenes* in minced silver carp. *Iran J Nutr Sci Food Technol* 2012; 6(4): 13-20. [In Persian].
- Kilic B. Current trends in traditional Turkish meat products and cuisine. *LWT- Food Science and Technology* 2009; 42(10): 1581-9. doi: 10.1016/j.lwt.2009.05.016.
- Teymori R, Ghazanfarirad N, Dehghan K, Asadzadeh J, Hajigholizadeh G, Bahmani M. A survey of bacteria and mold contamination of imported rice into West Azerbaijan Province, northwest of Iran. *Asian Pac J Trop Dis* 2014; 4 (Suppl 2): S833-5. doi: 10.1016/S2222-1808(14)60737-4.
- Kagkli DM, Vancanneyt M, Hill C, Vandamme P, Cogan TM. *Enterococcus* and *Lactobacillus* contamination of raw milk in a farm dairy environment. *Int J Food Microbiol* 2007; 114(2): 243-51. doi: 10.1016/j.ijfoodmicro.2006.09.016.
- Mosaferi M, Hajizadeh Y, Ostadrahimi AR, AslHashemi A. Importance of water quality control in food safety, case study: drinking, dairy and canning industries of eastern Azerbaijan. *Med J Tabriz Univ Med Sci Health Serv* 2007; 29(1): 93-97. [In Persian].
- Nascimento MS, Reolon EM, Santos ARB, Moreira VE, Silva N. *Enterobacteriaceae* contamination in chocolate processing. *Food Control* 2015; 47: 291-297.
- Garayoa R, Vitas AI, Díez-Leturia M, García-Jalón I. Food safety and the contract catering companies: food handlers, facilities and HACCP evaluation. *J Food Control* 2011; 22(12): 2006-12. doi: 10.1016/j.foodcont.2011.05.021.
- Vanderzant C, Splittstoesser DF. Compendium of methods for the microbiological examination of foods. 3rd ed. Washington, DC: American Public Health Association; 1992.
- Bolton DJ, Meally A, Blair IS, McDowell DA, Cowana C. Food safety knowledge of head chefs and catering managers in Ireland. *J Food Control* 2008; 19(3): 291-300. doi: 10.1016/j.foodcont.2007.04.006.
- Chapman B, Eversley T, Fillion K, Maclaurin T, Powell D. Assessment of food safety practices of food service food handlers (risk assessment data): testing a communication intervention (evaluation of tools). *J Food Prot* 2010; 73(6): 1101-7.
- Christison CA, Lindsay D, von Holy A. Microbiological surveys of ready-to-eat foods and associated preparation surfaces in retail delicatessens, Johannesburg, South Africa. *J Food Control* 2008; 19(7): 727-33. doi: 10.1016/j.foodcont.2007.07.004.
- Tavakoli HR, Karimi Zarchi AA, Izadi M. A survey on bacterial contamination of consumed foods in belonging centers of Baqiyatallah University of Medical Sciences. *Journal of Military Medicine* 2007; 9(2) 89-95. [In Persian].
- Salek Moghadam A, Forouhesh Tehrani H, Mozafari NA, Ansari H. Prevalence of virulence factors among *E. coli* isolated from food materials from Iran University of Medical Sciences' food microbial laboratory. *Feyz* 2000; 4(3): 32-40. [In Persian].
- Fadaei AM, Jamshidi A, Kheiri S. Comparison of bacterial contamination of raw and pasteurized milk used in Shahrekord in 2006. *Journal of Shahrekord University of Medical Sciences* 2008; 10(2): 37-44. [In Persian].
- Salek S. Microbial food intake of medical centers [dissertation]. Tehran: Shahid Beheshti University; 1999. [In Persian].
- Tavakoli H, Riazipour M. Microbial quality of cooked meat foods in Tehran University's restaurants. *Pak J Med Sci* 2008; 24(4): 595-9.
- Institute of Standards and Industrial Research of Iran. 2946. Microbiology of food and animal feeding stuffs - Detection and enumeration of presumptive *Escherichia coli* - most probable number technique. Karaj: Institute of Standards and Industrial Research of Iran; 2005. [In Persian].
- Institute of Standards and Industrial Research of Iran. Microbiology of food and animal feeding stuffs-horizontal method for the enumeration of positive staphylococci coagulase (number: 6806). Karaj: Institute of Standards and Industrial Research of Iran; 2005. [In Persian].
- Institute of Standards and Industrial Research of Iran. Microbiology of food and animal feeding stuffs - Detection and enumeration of presumptive *Escherichia coli* - Most probable number technique (number: 2946). 2nd ed. Karaj: Institute of Standards and Industrial Research of Iran; 2005. [In Persian].
- Institute of Standards and Industrial Research of Iran. Microbiology of food and animal feeding stuffs-horizontal method for detection of *Salmonella* (number: 1810). 3rd ed. Karaj: Institute of Standards and Industrial Research of Iran; 2005. [In Persian].
- Institute of Standards and Industrial Research of Iran. Microbiology of food and animal feeding stuffs horizontal method for the detection of *Shigella* spp (number: 2627). 2nd ed. Karaj: Institute Of Standards and Industrial Research of Iran; 2005. [In Persian].
- Institute of Standards and Industrial Research of Iran. Microbiology of foods and foodstuffs - Detection and enumeration of *Enterococcus* in food (number: 2198). Karaj: Institute of Standards and Industrial Research of Iran; 2005. [In Persian].
- Mirzabeygi M, Posti F, Rahbar Arasteh H. Study of

- Staphylococcus aureus* contamination in dairy and confectionery products from west of Tehran. Iranian Congress of Microbiology; 4-6 March 2008; Kerman. [In Persian].
28. Soltan Dallal MM, Sharifi Yazdi MK, Mirzaei N, Kalantar E. Prevalence of *Salmonella* spp. in packed and unpacked red meat and chicken in South of Tehran. *Jundishapur J Microbiol* 2014; 7(4): e9254. doi: 10.5812/jjm.9254 .
 29. Tavakoli HR, Farhang K, Karimi Zarchi A, Heydari E. Bacteriological quality of ready to eat food in four military restaurants. *Iranian Journal of Military Medicine* 2012; 13(4): 207-212. [In Persian].
 30. Faramarzi T, Jonidi Jafari A, Dehghani S, Mirzabeygi M, Naseh M, Arasteh RH. A survey of bacterial contamination of food supply in the West of Tehran. *Journal of Fasa University of Medical Sciences* 2012; 2(1): 12-18. [In Persian].
 31. Aycicek H, Cakiroglu S, Stevenson TH. Incidence of *Staphylococcus aureus* in ready-to-eat meals from military cafeterias in Ankara, Turkey. *J Food Control* 2005; 16(6): 531-4. doi: 10.1016/j.foodcont.2004.04.005.
 32. Meldrum RJ, Little CL, Sagoo S, Mithani V, de Pinna E. Assessment of the microbiological safety of salad vegetables and sauces from kebab take-away restaurants in the United Kingdom. *J Food Microbiol* 2009; 26(6): 573-7. doi: 10.1016/j.fm.2009.03.013.
 33. Sagoo SK, Little CL, Mitchell RT. Microbiological quality of open ready-to-eat salad vegetables: effectiveness of food hygiene training of management. *J Food Prot* 2003; 66(9): 1581-6.
 34. Young RE, Young MK, Yoon KS. Survival of pathogenic enterohemorrhagic *Escherichia coli* (EHEC) and control with calcium oxide in frozen meat products. *Food Microbiol* 2015; 49: 203-10. doi: 10.1016/j.fm.2015.02.010.