



Habitat destruction of local hedge plants of agricultural field by the modernization of agriculture in saoli region.

Watakhare Devilal N^{1*} and Kale MC²

¹Assistant Professor, RMG College Saoli

²Principal, Anand Niketan College, Warora

*Corresponding Author E-mail: dnwatakhare23@rediffmail.com

Manuscript details:

Available online on
<http://www.ijlsci.in>

ISSN: 2320-964X (Online)
ISSN: 2320-7817 (Print)

Editor: Dr. Arvind Chavhan

Cite this article as:

Watakhare Devilal N and Kale MC (2018) Habitat destruction of local hedge plants of agricultural field by the modernization of agriculture in saoli region, *Int. J. of Life Sciences*, Special Issue, A12: 204-206.

Copyright: © Author, This is an open access article under the terms of the Creative Commons Attribution-Non-Commercial - No Derives License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

ABSTRACT

Saoli region is basically an agriculture zone where majority of people survive on rice farming. In this area agricultural field is surrounded by the barbed wire fencing instead of natural fencing by hedge plants. There are several hedge plants which includes medicinal, ecological and wild fruit and vegetable plants. The present investigation focuses on the current status of hedge plants due to habitat destruction by modern agricultural techniques. The outcome of this research will open the debate on classical farming versus modern farming and also make awareness among the people about importance of hedge plants in nature.

Keywords: Saoli, Habitat, Hedge plants, Modernization, Barbed Wire Fencing.

INTRODUCTION

Habitat destruction occurs when a natural habitat, such as a forest or wetland, is altered so dramatically that it no longer supports the species it originally sustained. Plant and animal populations are destroyed or displaced, leading to a loss of biodiversity. Habitat destruction is considered the most important driver of species extinction worldwide (Pimm and Raven 2000). Humankind has dramatically transformed much of the Earth's surface and its natural ecosystems. This process is not new it has been ongoing for millennia but it has accelerated sharply over the last two centuries. Today, the loss and degradation of natural habitats can be likened to a war of erosion. Few habitats are destroyed entirely. Very often, habitats are reduced in extent and simultaneously fragmented, leaving small pieces of original habitat. In concert with habitat loss, habitat fragmentation is a grave threat to species survival (Laurance et al. 2002; Sekercioglu et al. 2002). Globally, agriculture is the biggest cause of habitat destruction. Other human activities, such as mining, clear cut logging, trawling, and urbanization also destroy or severely degrade habitats.

In developing nations, where most habitat loss is now occurring, the drivers of environmental change have shifted fundamentally in recent decades. Instead of being caused mostly by small-scale farmers and rural residents, habitat loss, especially in the tropics, is now substantially driven by globalization promoting intensive agriculture and other industrial activities. Destruction and fragmentation of natural habitats are the 2 most important factors in the current species extinction event (Groombridge 1992). Loss and fragmentation of habitat result in reduced population sizes, which increases the probability of extinction by demographic and/or environment (Burkey 1995).

A "hedge" is a living wall composed of plants around farm, garden and home lawns. A hedge is a line of closely spaced shrubs, climbers and sometimes trees, planted and trained to form a barrier or to mark the boundary of an area. The development of hedges over the centuries is preserved in their structure. The first hedges enclosed land for cereal crops during the Neolithic Age (4000–6000 years ago). Hedge is work as a decorative as well as security purpose. Hedge plants can give a higher level of security if farmers select shrubs or small trees that have thorns such as *Acacia sp.*, *Caesalpinia sp.*, *Bombax ceiba* etc. Hedges used to separate a road from adjoining fields or one field from another. Hedges also serve as wind breaks to improve conditions for the adjacent crops.

In India farmers were primarily planted natural hedge plants around the agriculture farm for security reason. as time goes on hedge plants uses increased from security to different purposes. Hedge plants are used by farmers as vegetables for instance several species of Fabaceae and Cucurbitaceae (*Lablab sp.*, *Memordica sp.*), for medicinal purpose - e.g. *Abrus*, *Acacia*, *Caesalpinia* etc., hedge plants is use as a fruit in villages for example *Zizyphus sp.*, as a timber plant e.g. *Acacia nilotica*, as fuel wood and uses goes on. Hedges acts as a wind breaks. Possibly the most salient point for many people, they work as an effective wind break but, unlike a fence, they allow some wind to pass through their foliage. When an 80mph wind blows. It bangs into a solid fence panel the pressure on the wood is immense. A hedge will allow that wind to pass through its foliage and slow it down which in turn protect the crops. Hedging is very vital for our wildlife, especially insects who need to travel from garden to garden. Fences stop these lovely creatures from getting to food, shelter and breeding sites but hedges allows it. Birds will take shelter in hedges and bees will collect pollen and nectar from

flowering varieties. Indirectly it increases the biodiversity of nature. Now a days due to modernization of agriculture, hedge plants around the farm is replaced by barbed wire fencing for higher security reason. it directly destruct the habitat of hedge plants. The goal of this study was to assess the habitat loss on population of hedge plant and extinction from Saoli town and nearby villages.

METHODOLOGY

Study Area:

Saoli is a tahsil in Chandrapur district of Maharashtra state. Location of Saoli is 20°06'41"N, 79°47'21"E. Saoli is come under the rice cultivation belt of Maharashtra. People of this village basically depend upon the rice farming. The main occupation is rice farming. Saoli is surrounded by large agricultural fields. The farming in this area is still not that much developed though the fencing of agriculture fields are barbed wire fence. It indirectly impact the habitat of hedge plants. The selected agricultural fields for study are from Saoli town, Malpiranji Village, Khedi Village, Sindola village, and Chakpiranji village. Figure 1 shows the location of the study area.




Fig 1: Location of study area

There are different methodologies proposed by ecologists for sampling of angiosperms. The most important and widely used method for a general assessment is belt transect method. The random sampling method in the selected area. The transect method was followed, and accordingly, transects or straight lines were marked starting from the base of the study area to the end of the Agricultural fields in each selected site. The length of a transect was 500 m to 1 km in each of the selected habitat. This is the standard scientific method followed by various workers in

Table

SN	Agriculture Site	GPS Location	Number of agriculture field surveyed in each site (500mt each)	Number of Hedge plant fencing present
01	Saoli	20°06'41"N 79°47'21"E	10	-
02	Chakpiranji	20°04'46"N 79°47'49"E	08	++
03	Khedi	20°04'31"N 79°46'40"E	07	++
04	Malpiranji	20°05'16"N 79°47'33"E	07	++
05	Shindola	20°04'19"N 79°48'57"E	08	-
Total	05		40	06

Note: +, - sign indicates the presence and absence of hedge plant fencing, number of + indicate The number of site where hedges present

respect of phytosociological studies (Cottam and Curtis 1956; Ralhan et al. 1982; Saxena and Singh 1982; Lu et al. 2004; Nautiyal 2008)

Observation: The main focus of this study is to assess the habitat of hedge plant is destroyed or not after doing transect method following observation has done:

Data Collection: The experimental site are selected randomly from each direction keeping in Saoli as center. Malpiranji is in north, Sandola is in south, Khedi is in west and Chakpiranji is in east with respect to Saoli as a center spot. Within each site agricultural field is surveyed. Number of field is depend upon the area of village. Total forty fields has been surveyed. Out of forty only six field has hedge plant fencing. The economical background of farmers also collected. Out of total maximum farmers has more than 5 acre of lands and those have a hedge plant fencing have less than 2 acre land.

RESULT AND DISCUSSION

The data indicates that loss of habitat of hedge plants is huge. Only six field out of forty has a natural hedge fencing it shows that marginalized farmers are still have natural hedge plant fence as they cannot afford barbed wire fencing. As the income of farmers raised this trend of natural hedge will decrease. The trend of use of modern equipment for agriculture is found in those villages which is nearer to city. The modernization of agriculture is causing more threat to habitat of hedge plants. This study shows the status of hedge plants near Saoli region.

REFERENCES:

- Pimm SL and Raven P (2000). Biodiversity: Extinction by numbers. *Nature*, 403: 843–845.
- Laurance WF, Lovejoy T, Vasconcelos H, et al. (2002). Ecosystem decay of Amazonian forest fragments: a 22-year investigation. *Conservation Biology*, 16: 605–618.
- Sekercioglu C H, Ehrlich PR, Daily GC, Aygen D, Goehring D, and Sandi R (2002). Disappearance of insectivorous birds from tropical forest fragments. *Proceedings of the National Academy of Sciences of the United States of America*, 99: 263–267.
- Groombridge B, Editor (1992). Global biodiversity: state of the earth's living resources. Chapman and Hall, New York, N.Y. 585.
- Burkey (1995). Extinction rates in archipelagos: implications for populations in fragmented habitats. *Conserv. Biol.* 9:527-541.
- Cottam G, Curtis JT (1956). The use of distance measures in phytosociological sampling. *Ecology* 37(3):451–460
- Ralhan PK, Saxena AK, Singh JS (1982). Analysis of forest vegetation at and around Nainital in Kumaun Himalaya. *Proc Indian Nat Sci Acad B* 48:121–137
- Saxena AK, Singh JS (1982) A phytosociological analysis of woody species in forest communities of a part of Kumaun Himalaya. *Vegetatio* 50(1):3–32
- Lu D, Mausel P, Brondizio E, Moran E (2004) Relationship between forest stand parameters and landsat TM spectral responses in the Brazilian amazon basin. *For Ecol Manage* 198:149–167
- Nautiyal S (2008) Structure of Central Himalayan forests under different management regimes: an empirical study. Working paper No 206. Institute for Social and Economic Change, Bangalore.

© 2018 | Published by IJLSCI