



---

## Nutritional profiling and free radical scavenging potentials of Garden egg stalk

Ogunka-Nnoka C.U., Ekrika D.D.

Department of Biochemistry, University of Port Harcourt, Choba Rivers State, Nigeria

**Abstract** This study was carried out to evaluate the nutritional properties of *Solanum macrocarpum* (garden egg) stalk using standard methods. Sulphates and chloride were the most abundant anions while calcium and magnesium were the predominant mineral cations. Results for the proximate composition showed high carbohydrates (27.70 %), fibre contents (22.94 %), and ash (18.83 %) while the phytochemical composition showed higher ribalinidine, catechin, and phenol contents. Vitamin B<sub>6</sub>, C, A, and B<sub>12</sub> were found in higher concentrations, while lysine (9.48 g/100g), phenylalanine (6.86 g/100g), leucine (6.59 g/100g), and arginine (5.66 g/100g) were the predominant essential amino acids. Results for the free radical scavenging potentials showed higher DPPH, nitric oxide, hydrogen peroxide, and ABTS radical scavenging potentials for the garden egg stalk, while their ferric reducing antioxidant power, superoxide and lipid peroxide scavenging were comparable to the standards used. This study has shown the high nutrient density of the garden egg stalk, thus encouraging its inclusion for dietary uses.

**Keyword:** Garden egg stalk; phytochemicals; micronutrients, amino acids; proteins

---

### 1. Introduction

Vegetables are useful components of the diet due to their contributions in forming a balanced diet. They are found among a vast amount of plant species that can serve as readily available proteinous sources and herbs. The efficacies of various kinds of herbs have been widely explored, and because of their distribution in different localities have availed the opportunity for traditional herbalists to ensure affordable treatment of ailments. Notwithstanding, the explosion in population have necessitated an extensive research for undiscovered or underexploited herbs. *Solanum* is a widely dispersed genus of the Solanaceae family with over a thousand species, a tenth of which are indigenous African species. The dominant species are *S. macrocarpon* and *S. aethiopicum* [1] and are commonly called garden eggs or locally known as Dauta in Hausa, Igbagba in Yoruba, and Afufa in Ibo. Garden eggs are highly regarded part of the regular daily diet and medicament in Nigeria. In other cases, egg plants are of cultural significance as they are used to show fruitfulness and blessings, and are frequently offered as kola to show goodwill during visits, ceremonies and other social gatherings. They can be consumed fresh, fried, and raw for soups, stew and sauce preparations. Bello *et al.*, [2] have reported their usage as medicament for weight loss, catarrh, rheumatism, skin diseases, digestive difficulties, and some allergies. In addition, Igwe *et al.*, [3] and Odetola *et al.*, [4] observed that egg plants possess potent glucose lowering, anti-inflammatory, cholesterol lowering and asthma relieving properties. Obviously, both the fruit and leaves of egg plants have been extensively researched on, but no study is yet to be undertaken on the nutrient properties of the stalks attached to the fruit which are usually discarded. This study was thus carried out to explore the nutrient density of the egg plant stalk.



## 2. Experimental Work

### 2.1. Sample collection and identification

Fresh garden egg (*Solanum macrocarpum*) fruits were bought from Choba Market, Port Harcourt, Rivers State. The stalk were collected and identified at the Department of Plant Science and Biotechnology, University of Port Harcourt, Choba, Rivers State.

### 2.2. Proximate composition and phytochemical analysis

Proximate content determination was carried out by the method of Ogunka-Nnoka *et al.* [5]. The method of Agomuo *et al.* [6] and as modified by Agomuo and Amadi, [7] was adopted for the chromatographic isolation of the phytoconstituents. A Bulk Scientific chromatography machine, equipped with a Hp-88 capillary column of 100m x 0.25mm dimension was used under some of the following conditions; detector type was flame ionization detector, split injection (1ml) mode at 220<sup>0</sup>C, detector & oven temperature were 250<sup>0</sup>C & 180<sup>0</sup>C respectively, and hydrogen as carrier gas. An internal standard was used for identification of the phytochemicals.

### 2.3. Determination of vitamins and minerals

For the vitamins determination, a 0.10g homogenized samples was pressed and 1.0ml of the extracts was transferred to a HP 6890N model of gas chromatography machine with a pulse flame photometric detector and a HP-5 30m x 25mm x 25µm column. A split injection mode of 20:1 was used for nitrogen as the carrier gas with a 1.0ml/min flow rate. For the oven conditions, 50<sup>0</sup>C was the initial temperature for 2min, and maintained the detector a 320<sup>0</sup>C. The mineral contents were estimated by the method of Amadi *et al.*, [8] by wet digestion of the samples in a mixture of Conc. nitric acid and HClO<sub>4</sub> (60% v/v) and afterwards estimated using an atomic mass absorption spectrophotometer. Sulphate was measured by turbidimetry, while chloride was by titrimetry.

### 2.4. Analysis of amino acids contents

Briefly, hydrolysates (150 µl) obtained following the method of Moore and Stein [9] was injected into the separation column of a Sykam-S7130 model of HPLC analyzer as described by Amadi *et al.*, [10]. The areas of the standards were used to determine the percentage amino acid composition of the sample. The amount of tryptophan was determined calorimetrically from alkaline hydrolysis of the sample according to Rama Rao *et al.* [11].

### 2.5. Determination of the antioxidant properties

2,2-azino-bis(3-ethylbenzthiazoline-6-sulphonic acid) (ABTS) radical scavenging assay was carried out as described by Agomuo and Amadi [7] at a wavelength of 734nm while 1,1-diphenyl-2-picrylhydrazyl (DPPH) radical scavenging assay was carried out at 517nm. The determination of ferric reducing antioxidant power (FRAP) was at a wavelength of 700nm following the procedure of Sutharsingh *et al.*, [12] while the method of Fontana *et al.*, [13] was adopted for the determination of superoxide radical scavenging assay. Nitric oxide scavenging activity was determined at a wavelength of 546nm with Greiss reagent as standard while the determination of lipid peroxidation inhibition potentials were in accordance with the method of Ruberto and Barrata [14].

## 3. Results

**Table 1:** Proximate composition of egg plant stalk

Proximate composition	Contents (%)
Ash	18.83±0.44
Moisture	9.43±1.65
Protein	15.41±1.99
Fat	5.68±0.31
Fibre	22.94±2.19
Carbohydrate	27.70±2.61

Values represent means ± S.D of duplicate determinations



**Table 2:** Phytochemical composition of egg plant stalk

Parameters	Concentration ( $\mu\text{g/g}$ )
Anthocyanin	0.37
Oxalate	0.79
Sapogenin	4.41
Rutin	4.42
Phenol	12.34
Saponin	7.05
Lunamarin	1.49
Ribalinidine	20.55
Phytate	1.13
Kaempferol	7.41
Catechin	13.46

**Table 3:** Vitamin composition of egg plant stalk

Parameters	Concentration ( $\mu\text{g/g}$ )
Vitamin A	21.87 $\pm$ 1.26
Vitamin D	9.63 $\pm$ 0.27
Vitamin E	7.82 $\pm$ 0.04
Vitamin C	46.26 $\pm$ 1.32
Vitamin B <sub>1</sub>	0.36 $\pm$ 0.04
Vitamin B <sub>2</sub>	0.34 $\pm$ 0.04
Vitamin B <sub>3</sub>	0.71 $\pm$ 0.05
Vitamin B <sub>6</sub>	116.48 $\pm$ 5.62
Vitamin B <sub>12</sub>	13.41 $\pm$ 0.65

Values represent means  $\pm$  S.D of duplicate determinations

**Table 4:** Mineral contents of egg plant stalk

Minerals	Content (mg/kg)
Sodium	7.91 $\pm$ 0.02
Potassium	9.16 $\pm$ 0.10
Magnesium	22.13 $\pm$ 0.22
Zinc	4.85 $\pm$ 0.12
Calcium	41.41 $\pm$ 1.53
Iron	0.25 $\pm$ 0.02
Copper	0.14 $\pm$ 0.02
Phosphorus	8.61 $\pm$ 0.50
Sulphate	111.12 $\pm$ 6.43
Chloride	173.24 $\pm$ 53.9

Values represent means  $\pm$  S.D of duplicate determinations

**Table 5:** Amino acid content of egg plant stalk

Components	Concentration (g/100g)
<b>Essential amino acids</b>	
Lysine	9.48
Histidine	5.35
Arginine	5.66
Methionine	1.54
Threonine	4.28
Isoleucine	4.49



Leucine	6.59
Valine	2.47
Tryptophan	1.24
Phenylalanine	6.86
<b>Non Essential amino acids</b>	
Glycine	3.89
Alanine	4.06
Serine	3.84
Proline	3.91
Aspartate	12.21
Glutamate	14.69
Tyrosine	2.97
Cysteine	3.38
Total Amino Acids	96.91
Total Essential Amino Acids	47.96
Total Non-Essential Amino Acids	48.95




Figure 1: DPPH radical scavenging potentials of garden egg stalk




Figure 2: Nitric oxide scavenging potentials of garden egg stalk






Figure 3: Hydrogen peroxide scavenging potentials of garden egg stalk




Figure 4: ABTS radical scavenging potentials of garden egg stalk




Figure 5: Ferric reducing antioxidant potentials of garden egg stalk




Figure 6: Superoxide anion scavenging potentials of garden egg stalk






Figure 7: Fatty acid peroxy radical scavenging potentials of garden egg stalk

#### 4. Discussion

The result for the proximate composition was presented in Table 1. The ash, moisture, and protein contents were 18.83%, 9.43%, and 15.41% respectively, while the fat, fibre and carbohydrate contents were 5.68%, 22.94%, and 17.70% respectively. The moisture content provides an indication as to its dry matter content, its susceptibility to microbial infestation, and the freshness of the food material. From the results of this study, the eggplant stalk contains poor moisture content, which was lower than the eggplant fruit itself [15]. The ash content directly relates to the amount of minerals in a food material, while the fibre content indicates the extent of gastric emptying and intestinal bulk, derivable from consuming the food material. Both the ash and fibre contents of the eggplant stalk in this study were greater than that of the fruit itself [15] as well as those for some other vegetables reported by Asaolu *et al.*, [16]. According to Ali, [17], food materials providing above 12% of proteins can be regarded as sufficient protein sources. This implies that the eggplant stalk is a sufficient protein source. The fat contents of the eggplant in this study compared with those of leafy vegetables [18], while the result of this study further indicated that the eggplant is a better carbohydrate source when compared to those for its fruit varieties; *S. aethiopicum* and *S. macrocarpum* [15].

Table 2 shows the phytochemical composition of eggplant stalk. Ribalinidine, catechin, phenols and kaempferol (20.55, 13.46, 12.34, and 7.41 µg/g) were the predominant phytochemicals in the eggplant stalk, while anthocyanin, oxalates, and phytates were the least abundant phytochemicals. Ribalinidine and lunamarine are recognized alkaloids with antimalarial and analgesic effects [6], hence, the eggplant stalk could potentially possess effective antimalarial and analgesic properties. In addition, the abundance of catechins and other phenols in the eggplant, implies that this plant material could be useful in management of blood pressure and cardiovascular diseases [19]. Also, coupled to the predominance of phenols, the kaempferol content of the eggplant stalk ensures its effectiveness in combating oxidative stress [20]. The low levels of the antinutrients; oxalates and phytates is an indication of higher mineral bioavailabilities.

Table 3 reveals the vitamin compositions of the egg plant stalk. The fat soluble vitamins; A, D, E were 21.86, 9.63, and 7.82 µg/g respectively, while vitamin B<sub>6</sub> (116.48 µg/g), C (46.26 µg/g), and B<sub>12</sub> (13.41 µg/g) were the most abundant water soluble vitamins. The results imply that all the fat soluble vitamins were below their respective RDA's of 900, 10, and 15 µg [5]. Vitamin A and E are potent antioxidants while vitamin D is essential for the maintenance of bone health, hence, the eggplant requires supplementation with richer sources of these vitamins on account of dietary usage. Notwithstanding that the amounts of the water soluble vitamins found in the eggplant stalk were lower than their recommended daily allowance, their levels presented in this study, were greater when compared to some plants with notable health benefits [21].

The mineral contents of egg plant stalk were presented in Table 4. The minerals ranged from 5073 mg/kg for the chloride content to 0.14 mg/kg for the copper level. The anions; chloride and sulphate were the most abundant minerals found in the egg plant stalk, followed by calcium and magnesium. The sodium and potassium contents of the eggplant stalk were greater than those reported for the eggplant fruit [22]. Also, the magnesium, zinc, and calcium contents of the eggplant stalk were lower than those found for commonly consumed vegetables like bitter



leaf and *Amaranthus hybridus* [16], while the iron, copper and phosphorus levels found for the eggplant stalk, were below the recommended daily intake [22]. The deficiencies regarding the amounts found for these minerals contained in the eggplant stalk in this study, implies that it requires supplementation with richer nutrient sources, for dietary purposes.

The amino acid composition of the eggplant stalk was shown in Table 5. Lysine (9.48 g/100g), phenylalanine (6.86 g/100g), leucine (6.59 g/100g), arginine (5.66 g/100g), and histidine (5.35 g/100g) were the predominant essential amino acids, while glutamate (14.69 g/100g) and aspartate (12.21 g/100g) were the most abundant of the non-essential amino acids. These amino acids were higher than those of the eggplant fruit, except for leucine and phenylalanine contents [23]. In addition, the threonine, isoleucine and tryptophan contents of the eggplant stalk in this study were higher than the WHO reference amino acid values [23]. Similarly, the eggplant stalk produced higher individual non essential amino acids than popularly consumed vegetables such as *A. cruentus*, *V. amygdalina*, and *T. triangulare* [24]. This implies that the eggplant is richly supplied with essential amino acids, and could serve as a dependable protein source.

Fig. 1-7 shows the in-vitro antioxidant potentials of the eggplant stalk. The result indicated a concentration dependent free radical scavenging potentials of the eggplants at all the assay carried out. The DPPH (Fig. 1), nitric oxide (Fig. 2), hydrogen peroxide (Fig. 3), and ABTS (Fig. 4) scavenging potentials of the garden egg stalk were higher than the standard ascorbic acid, while the  $Fe^{3+}$ , superoxide, and lipid peroxide scavenging potentials of the eggplant stalk as shown in Fig 5, 6, and 7 respectively, were equivalent to the standards used. The significant DPPH and ABTS scavenging potentials of eggplant stalk shown in this study, testifies to the potent hydrogen donating capacities of their phytoconstituents, and in addition, their nitric acid and hydrogen peroxide scavenging potentials directly implicate rich supplies of phenolic acids in the eggplant stalk [25].

## Conclusion

The nutritional evaluation of *S. macrocarpum* stalk in this study showed appreciable amount of proximate contents, micronutrients, and phytochemicals, while its free radical scavenging potentials further encourages its usage for relief of oxidative stress. Thus, this study has provided the needed scientific backing for the exploitation of garden egg stalk for dietary inclusion.

## Conflict of interest

None declared.

## References

1. Bonsu, K.O., Fontem, D.A., Nkansah, G.O., Iroume, R.N., Owusu, E.O. and Schippers, R.R. (2002) Diversity within the Gboma eggplant (*Solanum macrocarpon*), an indigenous vegetable from West Africa. *Ghana J. Horticulture*, 1, 50–58.
2. Bello, S.O., Muhammad, B.Y., Gammaniel, K.S., Abdu-Aguye, I., Ahmed, H., Njoku, C.H., Pindiga, U.H. and Salka, A.M. (2005) Preliminary Evaluation of the Toxicity and Some Pharmacological Properties of the Aqueous Crude Extract of *Solanum melongena*. *Res. J. Agric. Biol. Sci.*, 1(1), 1-9.
3. Igwe, C.U., Ibegbulem, C.O., Nwaogu, L.A., Ujowundu, C.O. and Okwu, G.N. (2013). Calcium, Zinc and phytate interrelationships in four lesser-known African seeds processed into food condiments. *J Adv. Chem.* 4(2): 288-292.
4. Odetola, A.A., Iranloye, Y.O. and Akinloye, O. (2004) Hypolipidaemic Potentials of *Solanum melongena* and *Solanum gilo* on Hypercholesterolemic Rabbits, *Pak J. Nutri.*, 3 (3), 180-187.
5. Ogunka-Nnoka, C.U., Amadi, P.U., Ikediashi, E.C. Nchendo, A. and Ugochukwu, P. (2017). Nutrient composition of *Dacryodes edulis* seed and seed coat mixture C.U. *PhytoChem & BioSub Journal*, 11(2), 81-88. DOI:10.163.pcbjsj/2017.11.-2.-81



6. Agomuo Emmanuel, Amadi Peter, Ogunka-Nnoka Charity, Amadi Benjamin, Ifeanacho Mercy, and Njoku Uche (2017). Chemical Composition of *Duranta repens* Leaf and Seed Oils. OCL Lipids, Oil Seeds & Crops, 24(6), 1-8. Doi: doi.org/10.1051/ocl/2017048
7. Agomuo, E.N. and Amadi, P.U. (2018). Nutritional and antioxidant properties of oils from selected bagasse, agricultural residues, and fodders. Journal of American College of Nutrition, Doi.org/10.1080/07315724.2018.1484307
8. Amadi, B.A., Peters, D.E., Agomuo, E.N. Amadi, P.U. and Denson, G.O. (2018) Nutrient composition of some selected diets of Ijaw people of Bayelsa state. Polish Journal of Natural Sciences, 33(1):59-74
9. Moore, K.J., Jung, H.G. 2001. Lignin and fiber digestion, J. Range Manage. 54, 420–430
10. Amadi, B., Ogunka-Nnoka, C., Ogaji, M. and Amadi, P. (2017) Nutrient Composition and Biochemical Effects of *Cocos nucifera* Products on Alloxan-induced Diabetic Wistar Rats. International Journal of Medicine, 5(2), 149-157. doi: 10.14419/ijm.v5i2.7647
11. Rama Rao, M.V., Tara, M.R. and Krishnan, C.K. (1974). Colour imetric estimation of tryptophan content of pulses. Journal of Food Science and Technology, (Mysore), 11: 13-216
12. Sutharsingh, D.C.R., Kavimani, S., Jayakar, B., Uvarani, M. and Thangathirupathi, A. (2011). Quantitative phytochemical estimation and antioxidant studies on aerial parts of *Naravelia zeylanica*. Int J Pharm Stud Res, 2(2), 52-56
13. Fontana, L., Cummings, N.E., Arriola, A., Sebastian, I., Neuman, J.C. and Kasza, I. (2016). Decreased consumption of branched-chain amino acids improves metabolic health. Cell Reports, 6:520–530. doi:doi.org/doi:10.1016/j.celrep.2016.05.092
14. Ruberto, G. and Barrata, M.T. (2000). Antioxidant activity of selected essential oil components in two lipid model system. *Food Chem* 69:167-74.
15. Nwodo, S.C., Abayomi, C.O., Eboji, O.K., Emiloju, O.C., Arinola, O.K. and Dania, D.I. (2011). Proximate and Phytochemical Analyses of *Solanum aethiopicum* L. and *Solanum macrocarpon* L. Fruits, Research Journal of Chemical Sciences 1(3), 63-71
16. Asaolu, S.S., Adefemi, O.S., Oyakilome, I.G., Ajibulu, K.E. and Asaolu, M.F. (2012) Proximate and Mineral Composition of Nigerian Leafy Vegetables, Canadian Center of Science and Education. Journal of Food Research; 1(3) 214-218
17. Ali, A. (2009). Proximate and mineral composition of the marchubeh (*Asparagus officinalis*). World Dairy and Food Science, 4(2), 142-149.
18. Udosen, E.O. (1995). Proximate and mineral composition of some Nigerian Vegetables. Discov. Inov., 7, 383-386.
19. Khalesi, S. (2014). Green tea catechins and blood pressure: a systematic review and meta-analysis of randomised controlled trials. Eur J Nutr 536:1299–1311.
20. Paolillo, R., Carratelli, C.R. and Rizzo, A. (2011). Effect of resveratrol and quercetin in experimental infection by *Salmonella enteric* Serovar typhimurium. International Journal of Immunopharmacology, 11:149-156.
21. Ikewuchi, C.C. and Ikewuchi, J.C. (2009) Comparative Study of the Vitamin Composition of Some Common Nigerian Medicinal Plants. 10 (1) 367-371.
22. Eze, S.O. and Kanu, C.Q. (2015). Assessment of the Phytochemical, Proximate, Vitamin and Mineral Composition of *Solanum gilo* L. International Research Journal of Pure & Applied Chemistry 5(1): 83-90.
23. Sodamade, A., Bolaji, O.S. and Owonikoko, A.D. (2015). The Nutritive Value and Amino Acid Characteristics of *Solanum Aethiopicum* Leaf Protein Concentrates. International Journal of Advanced Research in Chemical Science (IJARCS) 2(6), PP 28-33
24. Olubunmi, A.O., Olorunfemi, O., and Akinyeye, R.O. (2015) “Amino Acid Composition of Ten Commonly Eaten Indigenous Leafy Vegetables of South-West Nigeria.” *World Journal of Nutrition and Health*, 3(1) 16-21. doi: 10.12691/jnh-3-1-3.





25. Aluko, B.T., Oloyede, O.I and Afolayan, A.J. (2013). Polyphenolic contents and free radical scavenging potential of extracts from leaves of *Ocimum americanum* (L.). *Pakistan Journal of Biological Sciences*, 16(1): 22-30.

