



Original Article

Asian Pacific Journal of Reproduction

Journal homepage: www.apjr.net



doi: 10.4103/2305–0500.311615

Aphrodisiac potential of *Polyalthia bullata* (Tongkat Ali) in fowlJaya Vejayan¹✉, Yasmin Amira Che Yahya¹, Srikumar Chakravarthi², Halijah Ibrahim³, Aida Yun⁴¹Faculty of Industrial Sciences & Technology, Universiti Malaysia Pahang, Lebuhraya Tun Razak, 26300, Gambang, Kuantan, Pahang Darul Makmur, Malaysia²Faculty of Medicine, Biomedical Sciences and Nursing, MAHSA University, Jalan SP2, Bandar Saujana Putra, 42610 Jenjarom Selangor, Malaysia³Institute of Biological Sciences, University of Malaya, 50603 Kuala Lumpur, Malaysia⁴Ubat Tradisional Orang Asli, Dataran Sungai Perak Sebelah Utara, North–South Expressway, Kampung Menora, 33000 Kuala Kangsar, Perak Darul Ridzuan, Malaysia

ABSTRACT

Objective: To study the aphrodisiac potential of *Polyalthia (P.) bullata* in fowl.**Methods:** In this study, testosterone, as an indicator of the aphrodisiac potential of *P. bullata*, was investigated for its release from TM3 Leydig cells grown *in vitro* and in 4 fowls given capsules containing *P. bullata* at a dose of 10 mg in each capsule twice a day, for 50 days. In the latter *in vivo* evaluation, mating behaviours were additionally determined after the treated fowls were released to the individual hens, and their testes and liver were dissected for histological examinations. Blood drawn from the fowls was assessed for any changes in diagnostic parameters.**Results:** In the *in vitro* test (TM3 Leydig cells), *P. bullata* was able to increase testosterone to 0.48 nmol/L within 72 h of incubation, compared to the untreated control with only 0.18 nmol/L, *i.e.*, an increase of 170%. In the *in vivo* test, outcomes in the fowls dosed with *P. bullata* showed similar positive elevations of testosterone to (9.72±1.10) nmol/L in comparison to the controls that showed a level of only (4.05±0.84) nmol/L. Total frequencies of mating behaviours were observed (wing flapping, body shakes, crowing and beak pecking) to be 23 counts for the test compared to only 15 for the control fowls. Histological examination of the male reproductive organs provided evidence of testosterone boosting based on an observable increase in the activity at the seminiferous tubules of testis tissues without any damaging effects, compared to the controls. In the nine diagnostic blood parameters assessed, including alanine aminotransferase, aspartate aminotransferase, and gamma glutamyltransferase, none was remarkably elevated compared to the controls. The histological changes in the liver were not severe and mainly consisted of only localized moderate but recoverable obstructions and swellings of the vessels and tubules.**Conclusions:** *P. bullata* is able to boost testosterone both *in vitro* and *in vivo*, with no acute toxicities.**KEYWORDS:** Male hormone; Testosterone; Steroids; Leydig cells; Domestic chicken; Cock; Medicinal plant; *Polyalthia bullata*

1. Introduction

The use of aphrodisiac plants dates back centuries and includes plants either in their original form or made into consumable beverages, food or supplements for improving one's libido and sexual performance[1]. Not surprisingly, men are more inclined to resort to the use of aphrodisiacs. Many plants have been used for this purpose, with some being more popular than others in certain countries. Among them are *Microstylis wallichii* (Jeevak) and *Withania somnifera* (Ashwagandha) in India[2], *Butea superba* in Thailand[3], *Cynomorium coccineum* (Desert Thumb), *Prunus amygdalus* (almond) and *Zingiber officinale* (ginger) in Arab nations[4], and *Epimedium koreanum* (*Epimedium* leaf), *Dioscorea polystachya* (Chinese potato), *Ophiocordyceps sinensis* (caterpillar fungus), *Panax ginseng*, *Ginkgo biloba* and *Ganoderma lingzhi* in China and other pan-Asian countries[5]. In Malaysia, many plants have also been claimed to be aphrodisiacs, with some popularly used traditionally by the Malays and the indigenous people residing in the forest. Only a few of the aphrodisiac plants endemic to Malaysia have gone beyond the boundaries of its borders and been popularized internationally, including *Eurycoma (E.) longifolia* (Tongkat Ali) and *Smilax myosotiflora* (Ubi Jaga) for men and *Labisia pumila* (Kacip Fatimah) for women. Not known to many users of *E. longifolia* is that the term “Tongkat Ali” is synonymous with at least three plants in Malaysia. Perhaps the synonym originated through cultural practices over the years in defining

✉To whom correspondence may be addressed. E-mail: jayavejayan@ump.edu.my

This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial-Share Alike 4.0 License, which allows others to remix, tweak and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

©2021 Asian Pacific Journal of Reproduction Produced by Wolters Kluwer- Medknow. All rights reserved.

How to cite this article: Vejayan J, Yahya YAC, Chakravarthi S, Ibrahim H, Yun A. Aphrodisiac potential of *Polyalthia bullata* (Tongkat Ali) in fowl. *Asian Pac J Reprod* 2021; 10(2): 75–81.**Article history:** Received: 10 October 2020; Revision: 12 December 2020; Accepted: 25 December 2020; Available online: 29 March 2021

any plant with roots with a peculiar shape and claims of being an aphrodisiac. The three most widely used plants include *E. longifolia*, *Polyalthia (P.) bullata* and *Stema (S.) tuberosa*[6–8].

Of the three plants, *P. bullata* is the main focus of this study due to a lack of scientific investigations on its aphrodisiac claims. The whole plant is rarely seen in the modern landscapes of Malaysia but rather is confined to wild growth in forests located at moderate altitudes below 1 200 m[9]. The distinctive differences between the more popular Tongkat Ali, i.e., *E. longifolia*, and the lesser known *P. bullata* roots, where the colour is markedly visible as white or fairer for the former (hence simply termed as Tongkat Ali Putih in Malay), while *P. bullata* grows with black roots (Tongkat Ali Hitam)[10]. The root of *P. bullata* has been sold worldwide mostly in its original form of dry cut chips to users by e-commerce with claims of increasing sexual performances once its boiled preparations have been consumed regularly[7,11]. At times, this plant has been combined with other plants claimed to be aphrodisiacs, including the two other Tongkat Ali plants, i.e., *E. longifolia* and *S. tuberosa*, to become products of premixed coffee, capsules or tablets and other beverages or preparations[12,13]. Under such conditions, it has not been possible to ascertain whether the action of the products is due to this plant or merely due to the more established libido and testosterone boosting plant of *E. longifolia*. The latter has been well studied both clinically and in laboratory environments[13–18]. A study in 64 human subjects with confirmed moderate stress levels that were given *E. longifolia* extract for 4 weeks showed an increased testosterone level (+37%) and reduced cortisol level (-16%) compared to placebo[19].

P. bullata may not be a mere pseudosexual enhancer, since it has been well-known that this plant has been sold to repeat buyers and users worldwide for many years. As previous reports have almost no data on its aphrodisiac potential, this current *in vitro* and *in vivo* investigation was undertaken. This work used domesticated fowls similar to the study previously done for the other two Tongkat Ali plants i.e. *E. longifolia* and *S. tuberosa*[10].

2. Materials and methods

This work was conducted for a duration of 10 months (April 2019 through January 2020).

2.1. The *in vitro* Leydig cell evaluations

2.1.1. Extraction of crude plant samples

The highest quality roots of *P. bullata* were acquired by the indigenous Temiar group from the Perak forest, Malaysia, near Kampung Orang Asli Bukit Cermin (specimen deposited in University of Malaya with voucher number of HI1445). The group of 8 to 11 families belonging to a larger Senoic cluster have for generations harvested the thick tropical jungle of Malaysia for medicinal plants for their own use and as their livelihood in selling to herbal manufacturers involved in encapsulating them into registered products.

The raw Tongkat Ali (*P. bullata* and *E. longifolia*) was obtained in

the form of roots of various sizes and lengths. The roots were cut by using an electric saw into smaller pieces, followed by a slicing step that transformed the plant samples into chips. The chips were oven-dried at a temperature of 50 °C with convection. Tongkat Ali chips were turned into powder with a blender. The choice of the positive control for the *in vitro* studies was based on many reports on *E. longifolia* being a testosterone booster. Fifty grams of each powdered plant material was mixed with 500 mL of water and refluxed for 6 h. Upon completion, the particulates were filtered, and the filtrate was freeze-dried and kept until further use.

2.1.2. Growth of Leydig cells

TM3 Leydig cells (mouse origin), purchased from the US-based ATCC, were cultured at 37 °C in a 5% CO₂ incubator with complete medium consisting of Dulbecco's modified Eagle's medium/nutrient mixture F-12 medium (Nacalai Tesque, JP), 5% horse serum (Thermo Scientific, UK), 2.5% fetal bovine serum (Thermo Scientific, UK) and 1% penicillin-streptomycin (Nacalai Tesque, JP). The cells were allowed to reach approximately 80% confluence before passaging. Passaging of the cells was carried out according to Erasmus with slight modifications[20]. Briefly, the culture media was discarded, and the cells were washed with 4 mL Dulbecco-phosphate buffered saline (Nacalai Tesque, JP). Next, 1 mL of 10 × diluted 0.25% (w/v) trypsin with 0.53 mM ethylene diamine tetraacetic acid (Nacalai Tesque, JP) was added and incubated at 37 °C with 5% CO₂ for 5 min until the cells detached. A volume of 2 mL complete medium was added, and the cells were centrifuged at 730 × *g* for 5 min at 37 °C. Thereafter, the cell sediment (after discarding the supernatant) was resuspended in 1 mL complete medium, and 0.25 mL of the suspension was mixed with 4.75 mL of complete medium.

2.1.3. Evaluation of testosterone production

The cells (4 × 10³ cells/well) were seeded in a 96-well plate, treated with 50 µg/mL of *P. bullata* (test) and *E. longifolia* (a positive control) and incubated at 37 °C with 5% CO₂ for 24 h. After 72 h, the production of testosterone was assessed by using a testosterone enzyme linked immunosorbent assay kit according to the manufacturer's instructions (Elabscience, US). Briefly, the absorbances obtained at 450 nm were converted into the corresponding testosterone concentration with a plotted testosterone standard curve. The percentage of testosterone secretion was calculated with the following formula:

$$\text{Testosterone secretion (TS) (\%)} = \frac{\text{Sample TS} - \text{Control TS}}{\text{Control TS}} \times 100$$

2.2. *In vivo* evaluations

2.2.1. Fowl (*Gallus gallus*)

The animals selected for this experiment were domesticated red junglefowl (*Gallus gallus*), a type of rooster commonly found in Malaysia provided by Sapphire Enterprise (specialized in supplying experimental animals). The period of acclimatization was approximately 10 days following delivery. The control and test groups consisted of four fowls each. The fowls were 25–27 weeks

of age, *i.e.*, prior to reaching their fertile maturity (usually 28–38 weeks of age). The weights of the fowl ranged from 1.3–2.0 kg. The experiment conducted in controlled environment as indicated within the Animal Institutional Care and Use Committee approval of UMPIACUC_2018_01. The fowls were housed in cages at Experimental Animal Laboratory of Universiti Malaysia Pahang and fed *ad libitum* of chicken feed and drinking water.

2.2.2. Capsule preparation

The dosing was performed by using capsules containing the active ingredient of finely powdered *P. bullata* roots. A commercially available capsule maker used number “3” sized capsules to generate dosages of approximately 10 mg per capsule (adjusted to the average weight of fowl, based on capsules containing 250 mg of active ingredient per Bio-Root Tok Ali capsule that were given to adults weighing 70–90 kg). The type “3” capsule was able to contain up to 200 mg, and hence any empty space was filled with corn bran (blended chicken feed).

2.2.3. Dosing

The test and control fowl groups were made to swallow one capsule twice a day for a total of 50 days, a duration randomly selected considering no discernible increase of testosterone level at 30 days dosing was achieved (results not included). The fowls in the test group were given capsules containing 10 mg *P. bullata*, while the controls were given capsules with only blended chicken feed. The dose selected was similar to that used in the investigations of other Tongkat Ali plants in fowls[10]. The delivery of the capsule was performed by inserting it on the tongue of the forced opened beak, followed by induction of swallowing with a supply of water squirted gently and shutting the beak tightly. Consequently, the fowl was inspected by opening the mouth for successful dosing.

2.2.4. Evaluating the mating behaviours

Upon completing the dosing, the fowls, one at a time, were released to mate with a healthy hen and observed carefully for indices known as mating behaviours (wing flapping, body shakes, crowing and beak pecking) for fowl[21]. The frequencies of the mating behaviours were compared with the control fowls. The duration of observation was limited for an hour, and for accuracy in determining the frequencies, video recordings were performed for subsequent viewing. The mating experiment conducted in a closed area (dimension: 3 m × 10 m) fitted with a glass window within the Experimental Animal Laboratory of Universiti Malaysia Pahang.

2.2.5. Blood sampling

To determine changes in biochemical parameters of vital organs after 50 days of dosing, approximately 4 mL blood was carefully taken into a vacutainer by positioning the fowl to expose the wing, next identified the brachial veins (located at the ventral humoral region) followed by careful drawing of the blood using a 25 G needle connected to a 5 mL syringe. The vacutainer containing the blood was sent immediately to a diagnostic lab to measure the various biochemical parameters.

2.2.6. Microanatomy of the organs

To determine the histological changes in the organs, both the liver and testis were removed and preserved in 10 mL of 10% formalin. All tissue preparations and evaluations were performed as previously described[22]. Briefly, the tissue was cut into small blocks and stained in a xylene-ethanol mixture before being waxed in paraffin. A microtome was used to make a thin slice of approximately 4–5 microns, which was placed onto a microscope slide before being stained with haematoxylin and eosin (H & E). Consequently, a trained pathologist observed the slides under a light microscope (Nikon TS100 equipped with Eclipse software) at magnification of 400× and graded the tissues for changes in structure and function in comparison to the controls. Consequently, photos were taken using a camera linked to the microscope.

2.2.7. Statistical analysis

All data requiring statistical analysis were appropriately analysed by using IBM SPSS Statistics version 21. Analysis of variance was applied to determine the statistical significance of the data by calculating the variance in the data collected. A *P*-value less than 0.05 was considered as significant. To test the means of the population, the least significance formulae was used.

2.2.8. Ethics approval

This study was approved by the Institutional Animal Care and Use Committee of the Universiti Malaysia Pahang (ethics approval number: UMPIACUC_2018_01).

3. Results

3.1. In vitro evaluations of testosterone production

Initially, TM3 Leydig cells were used to examine the effects of *P. bullata* on testosterone production. As shown in Figure 1, *E. longifolia* (denoted as the positive control), which was the most popularly studied Tongkat Ali plant, had higher percentage of increase in testosterone secretion, at (0.80±0.02) nmol/L (350%), followed by *P. bullata* at (0.48±0.04) nmol/L (170%) compared to the untreated control cells with only (0.18±0.02) nmol/L (0%). The outcome showed almost twice as more testosterone produced by TM3 Leydig cells by *E. longifolia* compared to *P. bullata* in the *in vitro* test conducted. The results were significantly different compared to the testosterone produced by the untreated cells (*P*<0.05).

3.2. Frequency of mating behaviours in the fowl

A total of four mating behaviours were observed, and their frequencies were shown in Table 1. The mating behaviours included wing flaps (flapping with the head held high and chest outwards), body shakes (wing dropped down and held inside, followed by dance-like motion circling around the hen), crows (distinctive chicken noise to attract attention of hen) and beak pecking (using the beak to peck the back of the hen's neck). Based on five observations from video recordings of fowl mating with the hens, the total frequencies for all the behaviours were 23 counts for fowl dosed with *P. bullata* and only approximately 15 counts for the control fowl.




Figure 1. Percentage of testosterone production by TM3 Leydig cells after treatment with 50 µg/mL *Polyalthia bullata* and *Eurycoma longifolia* for 72 h compared to the untreated control cells. The increases in the *Polyalthia bullata* and *Eurycoma longifolia* groups are significantly different compared to the untreated control cells ($P < 0.05$).

3.3. Blood diagnostics

The levels of the selected blood diagnostics were given in Table 2. To date, no known normal range is available for any of the biochemical parameters for chickens, unlike for humans. The testosterone measured for the control fowl averaged (4.05±0.84) nmol/L, which more than doubled to (9.72±1.1) nmol/L for *P. bullata*-treated fowl. There was a significant difference in testosterone between

the two groups ($P=0.02$). Apart from the markedly lower detection of aspartate aminotransferase (AST), alanine aminotransferase (ALT) and bilirubin, all other parameters were within the levels of the controls and hence considered as insignificant and safe. Interestingly, the decrease in AST and ALT (biomarkers generalized for liver failure) and in bilirubin [approximately three times below the average of the controls, (50.74±2.97) µmol/L] suggested that this was not the result of acute toxicity but rather due to protective capabilities generally observed in some medicinal plants.

3.4. Histological profiles of liver and testis

The results gave the overall impressions of the *P. bullata*-treated fowl having livers with focal areas of congestion in the central veins and scattered inflammation within the hepatocytes, sinusoids and portal triads. However, only insignificant to relatively mild congestion was seen in all livers (Table 3).

A photomicroscopic section of the control liver microanatomy, shown in Figure 2a, showed no discernible evidence of congestion, a central vein that appeared normal and a portal triad geometrically well-organized throughout. The inspection of the sinusoidal capillary showed that it was similar to the natural structure found in liver cells. Figure 2b showed fowl-4 was graded as the most extreme among the four *P. bullata*-treated fowls, which had focal regions of congestion in the central vein and scattered periportal inflammation, collectively indicating that the dose given was able to lead to an initial change in the liver; however, the drug evaluated was not strong enough to cause severe necrosis or impairment.

Table 1. Mating frequency observed in fowls treated with *Polyalthia bullata*.

Behaviours	Average counts		<i>t</i> -value	<i>P</i> value
	Control	<i>Polyalthia bullata</i>		
Wing-flaps	4.0±0.4	5.8±0.8	2.887	0.01
Body shakes	4.3±0.2	6.0±1.4	2.552	0.01
Crows	3.5±0.3	5.3±0.4	2.465	0.01
Pecking	3.0±0.1	6.0±0.7	3.745	0.01
Total	14.8±0.3	23.1±0.7	3.659	0.01

Data are expressed as mean±SD; $n=4$ in each group. The control group is given capsules containing corn bran (blended chicken feed).

Table 3. Histopathological evaluations of the livers of the fowls treated with *Polyalthia bullata*.

<i>Polyalthia bullata</i>	Inflammation	Necrosis	Congestion
Fowl-1	+	-	+
Fowl-2	+	+	+
Fowl-3	+	-	+
Fowl-4	+	-	++

Note: Fowls 1-4 are dosed twice daily and at each instance with a capsule containing 10 mg of *Polyalthia bullata* as the active ingredient. The grading scale used is as follows: no structural changes (-), insignificant to mild changes (+), moderate changes (++) or severe changes (+++), compared to the histology of the control fowl.

Table 2. Biochemical parameters of the *Polyalthia bullata*-dosed and the control groups.

Biochemical parameters	Control	<i>Polyalthia bullata</i>	<i>t</i> -value	<i>P</i> value
Testosterone (nmol/L)	4.05±0.84	9.72±1.10	4.350	0.02
ALT (µ/L)	8.00±2.94	5.00±0.10	1.777	0.04
AST (µ/L)	312.67±49.80	222.00±33.50	2.119	0.03
GGT (µ/L)	22.01±3.52	21.50±3.30	1.510	0.07
Total protein (g/L)	50.76±2.98	46.25±3.42	1.601	0.06
Albumin/Globulin ratio (g/L)	0.53±0.05	0.65±0.08	1.605	0.06
Total bilirubin (umol/L)	50.74±2.97	18.25±1.78	3.250	0.02
Urea (mmol/L)	1.02±0.25	1.10±0.31	1.320	0.10
Creatinine (umol/L)	11.75±0.96	10.00±1.58	1.312	0.10
Uric acid (mmol/L)	0.13±0.02	0.14±0.02	1.319	0.10

Data are expressed as mean±SD; $n=4$ in each group. ALT: alanine aminotransferase; AST: aspartate aminotransferase; GGT: gamma glutamyltransferase.




Figure 2. Histology of normal liver (A) and fowl-4 (B) liver tissues stained with H & E under a magnification of 400×. The circled area highlights comparison between a selected region of the portal triad for the control (A) and fowl-4 (B) liver tissues. The outcome indicates no visible necrosis, mild inflammation and tolerable congestions within the tissues of the most extremely graded of the 4 fowls tested *i.e.* belonging to fowl-4.




Figure 3. Histology of normal testis (A) and a selected dosed fowl, fowl-4 (B) tissues stained in H & E under a magnification of 400×. The circled area highlights the increased activity within the seminiferous tubules. Overall, there is a progressive development of seminiferous tubules, spermatogonia, primary and secondary spermatocytes, spermatids and spermatozoa of fowl-4 compared to those belonging to the control.

The normal testicular tissue (Figure 3a) examined showed abundant seminiferous tubules with dimensions estimated to be approximately 150 microns but not exceeding 200 microns, while the spermatogonia, primary and secondary spermatocytes, spermatids and spermatozoa were structurally uniform and normal. Any Leydig cells visible were without inflammation or fibrosis. The testicular microanatomy of the fowl dosed with *P. bullata* was also examined, with that of fowl-4 was again chosen as the representative (Figure 3b). Some degree of increased seminiferous tubule size was evident, estimated to be up to 250–300 microns, which was apparently slightly larger than the control shown. The tubules were concluded to be apparently increased cellularly as well as in numbers. The same results were observed for the spermatogonia, primary and secondary spermatocytes, spermatids and spermatozoa, all of which indicated a progressive development in comparison to the controls. Throughout the inspection, no signs of inflammation or fibrosis were observed. Furthermore, the spermatogonia were intensely dense in the core of the lumen with long tails and a close grouping. All of these results indicated that *P. bullata* was capable of encouraging the positive development of seminiferous tubules and germ cell division followed by maturation better than the control.

4. Discussion

P. bullata is capable of releasing testosterone in TM3 Leydig-treated cells, although to levels not as high as the more popularly studied Tongkat Ali plant of *E. longifolia*. No similar study on this cell line has been reported for *P. bullata*. Mohamed *et al* reported

that *E. longifolia* possesses a greater ability to promote serum testosterone increase in rats than *P. bullata*[23].

Additionally, *P. bullata* provided a significant libido improvement based on sexual mating behaviours observed in the treated fowl. The fowl also showed an increased testosterone production by approximately 138%. The microanatomy of specific testicular tissue regions further supported the capability of *P. bullata* to stimulate the male sexual reproductive organ of the fowl. There are no available results assessing the ability of *P. bullata* to improve sexual function in chickens. However, it is known that the inclusion of testosterone alone by direct injection into fowl causes an increase in sexual behaviours such as crows and rooster dances upon facing hens[24].

The current work on *P. bullata* indicates that it has the potential to elevate testosterone and mating behaviours. In fowl, these effects were achieved only after a total of 50 days of dosing with a capsule containing 10 mg of active ingredient and required dosing twice a day. Such an activity can be considered as late-onset comparison to drugs such as Viagra or Cialis, which are effective within hours after consumption. Herbal aphrodisiacs in fact have a late-onset activity, as shown previously by similar work in fowl given capsules containing 12 mg of *E. longifolia* and *S. tuberosa*, both capable of increasing testosterone levels after 30 days when dosed individually[10]. The indigenous people of Kampung Orang Asli of Bukit Cermin revealed that some of their customers purchasing and using *P. bullata* for the very first time experienced none of the aphrodisiac effects of this plant in the first month of consumption. Due to this explanation given by the indigenous people as well as the initial trial of dosing the fowl twice a day with a 10 mg capsule for 30 days provided negative results (data not included), hence prompting another trial

of up to 50 days. Importantly, the results of this study provided evidence of the testosterone-boosting capabilities of *P. bullata* in fowl, similarly shown in only one other study, where this plant was administered to rats[23]. In contrast, *E. longifolia*, which shares the same name of Tongkat Ali, has been proven in numerous studies[25]. Some examples include the study of male rats and wild pigs fed *E. longifolia* that showed an increase in sexual desires and other important sexual parameters, establishing the usefulness of this plant in its aphrodisiac capabilities[26,27].

Personal communications with the families of indigenous people also highlighted that their returning customers had never complained about any adverse effects after consuming *P. bullata*. In this study of administering *P. bullata* for a duration of 50 days, none of the biochemical markers assessed were alarmingly elevated, and even in the worst graded fowl, the histology examination by the pathologist did not report severe findings, suggesting that overall, for short-duration or acute investigations, the plant is considerably safe. A study detected high levels of heavy metals in 25% of a total of 100 herbal products with *P. bullata* as the active ingredient[11]. This sort of metal ion accumulation is due to the root leaching water containing industrial contaminated soil[28]. The samples of *P. bullata* obtained by the indigenous people within virgin forests that were administered to the fowl are unlikely to be polluted by heavy metals; however, herbal use has been demonstrated to be toxic towards vital organs such as the liver, kidney or heart. Hence, *P. bullata* still requires chronic toxicity studies before concluding that it is safe. Popular Indian medicinal plants such as *Eclipta alba* and *Phyllanthus niruri* have apparently been shown to have adverse effects on the liver at high doses[29,30].

ALT and AST, if elevated to more than 5 times the highest values of their normal ranges, are considered to indicate the onset of likely liver failure according to the Association for the Study of Liver Diseases[31]. To date, unlike humans, there is no range for these biomarkers in chickens; hence, they can only be compared with the control fowl. In this study, the biochemical markers of the liver, namely, ALT and AST, were not elevated; instead, they were lowered, as was bilirubin. In chickens, abnormally elevated bilirubin levels, accompanied by obstructive jaundice, are indicators of active liver disease[32]. However, not only were the levels of ALT, AST and bilirubin lower than those of the controls, but histology examinations were also supportive of normal structural architectures and functions of the hepatocytes. Plants have already been reported to be able to lower bilirubin that was initially elevated dangerously high due to carbon tetrachloride exposure in liver cells[16,33,34]. Although none of these liver protective studies have ever been reported for *P. bullata*, it is likely that it may possess such potential. Importantly, *P. bullata* was found to be safe in comparison to *E. longifolia*, which was demonstrated in a similar study to induce an elevation in ALT that was noticeable after 30 days of dosing with 12 mg *E. longifolia* per capsule twice daily in fowl[10].

This study limited by the ability to use a larger number of animals due to 3Rs (Replacement, Reduction and Refinement) requirement

imposed by Institutional Animal Care and Use Committee. The researchers understood and accepted the requirement as it is for the welfare of the animal in study. This study of evaluating for 50 days was only an acute investigation on the efficacy and safety of *P. bullata*. A longer duration of at least 6 months can potentially provide the chronic outcomes.

In conclusion, *P. bullata* showed an increase in efficacy indicators in both *in vitro* (Leydig cells) and *in vivo* investigations performed in fowl (elevated frequencies of selected sexual mating indicators, boosted testosterone concentrations in blood, and evidence of a positive active testicular microanatomy). The onset of aphrodisiac activities nevertheless required up to 50 days of dosing in the fowl.

Conflict of interest statement

The authors declare that there is no conflict of interest.

Funding

This work was funded by grants awarded by Malaysian Technological University Network (MTUN) with grant number of UIC191201 as well as grants awarded by Universiti Malaysia Pahang with grant numbers of PDU203209 (product development grant) and PGRS1903203 (postgraduate grant).

Authors' contributions

The main author, Assoc. Prof Dr Jaya Vejjayan, was the project leader, involved in conceptualizing, designing and executing the project, as well as in writing the manuscript from an initial rough draft stage to its current status and in editing and submitting the manuscript. The second author, Yasmin Amira Che Yahya, contributed to performing the bench work to obtain all raw data and ensure statistical validity. She also contributed to collating the results into the initial rough draft of the manuscript for the purpose of refining by other authors. Professor Dr Srikumar Chakravarthy, with expertise as a pathologist, contributed to preparing the histology tissues and examining and grading them. Next, Professor Dr Halijah Ibrahim was involved in the initial conceptualization of the research into the aphrodisiac plants, editing the manuscript draft and providing her expertise in data analysis. Last, Aida Yun, as an indigenous person, was of the utmost assistance in acquiring precise plants from the wild for the numerous samples required for the completion of the study.

References

[1] Chauhan NS, Sharma V, Dixit VK, Thakur M. A review on plants used

- for improvement of sexual performance and virility. *BioMed Res Int* 2014; **2014**: 868062.
- [2] Gupta G, Rana AC. *Withania somnifera* (Aswagandha). A review. *Phcog Rev* 2007; **1**(1): 129-136.
- [3] Tocharus C, Jeenapongsa R, Teakthong T, Smitasiri, Y. Effects of long-term treatment of *Butea superba* on sperm mobility and concentration. *Naresuan Univ J* 2005; **3**: 11-17.
- [4] Qureshi S, Shah AH, Tariq M, Ageel AM. Studies on herbal aphrodisiac used in Arab system of medicine. *Am J Chin Med* 1989; **17**: 57-63.
- [5] Cho WK, Weeratunga P, Lee BH, Park JS, Kim CJ, Ma JY, et al. *Epimedium koreanum* Nakai displays broad spectrum of antiviral activity *in vitro* and *in vivo* by inducing cellular antiviral state. *Viruses* 2015; **7**(1): 352-377.
- [6] Abdullah N, Hosseinpour Chermahini DS, Chua LS, Sarmidi M. *Labisia pumila*: A review on its traditional, phytochemical and biological uses. *World Appl Sci J* 2013; **27**(10): 1297-1306.
- [7] Chee BJ. Men's health: Tongkat Ali Hitam (*Polyalthia bullata*). In: Vimala S. (ed.) *Malaysian herbal heritage*. Kepong, Selangor: Forest Research Institute Malaysia; 2013.
- [8] Vejjayan J, Mohamed AA, Zulkifli AA, Yahya YAC, Munir N, Yusoff MM. Marker to authenticate *Eurycoma longifolia* (Tongkat Ali) containing aphrodisiac herbal products. *Curr Sci* 2018; **115**(5): 886-894.
- [9] Tnah L, Lee S, Tan A, Lee C, Ng K, Ng C, et al. DNA barcode database of common herbal plants in the tropics: A resource for herbal product authentication. *Food Control* 2019; **95**: 318-326.
- [10] Vejjayan J, Che Yahya YA, Chakravarthy S, Bathmanathan R, Ibrahim H, Yun A. Tongkat Ali plants of *Eurycoma longifolia* and *Stema tuberosa* stimulate sexual arousal in domestic cocks. *Malays J Sci* 2020; **39**(1): 1-14.
- [11] Ang HH, Lee KL. Contamination of mercury in Tongkat Ali hitam herbal preparation. *Food Chem Toxicol* 2006; **44**(8): 1245-1250.
- [12] Effendy NM, Mohamed N, Muhammad N, Mohamad IN, Shuid AN. *Eurycoma longifolia*: Medicinal plant in the prevention and treatment of male osteoporosis due to androgen deficiency. *Evidence-Based Complement Altern Med* 2012; **2012**: 125761.
- [13] Mohamed AN, Vejjayan J, Yusoff MM. Review on *Eurycoma longifolia* pharmacological and phytochemical properties. *J Appl Sci* 2015; **15**(6): 831-844.
- [14] Ang HH, Cheang HS, Yusof APM. Effects of *Eurycoma longifolia* Jack (Tongkat Ali) on the initiation of sexual performance of inexperienced castrated male rats. *Exp Anim* 2000; **49**(1): 35-38.
- [15] Bhat R, Karim AA. Tongkat Ali (*Eurycoma longifolia* Jack): A review on its ethnobotany and pharmacological importance. *Fitoterapia* 2010; **81**(7): 669-679.
- [16] Rehman S, Choe K, Yoo H. Review on a traditional herbal medicine, *Eurycoma longifolia* Jack (Tongkat Ali): Its traditional uses, chemistry, evidence-based pharmacology, and toxicology. *Molecules* 2016; **21**(3): 331.
- [17] Scarano WR, Messias AG, Oliva SU, Klinefelter GR, Kempinas WG. Sexual behaviour, sperm quantity and quality after short-term streptozotocin-induced hyperglycaemia in rats. *Int J Androl* 2006; **29**(4): 482-488.
- [18] Tambi MIBM, Imran MK, Henkel RR. Standardised water-soluble extract of *Eurycoma longifolia*, Tongkat ali, as testosterone booster for managing men with late-onset hypogonadism. *Andrologia* 2012; **44**(Suppl 1): 226-230.
- [19] Talbott SM, Talbott JA, George A, Pugh M. Effect of Tongkat Ali on stress hormones and psychological mood state in moderately stressed subjects. *J Int Soc Sports Nutr* 2013; **10**(1): 28.
- [20] Erasmus N. *Investigations on the in vitro effects of aqueous Eurycoma longifolia Jack extract on male reproductive functions*. MSc thesis. University of the Western Cape, Bellville, South Africa; 2013.
- [21] Leonard ML, Zanette L. Female mate choice and male behaviour in domestic fowl. *Anim Behav* 1998; **56**(5): 1099-1105.
- [22] Haleagrahara N, Jackie T, Chakravarthy S, Rao M, Kulur A. Protective effect of *Eilingeria elatior* (torch ginger) extract on lead acetate-induced hepatotoxicity in rats. *J Toxicol Sci* 2010; **35**(5): 663-671.
- [23] Mohamad M, Mohsin HF, Singh GKS. A study on the effects of *Eurycoma longifolia* and *Polyalthia bullata* on reproductive organs and androgens of male Sprague-Dawley rats. *J Engineer Appl Sci* 2017; **12**(5): 6994-6999.
- [24] Davis DE, Domm LV. The sexual behavior of hormonally treated domestic fowl. *Proc Soc Exp Biol Med* 1941; **48**(3): 667-669.
- [25] Hamdi AHA, Ming LC, Kassab YW. A review on safety and efficacy of products containing *longifolia*. *Arch Pharm Pract* 2016; **7**(5): 39.
- [26] Ho CC, Tan HM. Rise of herbal and traditional medicine in erectile dysfunction management. *Curr Urol Rep* 2011; **12**(6): 470-478.
- [27] Frydrychova S, Opletal L, Macáková K, Lustyková A, Rozkot M, Lipensk J. Effects of herbal preparation on libido and semen quality in boars. *Reprod Domest Anim* 2011; **46**(4): 573-578.
- [28] Yang B, Gao Y, Zhang C, Zheng X, Li B. Mercury accumulation and transformation of main leaf vegetable crops in Cambosol and Ferrosol soil in China. *Environ Sci Pollut Res* 2019; **27**: 391-398.
- [29] Singh T, Sinha N, Singh A. Biochemical and histopathological effects on liver due to acute oral toxicity of aqueous leaf extract of *Ecliptaalba* on female Swiss albino mice. *Indian J Pharmacol* 2013; **45**(1): 61-65.
- [30] Singh T, Ruchi Kumar R, Singh A. Evaluation of biochemical and histological effects on liver of Swiss albino mice due to acute oral toxicity of aqueous leaf extract of *Phyllanthus niruri*. *Int J Pharmacogn Phytochem Res* 2016; **8**(1): 85-90.
- [31] Kim WR, Flamm SL, Di Bisceglie AM, Bodenheimer Jr HC. Serum activity of alanine aminotransferase (ALT) as an indicator of health and disease. *Hepatology* 2008; **47**(4): 1363-1370.
- [32] Sabater M, Forbes N. Avian haematology and biochemistry 2. *Biochemistry. Practice* 2015; **37**(3): 139-142.
- [33] Mohamed MA, Eldin IM, Mohammed AE, Hassan HM. Effects of *Lawsonia inermis* L. (Henna) leaves' methanolic extract on carbon tetrachloride-induced hepatotoxicity in rats. *J Intercult Ethnopharmacol* 2015; **5**(1): 22-26.
- [34] Parvez MK, Al-Dosari MS, Arbab AH, Alam P, Alsaied MS, Khan AA. Hepatoprotective effect of *Solanum surattense* leaf extract chemical-induced oxidative and apoptotic injury in rats. *BMC Complement Altern Med* 2019; **19**(1): 154.