

## RESEARCH ARTICLE

## High Keratin Secretion of T47D Cell under Hypoxic Condition

Annisa<sup>1</sup>, Elizabeth Henny Herningtyas<sup>2</sup>, Dewajani Purnomosari<sup>3,\*</sup>

<sup>1</sup>Post Graduate Program of Biomedical Sciences Faculty of Medicine, Public Health and Nursing, Universitas Gadjah Mada, Jl. Farmako, Sekip Utara, Yogyakarta 55281, Indonesia

<sup>2</sup>Department of Clinical Pathology and Laboratory Medicine, Faculty of Medicine, Public Health and Nursing, Universitas Gadjah Mada, Jl. Farmako, Sekip Utara, Yogyakarta 55281, Indonesia

<sup>3</sup>Department of Histology and Cell Biology, Faculty of Medicine, Public Health and Nursing, Universitas Gadjah Mada, Jl. Farmako, Sekip Utara, Yogyakarta 55281, Indonesia

\*Corresponding author. E-mail: d.purnomosari@ugm.ac.id

Received date: Jul 13, 2022; Revised date: Aug 5, 2022; Accepted date: Aug 16, 2022

### Abstract

**BACKGROUND:** Hypoxia adaptation in cancer cells is mediated by hypoxia inducible factor (HIF)-1 that induces the expression of various proteins. Up to date, the analysis of T47D breast cancer cells-secreted protein is still limited. The aim of this study was to compare the protein profile secreted by T47D breast cancer cells under hypoxic and to normoxic conditions and identify the proteins as candidate for hypoxia marker proteins in T47D breast cancer cells.

**METHODS:** T47D breast cancer cells were cultured under standard conditions. Cells were subcultured in normoxic and hypoxic conditions. The normoxic group was incubated with 20% oxygen and the hypoxic groups were incubated in a hypoxic chamber with 0.5% and 5% oxygen for 6, 24, and 48 hours in serum free medium. Proteins in the culture media were isolated and precipitated. Protein concentrations

released by the cells were then measured by bicinchoninic assay (BCA). Protein bands were visualized by the sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) method. The bands that had differences between the hypoxic and normoxic groups were further analyzed by liquid chromatography-mass spectrophotometer (LC-MS).

**RESULTS:** Hypoxic groups showed higher secretory protein than normoxic group. Protein bands were found in the 0.5% hypoxic group with a size of 50-75 kDa. Secretory proteins identified by LC-MS were keratin 1, 2, 9, and 10.

**CONCLUSION:** The T47D cell line under 0.5% hypoxic treatment showed higher secretory proteins that identified as keratin 1, 2, 9, and 10.

**KEYWORDS:** hypoxia, secretory protein, liquid chromatography-mass spectrophotometer

*Indones Biomed J. 2022; 14(4): 376-81*

### Introduction

In solid tumors, excess proliferation increases the distance of tissue oxygen diffusion from the vessels leading to insufficient angiogenesis in the growing tumor lesion. (1) Areas of hypoxia in solid tumors can cause various changes in tumor cell regulation as an adaptation to hypoxic conditions. Hypoxia in tumor cells is known to play an

important role in the development, resistance to anti-tumor agents and metastasis.(2,3)

Cellular responses to hypoxic conditions are mediated by the transcription factor, hypoxia inducible factor (HIF)-1, which can influence the expression of various proteins. Proteins can be released by tumor cells into the extracellular matrix to support the tumor microenvironment. Protein release under hypoxic conditions can occur directly through the activation of HIF, as well as the activation of cellular

signaling pathways. Hypoxia in tumor cells is known to increase the release of proteins and other effectors that can trigger changes in cell behavior. These proteins can cause tumor cells to develop with unrestricted growth characteristics and can invade surrounding tissues.(4,5) Increased HIF expression as a result of hypoxia is known to occur in 56-57% of breast cancers.(6)

Identification of hypoxia biomarker in human umbilical cord mesenchymal stem cells (hUC-MSCs) revealed that secretion of Vascular Endothelial Growth Factor (VEGF) reached the highest level at 96 hours hypoxic condition.(7,8) Identification of candidate biomarker secreted protein released by cancer cells using mass spectrophotometry already used in human endometrial cancer, cervical cancer cells (9,10), lung cancer (11), head and neck cancer (12), colorectal cancer (13,14), and many more. Meanwhile, secretome analysis of previous *in vitro* study in the hypoxic cancer secretome from MDA-MB 231 breast cancer cell analysis revealed that high expression of lysyl oxidase in breast cancer may contribute to osteolytic lesion formation.(6,15)

Among various types of breast cancer, Luminal A breast cancer subtype is the most prevalent breast cancer. The nowadays available cell lines that represent luminal A type breast cancer are T47D and MCF7 cells. Secretory protein analysis in MCF7 breast cancer cell line under hypoxic condition showed stem cell factor (SCF) and vascular endothelial growth factor (VEGF) as the main secretory protein.(16) However, the secretory protein analysis of T47D breast cancer cell line has not been conducted before, thus different profile of secretory proteins in hypoxic T47D breast cancer cell line might be suggested. Studies comparing hypoxic secretory protein profiles in breast cancer with modeling of the T47D cell line versus normoxia are not available yet. The aim of this study was to compare the secretory proteins produced by the T47D breast cancer cell line cultured under hypoxic and normoxic conditions.

## Methods

### T47D Cell Culture

T47D breast cancer cell line was obtained from Elabscience (Cat. No. EP-CL-0228, Texas, USA) with certificate of analysis for authentication. T47D cells were grown in 100 mm dishes using culture media (RPMI 1640 with 10% FBS, 0.2 U/mL insulin, 1% Penicillin-Streptomycin, Amphotericin; all materials obtained from Capricorn

Scientific, Ebsdorfergrund, Germany). T47D subculture was carried out when it reached 70-90% confluent with a planting density of  $4 \times 10^4$  cells/cm<sup>2</sup>. Cells were harvested and  $2.2 \times 10^6$  cells were grown in 100 mm petri dishes (Iwaki, Woodland, Singapore). The culture cell medium was replaced with new media every 48 hours and incubated in an incubator at 37°C with 5% CO<sub>2</sub>.

### Hypoxic Treatment

Cells were harvested and  $2.2 \times 10^6$  cells then planted and grown in 100 mm plates until confluence reached 60-70%. When confluency reached 70%, cells were washed with PBS three times then the medium was replaced with serum free medium. The normoxic group was incubated in 37°C and 20% oxygen. The hypoxic treatment group was incubated in a hypoxic chamber (Stem Cell®, Vancouver, Canada) which flowed with a mixed gas with an oxygen concentration of 0.5% and 5% for 4 minutes at a flow of 20 L/minute. The medium from T47D cell cultured was then collected and transferred to the tube according to the hypoxic treatment time 6 hours, 24 hours and 48 hours for further analysis. All treatment groups were conducted in triplicate.

### Protein Isolation and Precipitation

The culture media were centrifuged at 800 rpm for 10 minutes and filtered with a 0.2 µm syringe filter to eliminate cells and debris. Media were added with 10% trichloroacetic acid (TCA) (Merck, Kenilworth, NJ, USA). The medium then precipitated on ice for 2 hours followed by centrifugation for 30 minutes at 14000 rpm. Washing was carried out with cold acetone 2 times. The washed precipitate protein was resuspended with 50 Mm ammonium bicarbonate. Protein concentration was measured by Bicinchoninic Assay/BCA kit (Thermo Fisher Scientific, Rockford, IL, USA).

### BCA Assay

Measurement of protein precipitate concentration was conducted by the BCA kit (Thermo Fisher Scientific). After reagent preparation by mixing reagent A and B, a total of 25 µL of standard protein and samples were put into the 96-well microplate (Iwaki). Working reagent (200 µL) was added to each well and incubated in an incubator at 37°C for 30 minutes. Samples were read with a spectrophotometer at a wavelength of 570 nm.

### Protein Separation with Sodium Dodecyl Sulfate Polyacrylamide Gel Electrophoresis (SDS-PAGE)

We used TGX FastCast Acrylamide Kit 10% (BioRad, California, USA) for SDS-PAGE methods. The step


began with making running gel/resolving gel and stacking gel according to the product manual. A total of 20  $\mu\text{L}$  of sample protein was transferred into the well and electrophoresis was performed at 100 v for 2 hours. The gel was then stained with 1% Coomassie Brilliant Blue R250 on a shaker until the bands were visible. The gel bands with different expressions were further diced using scalpel and transferred to a micro-tube containing 50 mM Ammonium Bicarbonate/50% acetonitrile then vortexed for 10 minutes.

### Liquid Chromatography-Mass Spectrophotometer (LC-MS) Preparation

Protein samples were incubated for 60 minutes in 10 mM Dithiothreitol (DTT) for protein unfolding. After reduction, protein samples were alkylated by adding 54 mM Iodoacetamide (IAM) and incubated in the dark at room temperature for 30 minutes. Supernatant was removed and gel cubes were vortexed in 400  $\mu\text{L}$  50 mM  $\text{NH}_4\text{HCO}_3$  for 10 minutes for two times. Supernatant then discharged and alkylated protein samples were digested using trypsin in 50 Mm ammonium bicarbonate with incubation at 37°C overnight. Digested protein then dried using the freeze drying and redissolved in 0.1% formic acid.

### Protein Analysis Using LC-MS

The redissolved digested samples were placed in the automatic sample injector on the LC-MS device. The tool used is the NanoLC Ultimate 3000 Series System Tandem Q Exactive Orbitrap HRMS Thermo Scientific™. The chromatographic system used was reverse phase. The stationary phase used was C18 (PepMap RSLC). The mobile phase used was water with 0.1% formic acid and acetonitrile, 0.1% formic acid. Separation was carried out using a mobile phase gradient: gradient B 2% for 3 minutes, 2%-35% B for 27 minutes, 35-90% B for 15 minutes, 90% B for 15 minutes, 2%B for 15 minutes. LC-MS data was in the form of a chromatogram that showed the ratio of mass to electric charge (m/z) for each compound. The data were analyzed by Proteome Discoverer 2.2 (Thermo fisher scientific, Rockford, USA) to identify the protein present in the sample (Figure 1).



**Figure 1. Flowchart of mass spectrophotometric liquid chromatography examination.** The sample was injected and passed through an ion source for ionization. The peptides were separated based on the charge of the ions in the analyzer. The mass analyzer measured the Mass-to-charge ratio (m/z). The detector captured the ions and measures the intensity of each ion. Proteins were identified through bioinformatics analysis.

## Results

Secretory protein concentration in T47D cell culture medium that had been precipitated with 10% (v/v) TCA was measured by BCA. The total protein concentration in the culture media of the hypoxic treatment groups were higher than the normoxic group (Table 1), and while compared to the normoxic group, there were increments of protein ratio from the hypoxic groups at the range of 1.73-22.74 folds. The increments in protein concentration were in accordance with the increment of the hypoxic concentration and the duration of treatment. The lowest protein concentration was found in the normoxic group while the highest secretory protein concentration was found in the 48 hours 0.5% hypoxic group. There was an increased concentration of secretory proteins in the 5% hypoxic group compared to the normoxic group, but the increase was not as much as the 0.5% hypoxic group.

The proteomic analysis method in this study combined SDS-PAGE electrophoresis and mass spectrophotometry. The SDS-PAGE method can identify proteins based on their molecular weight and isoelectric strength. The SDS-PAGE gel showed a band in the 0.5% hypoxic treatment group with a molecular weight between 50 kDa-75 kDa which did not appear in the 5% hypoxic or normoxic groups (Figure 2). Protein identification was carried out using the LC-MS method on the eluted SDS-PAGE band. The proteins in the SDS-PAGE band were identified as keratin 1, keratin 2, keratin 9, keratin 10 and cDNA FLJ54371 proteins/serum albumin (Table 2).

## Discussion

The increased secretory protein concentration obtained from culture medium under hypoxic conditions may be related to the expression of HIF-1 $\alpha$ .(17) Previous study showed that HIF-1 $\alpha$  expression is known highest with 0.5% oxygen and significantly decreases above 2% to 6% oxygen.(18) HIF-1 $\alpha$  expression in various types of breast cancer cells is known to reach a peak between 4 hours to 8

**Table 1. Protein concentration from T47D cell culture medium with 0.5%, 5% hypoxic, and normoxic treatment.**

Group	Protein Concentration ( $\mu\text{g/mL}$ )			Protein Concentration Average ( $\mu\text{g/mL}$ )	Protein Ratio (Hypoxic/normoxic)
	1	2	3		
0.5% hypoxic, 6 hours	130.63	92.51	83.43	102.19	1.95
0.5% hypoxic, 24 hours	119.49	111.83	128.11	119.81	2.29
0.5% hypoxic, 48 hours	178.97	126.33	124.67	143.33	2.74
5% hypoxic, 6 hours	132.35	128.91	79.31	113.52	2.17
5% hypoxic, 24 hours	130.63	72.71	69.48	90.94	1.73
5% hypoxic, 48 hours	87.53	170.03	82.60	113.39	2.16
Normoxic	61.33	35.00	60.55	52.30	1.00

hours of hypoxia and begins to decrease after 12 hours of treatment.(19)


Protein release by the tumor cells is carried out mainly through extracellular vesicles. Hypoxia is known to increase biogenesis and release of secretory vesicles. Several mechanisms that can increase the release of these secretory vesicles are through HIF, Rab-GTPase, nuclear factor kappa-B (NF- $\kappa$ B) and tetraspanin signaling.(20) These extracellular vesicles have an important role in intercellular communication through carried biological molecules, including proteins. In tumor cells, these extracellular vesicles are mainly released into the microenvironment around the tumor (tumor microenvironment) and play a role in angiogenesis, invasion and metastasis.

Some factors that might cause the disappearance of the band in the 5% hypoxic and the normoxic group are the small sample concentration or whether the protein has been degraded. Type of gel dye is also known to affect the quality of the band on the gel. The gel staining used in this study was coomassie blue R250, which is known to have a sensitivity about 3-10 ng. Several stainings are known to have a higher sensitivity than Coomassie Blue staining such as silver stain (0.25-5 ng).(21)

Keratin (formerly known as cytokeratin), is an intermediate filament present in various epithelial cells. Keratin in humans is encoded by 54 functional genes, and is expressed specifically depending on the cell type. Keratin plays an important role in maintaining cellular integrity and protecting cells from injury. There are 2 types of keratin, namely type I (K9-K19), keratin type II (K1-K8). Keratin is known to be very dynamic to cellular stress and environmental changes.(22)

Keratin was recognized as a fragmented secretory product due to the degradation of dead cells. Several types of colorectal cancer cells (HT29, HCT116, Caco-2) and breast cancer cells (MCF-7, SKBR3, MDA MB-231) showed that cancer cells released cytokeratin 19 (CK-19) in a full length.(23) The cytokeratin released by breast cancer cells is thought to have a role in metastasis and tumor growth through an unexplained mechanism.

Proteomic analysis of keratin as a secretory product of cancer cells is still limited. An expression of keratin 9 is found in a proteomic analysis of breast cancer cell exosomes.(24) The expression of keratin 9 and several other cytoskeletal proteins such as tropomyosin, and transgelin were also found in a proteomic characterization study of



**Figure 2. Protein bands visualization on SDS-PAGE gel. A:** Normoxic group; **B:** 0.5% hypoxic group; **C:** 5% hypoxic group. There was a band between 50-75 kDa in the 0.5% hypoxic group, but not in the 5% hypoxic and normoxic groups.

**Table 2. Protein identified from SDS-PAGE band of T47D cell culture medium under 0.5% hypoxic treatment for 24 hours and 48 hours.**

Accession Number	Protein Identity	Protein Coverage (%)	Peptide Number	Molecule Weight (kDa)
0.5% hypoxic group, 24 hours				
H6VRG1	Keratin 1	11	6	66.1
P35527	Keratin 9	4	2	62.0
B4DPP6	Serum Albumin	2	1	70.3
0.5% hypoxic group, 48 hours				
P35527	Keratin 9	9	3	62.0
H6VRG1	Keratin 1	7	3	66.1
P35908	Keratin 2	4	2	65.4
P13645	Keratin 10	3	1	58.8
B4DPP6	Serum Albumin	2	1	70.3

breast cancer exosomes associated with metastatic ability. These proteins are known to play a role in signaling and regulation of cellular structures.(25)

Effect of hypoxia on keratin release research is still very limited and has not been fully elucidated. Hypoxia through the HIF transcription factor is known to increase the transcription of various genes. Hypoxia can cause increases in the expression of several keratin coding genes including KRT14, KRT16, KRT18, KRT19 and KRT20.(26,27) There is no data yet regarding KRT1, KRT2, KRT9, KRT10.

The mechanism of keratin release under hypoxic conditions in tumor cells has not been fully elucidated. A study on the effect of hypoxia on alveolar cells revealed that hypoxia is known to induce cellular stress by increasing reactive oxygen (ROS), which can cause reorganization of keratin. Through the activation of mitogen-activated protein kinase (MAPK) p38, keratin 8 will be phosphorylated resulting in keratin disassembly and degradation. Disassembly and degradation of keratin will reduce cell integrity and increase the ability to migrate.(28) The mechanism of keratin release under hypoxic conditions of tumor cells may have the same mechanism and further research needs to be done.

Evidence about expression of keratin as a secretory product of cancer cells under hypoxic conditions is limited. While, keratin is often referred to as a contaminant in proteomic examination. Contamination of keratin in this examination can come from skin, hair or air during sample processing. Contamination is difficult to avoid even though it is carried out in a laboratory with a high standard protocol. (29) Possible sources of keratin as a contaminant in this study can occur and have not been ascertained.

Comparison of cells in the hypoxic treatment group with the normoxia group was carried out to assess

the protein released by breast cancer cells is the protein released due to hypoxia treatment. Several materials in this experiment which are not mass spectrophotometry/MS-grade become the weakness of this study and can increase the risk of contamination. Additional examination can be done by specific examinations such as immunoeptithelial spots combined with keratin expression in culture medium. Immunoeptithelial spots can be performed to observe the expression of keratin in cells. Expression of keratin in cells could be associated with the release of proteins identified through LC-MS. In this study, immunoeptithelial spot examination could not be performed because T47D cells were used for other research. The role of hypoxia on keratin expression, especially keratin 1, 2, 9 and 10 needs to be studied further.

## Conclusion

The T47D cell line under 0.5% hypoxic treatment showed higher secretory proteins that identified as keratin 1, 2, 9, and 10. The concentration of secretory protein released by the T47D breast cancer cells in the hypoxic group was higher than the normoxic group. Expression of keratin 1, 2, 9, and 10 proteins were obtained on T47D culture medium with 0.5% hypoxia treatment.

## Acknowledgments

We are grateful to Histology Department staff Dewi Kartikawati Paramita, PhD for handling T47D until cell stock achieved and the technical assistant of all laboratory staff Laboratorium Riset Terpadu, Dr. Sardjito General

Hospital and in Histology and Cell Biology, Universitas Gadjah Mada, Yogyakarta, especially Mrs. Sumartiningsih.

## Authors Contribution

DP contributed in planning and supervised the study. EH supervised experiment method. AN working in the laboratory experiment and drafting the manuscript. DP, EH reviewed and revised the manuscript. All authors discussed the results and commented on the manuscript.

## References

- Barrak NH, Khajah MA, Luqmani YA. Hypoxic environment may enhance migration/penetration of endocrine resistant MCF7- derived breast cancer cells through monolayers of other non-invasive cancer cells in vitro. *Sci Rep.* 2020; 10(1): 1127. doi: 10.1038/s41598-020-58055-x.
- Gilkes D, Semenza G. Role of hypoxia-inducible factors in breast cancer metastasis. *Futur Oncol.* 2015; 9(11): 1623-36.
- Rahman MN, Wijaya CR, Novalentina M. Survivin clinical features in cervical cancer. *Mol Cell Biomed Sci.* 2017; 1(1): 6-16.
- Caroline W, Sven P, Bexell, Daniel. Therapeutic targeting of hypoxia and hypoxia-inducible factors in cancer. *Pharmacol Ther.* 2016; 164: 152-69.
- Iwona B, Latosinska A, Metzger J, Marx D, Vlahou A, Mischak H, *et al.* Proteomics biomarkers for solid tumors: Current status and future prospects. *Mass Spec Rev.* 2017; 38(1): 49-78.
- Kozakiewicz AMB, Budzik MP, Przybylsky J. Hypoxia in breast cancer. *Pol J Pathol.* 2015; 66(4): 337-46.
- Sidharta VM, Herningtyas EH, Lagonda CA, Fauza D, Kusnadi Y, Susilowati R, *et al.* High VEGF level is produced by human umbilical cord mesenchymal stem cells (hUC-MSCs) in amino acid-rich medium and under hypoxia condition. *Indones Biomed J.* 2018; 10(3): 222-30.
- Meiliana A, Dewi NM, Wijaya A. Cancer stem cell hypothesis: Implication for cancer prevention and treatment. *Indones Biomed J.* 2016; 8(1): 21-36.
- Li H, DeSouza LV, Ghanny S, Li W, Romaschin AD, Colgan TJ, *et al.* Identification of candidate biomarker proteins released by human endometrial and cervical cancer cells using two-dimensional liquid chromatography/tandem mass spectrometry. *J Proteome Res.* 2007; 6(7): 2615-22.
- Lilis L, Siregar NC, Andrijono A. The association of parametrial invasion with the expression of vascular endothelial growth factor-C and other factors in squamous cell cervical carcinoma stage IB and IIA. *Indonesian Biomedical Journal.* 2019; 11(3): 279-85.
- Huang LJ, Chen SX, Luo WJ, Jiang HH, Zhang PF, Yi H. Proteomic analysis of secreted proteins of non-small cell lung cancer. *Ai Zheng.* 2006; 25(11): 1361-7.
- Ralhan R, Desouza LV, Matta A, Tripathi SC, Ghanny S, Datta Gupta S, *et al.* Discovery and verification of head-and-neck cancer biomarkers by differential protein expression analysis using iTRAQ labeling, multidimensional liquid chromatography, and tandem mass spectrometry. *Mol Cell Proteomics.* 2008; 7(6): 1162-73.
- Shin J, Kim HJ, Kim G, Song M, Woo SJ, Lee ST, *et al.* Discovery of melanotransferrin as a serological marker of colorectal cancer by secretome analysis and quantitative proteomics. *J Proteome Res.* 2014; 13(11): 4919-31.
- Wanandi SI, Lestari DR, Hilbertina N, Siregar NC, Jusman SW, Abdullah M. Secretomes of primary cancer-associated fibroblasts upregulate the expression of stemness markers in HT-29 human colorectal carcinoma cells. *Indones Biomed J.* 2020; 12(4): 333-9.
- Widowati W, Jasaputra DK, Sumitro SB, Widodo MA, Afifah E, Rizal R, *et al.* Direct and indirect effect of TNF $\alpha$  and IFN $\gamma$  toward apoptosis in breast cancer cells. *Mol Cell Biomed Sci.* 2018; 2(2): 60-9.
- Han ZB, Ren H, Zhao H, Chi Y, Chen K, Zhou B, *et al.* Hypoxia-inducible factor (HIF)-1 $\alpha$  directly enhances the transcriptional activity of stem cell factor (SCF) in response to hypoxia and epidermal growth factor (EGF). *Carcinogenesis.* 2008; 29(10): 1853-61.
- Mbeunkui F, Metge BJ, Shevde LA, Pannell LK. Identification of differentially secreted biomarkers using LC-MS/MS in isogenic cell lines representing a progression of breast cancer. *J Proteome Res.* 2007; 6(8): 2993-3002.
- Qutub AA, Popel AS. A computational model of intracellular oxygen sensing by hypoxia-inducible factor HIF1 alpha. *J Cell Sci.* 2006; 119(Pt 16): 3467-80.
- Lundgren K, Nordenskjöld B, Landberg G. Hypoxia, Snail and incomplete epithelial-mesenchymal transition in breast cancer. *Br J Cancer.* 2009; 101(10): 1769-81.
- Jafari R, Rahbarghazi R, Ahmadi M, Hassanpour M, Rezaie J. Hypoxic exosomes orchestrate tumorigenesis: molecular mechanisms and therapeutic implications. *J Transl Med.* 2020; 18(1): 474. doi: 10.1186/s12967-020-02662-9.
- Welsh JA, Jenkins LM, Kepley J, Lyons GC, Moore DM, Traynor T, *et al.* High sensitivity protein gel electrophoresis label compatible with mass-spectrometry. *Biosensors (Basel).* 2020; 10(11): 160. doi: 10.3390/bios10110160.
- Jacob JT, Coulombe PA, Kwan R, Omary MB. Types I and II keratin intermediate filament. *Cold Spring Harb Perspect Biol.* 2018; 10: a018275. doi: 10.1101/cshperspect.a018275.
- Alix-Panabières C, Vendrell JP, Slijper M, Pellé O, Barbotte E, Mercier G, *et al.* Full-length cytokeratin-19 is released by human tumor cells: a potential role in metastatic progression of breast cancer. *Breast Cancer Res.* 2009; 11(3): R39. doi: 10.1186/bcr2326.
- Palazzolo G, Albanese NN, DI Cara G, Gyax D, Vittorelli ML, Pucci-Minafra I. Proteomic analysis of exosome-like vesicles derived from breast cancer cells. *Anticancer Res.* 2012; 32(3): 847-60.
- Green TM, Alpaugh ML, Barsky SH, Rappa G, Lorico A. Breast Cancer-Derived Extracellular Vesicles: Characterization and Contribution to the Metastatic Phenotype. *Biomed Res Int.* 2015; 2015: 634865. doi: 10.1155/2015/634865.
- Lee JW, Bae SH, Jeong JW, Kim SH, Kim KW. Hypoxia-inducible factor (HIF-1)alpha: its protein stability and biological functions. *Exp Mol Med.* 2004; 36(1): 1-12. doi: 10.1038/emmm.2004.1.
- Hu X, Wu R, Shehadeh LA, Zhou Q, Jiang C, Huang X, *et al.* Severe hypoxia exerts parallel and cell-specific regulation of gene expression and alternative splicing in human mesenchymal stem cells. *BMC Genomics.* 2014; 15: 303. doi: 10.1186/1471-2164-15-303.
- Ni N, Navdeep S, Juan L, Karen M. Mitochondria; reactive oxygen species are require for hypoxia-induced degradation of keratin intermediate filaments. *FASEB J.* 2010; 24(3): 799-809.
- Lubec G, Afjehi-Sadat L. Limitations and pitfalls in protein identification by mass spectrometry. *Chem Rev.* 2007; 107(8): 3568-84.