

Digestibility Value and Fermentation Level of Local Feed-Based Ration for Sheep

Hamdi Mayulu¹, Nur Risma Fauziah¹, Muh Ichsan Haris¹, Marry Christiyanto², and Sunarso²

¹Animal Sciences Departement of Agricultural Faculty, Mulawarman University, EastKalimantan

²Diponegoro University, Semarang, Indonesia

Corresponding author email: mayoelohsptno@yahoo.com

Abstract. Feed Digestibility is an important indicator which can be used as a guideline to determine the amount of nutrient and feed that can be absorbed by digestive tract. The objective of the research was to determine the value dry matter digestibility (DMD), organic matter digestibility (OMD), NH₃ fermentation level and volatile fatty acid (VFA) fermentation level. The research was conducted on January 2018 in Animal Feed and Nutrition Laboratory of Animal Husbandry and Agriculture Faculty, Diponegoro University Semarang. The research method used was *in vitro* analysis with complete randomized design with five treatments and four replicates. The data were analyzed with analysis of variance at 95% significant level, followed by Duncan Multiple Range Test. The results showed that the highest score of DMD was T₅=56.51±0.33%, T₄=56.41±2.11; T₃=55.86±0.83; highest score of OMD was T₅=62.48±0.39%, T₄=61.87±1.60; T₃=60.86±1.10, the highest NH₃ concentration was T₃=6.07±1.55mM, and the highest VFA concentration was T₅=55.25±3.40 mM. Sheep's local feed-based ration was significantly (P<0.05) influenced *in vitro* DMD, OMD, NH₃ fermentation level and VFA fermentation level.

Keywords: Local feed, ration, *in vitro*, digestibility, dry matter, organic matter.

Abstrak. Kecernaan pakan merupakan indikator yang penting sebagai petunjuk menentukan jumlah nutrisi pakan yang mampu diserap oleh saluran pencernaan. Penelitian bertujuan untuk mengetahui nilai kecernaan bahan kering (KcBK), kecernaan bahan organik (KcBO), tingkat fermentasi NH₃ dan tingkat fermentasi *volatile fatty acid* (VFA). Penelitian dilaksanakan pada bulan Januari 2018 di Laboratorium Ilmu Nutrisi dan Makanan Ternak Fakultas Peternakan dan Pertanian Universitas Diponegoro Semarang. Metode Penelitian menggunakan analisis *in vitro* dengan rancangan acak lengkap (RAL) lima perlakuan, dan empat ulangan. Data dianalisis menggunakan *analysis of variance* pada taraf signifikansi 95%, dilanjutkan uji *Duncan Multiple Range Test*. Hasil penelitian menunjukkan bahwa nilai KcBK terbaik berturut-turut T₅=56,51±0,33%, T₄=56,41±2,11; T₃=55,86±0,83; nilai KcBO terbaik T₅=62,48±0,39%, T₄=61,87±1,60; T₃=60,86±1,11; konsentrasi NH₃ terbaik T₃=6,07±1,55mM, dan konsentrasi VFA terbaik T₅=55,25±3,40 mM. Ransum domba berbasis pakan lokal secara *in vitro* berpengaruh nyata (P<0,05) terhadap KcBK, KcBO, tingkat fermentasi NH₃ dan tingkat fermentasi VFA. Kata Kunci: Pakan lokal, ransum, *in vitro*, kecernaan, bahan kering, bahan organik.

Introduction

Sheep is one of meat-producer commodity which is very potential to be promoted to become superior product in order to fulfill domestic or even export market (Mansjoer et al., 2007). Sheep livestock enterprise is special as it can give farmers a sustainable economy; however the average productivity hasn't yet optimum. This phenomenon is closely related due to insufficient feed supply, the genetic quality of the breeding and fault in the raising system (Khanum et al., 2007). Sheep population is very promising which can be seen from the population in 2016 achieved to number 16.462.274 heads. The increasing population

may happen in East Kalimantan because of large field but currently the sheep population is still few only 287 heads in 2016 (Ministry of Agriculture, 2017).

Livestock productivity is influenced by genetic, feed and raising management (Wiyatna et al., 2012). A qualified feed is an important factor to fulfill livestock growth needs to achieve maximum beef production as well as to achieve a successful livestock enterprise (Yakin et al., 2012) so income of farmers increase. Ruminant is a livestock commodity that can utilize a low feedstuffs quality and high crude fiber and then digests into a qualified feeds (Lunagariya et al., 2017). feedstuffs used to formulate ration should contain adequate nutrient content (protein,

carbohydrate, fat, mineral, vitamin and water) for sheep thus can produce optimum result. Ration fed into sheep should contain and formulated in a certain procedure.

Utilization of local feedstuffs material which doesn't compete with human needs is very potential to be researched and studied. A new resource of feedstuffs which doesn't compete with human food chain is needed to be explored. Utilization of local feedstuffs for sheep is an alternative to provide sustain feedstuffs a whole year with quantity and quality that meet sheep's needs standard. The character of sheep itself is very selective to their feeding supply so sheep needs better digestibility value compared with cattle (Adiwimarta, 2012). *Supan-supan* legume (*Neptunia plena* L. Benth) or what is known as *Neptunia plena* and Kolomento grass (*Leersia hexandra* Swartz) or what is known as *Leersia hexandra* are alternative feeds that can be given to sheep.

Forage as the main feed has not yet produced maximum production for sheep, thus a qualified local feedstuffs is very recommended to fulfill the nutrient needs of sheep. The nutrient composition of a ration can be determined based on the ration-chemical composition, while the nutrient utilization can be biologically determined from digestibility and fermentation level test by using *in vitro* method (Khanum et al., 2007). Therefore, the objective of this research was to determine the digestibility score and fermentation rate of sheep's local feedstuffs-based ration.

Materials and Method

The research was conducted on January 2018 at the Laboratory of Animal Feed and Nutrition, Faculty of Animal and Agricultural Sciences, Diponegoro University, Semarang. The materials used in the study were: a) ration consisting of *Neptunia plena* and *Leersia hexandra*, other feed

ingredients (ricebran, corn, palm oil cake and calliandra); b) *in vitro* material using: sheep rumen fluid taken from Boestaman Semarang Slaughterhouse, Mc Dougall solution (artificial saliva), pepsin-HCL solution as protein degrading enzyme, ice water to stop the fermentation process, aquades, CO₂, indicator of metyl red and bromycresol green, saturated sodium carbonate (Na₂CO₃), boric acid solution, 0.5% HCl, phenolptalein 1% indicator, sulfuric acid 0.0055N, vaselin, sulfuric acid (H₂SO₄) 15% and 0.5N NaOH and whatman filter paper 41. The ration was prepared with crude protein content of 11-14% and 60% TDN content.

Experimental Design

The experimental design used in the research was complete randomized design (CRD) with 5 treatments, as shown in Table 1.

In vitro Analysis

In vitro analysis to determine dry matter digestibility (DMD), organic matter digestibility (OMD), NH₃ fermentation rate, and volatile fatty acid (VFA) fermentation rate was carried out using the method of Tilley and Terry (1963).

1. Equation for DMD:

$$\text{DM Digestibility} = \frac{\text{DM weight of sample} - (\text{DM residue} - \text{blanks})}{\text{DM weight of sample}} \times 100\% \dots (1)$$

2. Equation for OMD:

$$\text{OM Digestibility} = \frac{\text{OM weight of sample} - (\text{OM residue} - \text{blanks})}{\text{OM weight of sample}} \times 100\% \dots (2)$$

DM sample = sample weight x % DM

DM residual = weight after oven-CP-filter paper

Blanks = weight after oven-CP-filter paper

OM sample = DM weight of sample x % OM

% OM = 100% DM - (% Ash content in DM)

OM residue = weight after oven - weight after tanur-filter paper

3. Equation for NH₃ Amonia Production:

NH₃ Production (mM) = (ml titran x N H₂SO₄ x 1000) ..(3)

N = Solution normality of H₂SO₄

Table 1. Ration Feedstuffs and Chemical Composition (% DM)

Composition	Treatment				
	T ₁	T ₂	T ₃	T ₄	T ₅
(%).....				
Feedstuffs:					
<i>Leersia hexandra</i>	100.00	-	15.00	20.00	25.00
<i>Neptunia plena</i>	-	100.00	15.00	20.00	25.00
Corn	-	-	3.00	39.00	42.00
Rice bran	-	-	14.00	9.50	1.00
Palm oil cake	-	-	14.50	3.00	2.00
Calliandra	-	-	7.50	8.50	5.00
Total	100.00	100.00	100.00	100.00	100.00
Feed Nutrient:					
DM	85.09	86.89	89.92	89.65	88.69
OM	90.43	95.18	94.30	94.27	94.42
CP	11.28	15.49	12.00	11.92	11.68
TDN*	40.88	38.38	60.00	59.80	59.39

Source: Proximate Analysis Results at the Animal Feed and Nutrition, Faculty of Animal and Agricultural Sciences, Diponegoro University, Semarang (2017)

*Calculation Results according to Sutardi (2001).

4. Equation for VFA:

$$\text{VFA Prod (mM)} = (a-b) \times N \text{ HCl} \times 1000/5 \dots (4)$$

a = Blanks Titran Volume (ml)

b = Sample Titran Volume (ml)

Data Analysis

Data generated from in vitro test were analyzed using Costat program for variance analysis (ANOVA) at 95% significance level followed by a Duncan Multiple Range Test (DMRT) test if there were differences between treatments.

Results and Discussion

Diversity of nutrient value from different feedstuffs requires easier and more efficient nutrient evaluation method (Khanum et al., 2007). Ration fed to sheep should be specially composed and formulated. The nutrient content of ration can be determined from its chemical ration, while nutrient intake can be determined biologically from digestibility test and fermentation rate by means of in vitro method (Khanum et al., 2007). Tilley and Terry's in vitro method is the commonly method used to

evaluate feed's nutrient intake of ruminant and proven to be more accurate to predict digestibility (Mabjeesh et al., 2000). This method has two stages i.e. digested using rument liquid buffer for 48 hours and digested by using pepsin-HCL liquid for 48 hours (Tilley and Terry, 1963).

Feed intake is an important indicator to predict feed's nutrient intake that can be digested by digestive system. Low digestibility indicator that the feedstuffs have less ability to supply nutrient for production and reproduction needs (Mayulu, 2014). In vitro analysis can determine dry matter digestibility (DMD), organic matter digestibility (OMD), NH₃ fermentation level, and volatile fatty acid (VFA) fermentation level (Tilley and Terry, 1963). Average of in vitro analysis of sheep's local feed ration is shown in Table 2.

Dry Matter and Organic Matter Digestibility

Digestibility is an initial indication of various nutrients availability contained in a feedstuffs to be consumed by the livestock. Large amount of digested nutrient intake shows that feed has high digestibility. Feed that has low digestibility shows

that thus feed has less supply of nutrient to livestock (Mayulu, 2012).

Table 2. Average of in vitro DMD, OMD, NH₃ and VFA analysis of sheep fed with local feed ration

Treatment	DMD (%)	OMD (%)	NH ₃ (mM)	VFA (mM)
T ₁	41.93 ^b	52.72 ^b	4.61 ^c	47.75 ^{ab}
T ₂	42.71 ^b	49.13 ^b	4.14 ^c	24.50 ^c
T ₃	55.86 ^a	60.86 ^a	6.07 ^a	41.50 ^b
T ₄	56.41 ^a	61.87 ^a	4.79 ^{bc}	44.00 ^{ab}
T ₅	56.51 ^a	62.48 ^a	5.85 ^{ab}	55.25 ^a

Remarks: Different superscripts show significant different (P<0.05)

T₁=100% of *Leersia hexandra*

T₂= 100 % of *Neptunia plena*

T₃ = Ration (15% *Neptunia plena*+ 15% *Leersia hexandra*+ 70% other feedstuffs)

T₄ = Ration (20% *Neptunia plena*+ 20% *Leersia hexandra*+ 60% other feedstuffs)

T₅ = Ration (25% *Neptunia plena*+ 25% *Leersia hexandra*+ 50% other feedstuffs)

Analysis of variance result on in vitro DMD analysis of sheep's local feed ration (Table 2) showed that the score of DM digestibility from highest to lowest were T₅, T₄, T₃, and followed by T₂, T₁. Average of DM digestibility from those five treatments showed significant difference (P<0.05). DMRT test result of average DMD showed that T₅ was significantly different to T₂ and T₁ but not significantly different to T₄ and T₃, average DMD of T₂ wasn't significantly different to T₁. Analysis result showed that in vitro DMD analysis was influenced by sheep's local feed ration.

The result of this research showed that utilization of local feed as ration with CP 11.68% and TDN 59.39% on the five treatments showed that DM digestibility (56.51%) was better than consumption of single local feed in T₁ (100% of *Leersia hexandra*) (41.93%) and T₂ (100% of *Neptunia plena*) (42.71%) (Table 2). Thus was caused by digestibility was influenced by several factors including the physical shape of feedstuffs, ration composition, comparison composition

amongst the feedstuffs material, feed treatment, enzyme supplementation inside the feed, livestock and feed provision level (Sandi et al., 2015; Priyanto et al., 2017). Research resulted by Trisnadewi et al., (2014) on in vitro DM digestibility of gamal substitution (*Gliricidia sepium*) with calliandra (*Calliandra calothyrsus*) on ration resulted the highest DM digestibility was 38.85%. Thus result was still lower from this research's result i.e. 56.51%.

A decreasing DM digestibility is clearly associated with the increasing fraction of Neutral Detergent Fiber (NDF) and Acid Detergent Fiber (ADF) and increasing of lignification process and decreasing comparison between leaves and stem. Changing in leaves and stem ratio will change the leaves chemical composition, leaves contain of easier digested-feed composition compared with stem. Stem is a part of plant which support the plant structure thus has higher lignin content (Bonneau and Laarveld, 1999).

Feed intake depends on rument microorganism activities as they contribute in fermentation process, while rumen microorganism activities are influenced by food substances contained inside the feedstuffs (Sandi et al., 2015). Number and activity microorganisms reflects increased digestibility (Mayulu, 2014). Ration's feedstuffs which contain of high crude fiber, thicker and more compact cell wall cause lower feed intake. Meanwhile, feedstuffs having lower crude fiber generally has higher feed intake because the cell wall is thin thus can be penetrated by digestive latex (Riswandi et al., 2015). Rumen microorganism can synthesize and secrete cellulose enzyme β 1-4 which roles on hydrolyzing plant cell wall thus high crude fiber feedstuffs can be degraded into high quality feedstuffs (Lunagariya et al., 2017). Microorganism of synthesis up to 30-40 g/kg of digestible organic matter or equal to 187.5-250 g protein of microorganism/kg of digestible organic matter (Mayulu, 2014).

Analysis of variance result on in vitro OMD of sheep's local feed ration is shown in Table 2. It showed average of OM digestibility from the largest to lowest was T₅, T₄, T₃, and followed by T₂, T₁. The score showed that the five treatments were significantly different (P<0.05). Duncan Multiple Range Test result test of OM digestibility showed that T₅ was significantly different to T₂ and T₁, but not significantly different to T₄ and T₃; and T₂ was not significantly different to T₁. Analysis result showed that in vitro OM digestibility score was significantly influenced by sheep's local feed ration.

The result of this research proved that the utilization of local feed as ration with CP 11.68% and TDN 59.39% in T₅ showed that OM digestibility was better (62.48%) compared with single feedstuffs material in T₁ with 100% *Leersia hexandra* (52.72%) and T₂ with 100% *Neptunia plena* (49.13%). Organic matter digestibility is closely associated with DMD because part of DM is OM which contains of CP, CFa, CF and BETN. Result of Trisnadewi et al. (2014) on gamal substitution (*Gliricidia sepium*) with calliandra (*Calliandra calothyrsus*) in ration to in vitro digestibility produced the highest OM was 41.21%. That result was lower than this research's result (62.48%).

Fermentation Level of NH₃ and VFA

Analysis result of in vitro NH₃ fermentation level of sheep's local feed ration (Table 2) showed the highest NH₃ fermentation level was found in T₃ and followed by T₅, T₄, T₁, and T₂. Average of NH₃ fermentation level on five treatments showed significantly different (P<0.05). Duncan Multiple Range Test result on average of NH₃ fermentation concentration showed that T₃ was not significantly different to T₅ and T₄ but significantly different to T₁ and T₂. Analysis result proved that in vitro NH₃ fermentation level was influenced by sheep's local feed ration.

The result of this research proved that sheep ration with CP 12.00% and TDN 60% in T₃ showed

that NH₃ fermentation level was 6.07 mM. Concentration of NH₃ 6,07 mN might be caused by high CP content in ration (12.00%) and nitrogen non protein (NPN) supply in ration as well as high number of microbe contained inside the rumen that used as buffer. Aderinboye et al. (2016) stated that inoculum source (rumen liquid) influenced the NH₃ production. This value showed that T₃ ration can support the microbe growth in rumen because the optimum NH₃ ranging between 3.57-7.14 mM. Low concentration of NH₃ can be caused by low content of protein or protein resistant to degradation which results in inhibition of carbohydrate degradation. Concentration of NH₃ can be a good reference in determining soluble protein supplementation. NH₃ synthesis rate inside the rumen highly depends on chemical structure of protein of the ration (Cristiyanto, 2011). High crude protein content in ration is highly required by microbe as nitrogen source for microbe to build up their body protein (Aderinboye et al., 2016).

Ammonia is a product of fermentation activity in the rumen from protein degradation derived from feed and is the main source of nitrogen (N) for bacteria because most (82%) of bacteria utilizes NH₃ as a source of N. Feed nitrogen 40-60% will be changed into NH₃ by rumen microbe and 50-70% NH₃ will be utilized for rumen microbe protein synthesis. Different NH₃ production can be caused by different crude protein content among treatments. Feed protein in the rumen will be hydrolyzed by proteolytic enzymes rumen microbes to produce oligopeptides which then undergo further digestion into peptides, some pass rumen degradation and some are hydrolyzed into amino acids (Trisnadewi et al., 2014; Sandi et al., 2015).

Concentration of NH₃ in the rumen is a very important quantity to control, because it determines the optimization of rumen microbial growth. High NH₃ concentration is determined by the level of feed protein consumed, degree of degradability, length of feed in the rumen, and

rumen pH. Level of NH_3 needed to support maximum rumen microbial biosynthesis is 3.57-7.14 mM (Sunarso, 2003). The growth of microorganism in the rumen requires an adequate supply of NH_3 derived from feed protein and NPN supplementation in the ration (Riswandi et al., 2017). Rumen microorganism cannot be reproduced if the supply of N in the rumen is limited (Arora, 1995).

The microbial protein precursors are NH_3 and carbon source compound, the increasing NH_3 content in the rumen, the more likely microbial protein are formed and role as the protein source for the body. This condition is a reflection of the contribution of NH_3 to ruminant (Indriani et al., 2013). The energy available after fiber degradation process and rumen pH which is suitable for proteolytic microbial activity affects the concentration of NH_3 (Usman, 2013). Availability of NH_3 is the most important determinant of microbial protein production because the majority of rumen bacteria use NH_3 as a nitrogen source, so it is important to determine NH_3 concentration that supports microbial growth by utilizing NPN (Phesatcha and Wanapat, 2016).

The result of ANOVA on in vitro VFA fermentation level in sheep's local feed ration (Table 2), showed the highest average value of VFA fermentation level, namely at T_5 and sequentially followed by treatment T_1 , T_4 , T_3 , and T_2 . The average fermentation rate of VFA on the five treatments showed very significant differences ($P < 0.05$). The DMRT test result on the average fermentation rate of VFA indicated that T_5 was not significantly different with T_1 , T_3 and T_4 but significantly different with T_2 . The result of the analysis proved that in vitro VFA fermentation level was influenced by sheep's local feed ration.

The result showed that sheep ration with CP 11.68% and TDN 59.39% in the T_5 showed a value of VFA fermentation level of 55.25 mM. This result was still lower than the VFA value for optimum microbial growth ranging from 80-160

mM and the value was influenced by the type of feed given. Volatile fatty acid concentration is influenced by basal feed, type of carbohydrate feed, physical form of feed, consumption level, frequency of feed, and use of chemical additives. Higher VFA concentration indicates an increasing of rumen microbial activity because more organic material is fermented in the rumen (Madrid et al., 2002).

Food substances in the form of carbohydrate and protein strongly support the continuity of metabolic activity in the digestive tract of ruminant. Available carbohydrate is needed by microbes for carbohydrate fermentation and VFA production. The high level of VFA production is an indication that the organic material contained in the ration is very easily degraded by rumen bacteria. Volatile Fatty Acid is formed from the hydrolysis of carbohydrate polysaccharide converted into monosaccharide, especially glucose, then transformed into acetate (C_2), propionate (C_3), butyrate (C_4), isobutane, valerate, isovalerate, methane (CH_4) and CO_2 . Acetate is the main acid formed in the rumen from the degradation of sugar by microbes and is the main energy source, the higher the VFA concentration indicates the fermentation process is more effective, however the concentration of VFA that is too high can cause disruption of rumen system balance (Sandi et al., 2015). In general, comparison of VFA in the rumen ranges from 65% acetate, 20% propionate, 10% butyrate, 5% valerate (Rahmadi et al., 2010). The formation of VFA in the rumen is very important because 70-85% of ruminant energy comes from VFA (Trisnadewi et al., 2014). The sheep's rumen fluid contains of 53-66% acetate, 19-27% propionate, and 12-17% butyrate (Arora, 1995).

Volatile Fatty Acid fermentation in the rumen is influenced by the type of food consumed. Chemical composition, the physical form of feed consumed affect the retention and digestibility of digesta of the rumen and reticulum. Fibrous feeds that have low digestibility will experience

gradual degradation because the first physical contact is slow. This condition results in delayed enzyme work and retention in the rumen, so that only small particles can pass the rumen (Usman, 2013).

Site specific local feed resources (*Leersia hexandra* and *Neptunia plena*) need to be utilized, and started to be given to livestock. Should be carried out by farmers in order to obtain nutrient diversity to meet the needs of sheep and/or ruminants in general. The government is obliged to carry out a socialization of the utilization of site specific local feed resources, so that farmers are able to increase their farming productivity through the use of economic feedstuffs.

Conclusions

Based on the assessment of variance analysis, digestibility value and fermentation level evaluation on sheep's local feed ration under in-vitro method can be concluded as follows:

1. Dry matter digestibility of local feed-based ration had a very significant effect on treatments T₅, T₄, dan T₃ respectively at 56.51%; 56.41%; and 55.86%;
2. Organic matter digestibility of local feed-based sheep ration had a very significant effect on treatments T₅, T₄, and T₃ respectively at 62.48%; 61.87%; and 60.86%;
3. Fermentation of NH₃ level of local feed-based ration had a significant effect, the highest concentration occurred in T₃ treatment i.e. 6.07 mM; and
4. Volatile fatty acid (VFA) fermentation level of local feed-based ration had a significant effect, the highest concentration occurred in T₅ treatment i.e. 55.25 mM.

Acknowledgment

The author would like to say thank you to head and staffs Laboratory of Animal Feed and Nutrition, Faculty of Animal and Agricultural Sciences, Diponegoro University, for the supports during the research.

References

- Aderinboye, R.Y., A.O Akinlolu, M. A. Adeleke, G.O. Najeem, V.O.A. Ojo, O. A. Isah, O. J. Babayemi. 2016. *In vitro* gas production and dry matter degradation of four browse leaves using cattle, sheep and goat inocula. *Slovak J. Anim. Sci* 49 (1): 32-43.
- Adiwimarta. K. I. S. 2012. Pengukuran Kualitas Pakan Sapi (*Cattle Feed Quality Measurement*). PT. Citra Aji Parama. Jakarta.
- Arora, S.P. 1995. Pencernaan Mikroba Pada Ruminansia (*Microbial Digestion in Ruminants*). Gadjah Mada University Press. Cetakan kedua (*Second Edition*). Yogyakarta.
- Bonneau, M., B. Laarveld. 1999. Biotechnology in animal nutrition, physiology and health. *Livestock Production Science* 59: 223–241.
- Christiyanto, M. 2011. Evaluasi Nutrisi Ransum Berpakan Basal Rumput Raja Dengan Ketersediaan Prekursor Nitrogen-Energi Pada Kecernaan Dan Konsumsi Sapi Peranakan Friesian Holstein (*Nutrient Evaluation of King Grass Basal Feed Ration with Availability of Nitrogen-Energy Precursor in Digestion and Consumption of Friesian Holstein Breeds*).Dissertation.Gadjah Mada University. Yogyakarta.
- Indriani, N., T. R. Sutardi and Suparwi. 2013. Fermentasi limbah soun dengan menggunakan *Aspergillus niger* ditinjau dari kadar volatile fatty acid (VFA) total dan amonia (NH₃) secara *in vitro* (*Fermentation of Cassava Flour Noodle Waste Using Aspergillus niger: A Review on Volatile Fatty Acid (VFA) and Ammonia (NH₃) by In Vitro*). *Jurnal Ilmiah Peternakan* 1 (3): 804-812 (*J. of Animal Science*).
- Kementerian Pertanian (*Ministry of Agriculture*). 2017. Sub Sektor Peternakan untuk populasi domba (*Livestock Sub Sector for Sheep Population*). [Downloaded on 01 December 2017]. Available at: http://pertanian.go.id/ap_pages/mod/datanak.
- Khanum, S.A., T. Yaqoob, S. Sadaf, M. Hussain, M. A. Jabbar, H. N. Hussain, R. Kausar, and S. Rehman. 2007. nutritional evaluation of various feedstuffs for livestock production using in vitro gas method. *Pakistan Vet. J* 27 (3): 129-133.
- Lunagariya. P. M., R. M. Gupta, dan S. Parnerkar. 2017. in vitro evaluation of total mixed ration supplemented with exogenous fibrolytic enzymes for crossbred cows. *Veterinary World* 2231-0916: 281-285.
- Madrid, J., M. D. Megias, dan F. Henandeminaz. 2002. In vitro determination of ruminal dry matter and cell wall degradation, and production of fermentation end-product of various by-products. *Anim. Res.*51: 189-199.

- Mabjeesh. S. J., M. Cohen, dan A. Arieli. 2000. In vitro methods for measuring the dry matter digestibility of ruminant feedstuffs: comparison of methods and inoculum source. *Journal of dairy science* 83 (10): 2289-2294.
- Mansjoer, S. S., T. Kertanugraha, and T. Sumantri. 2007. Estimasi jarak genetik antar Domba Garut tipe tangkas dengan tipe pedaging (*Estimation of genetic distance between fighting and meat types of Garut sheep*). *Media Peternakan (J. of Animal Science and Technology)* 30. 129-138.
- Mayulu, H. 2012. Optimasi Sumberdaya Pakan Ruminansia Berbasis Agroekosistem Perkebunan Kelapa Sawit Melalui Teknologi *Complete Feed* (Optimization of Ruminant Feed Resources Based on Oil Palm Plantation Agroecosystems through Complete Feed Technology). *Dissertation*. Diponegoro University. Semarang.
- Mayulu, H. 2014. The nutrient digestibility of locally sheep fed with amofer palm oil byproduct-based complete feed. *Internat. J. Sci. Eng* 7 (2): 106-111.
- Phesatcha. K dan M. Wanapat. 2016. Improvement of nutritive value and *in vitro* ruminal fermentation of *Leucaena* silage by molasses and urea supplementation. *Asian Australas. J. Anim. Sci* 29 (8): 1136-1144.
- Priyanto, A., A. Endraswati, Rizkiyanshah, N. C. Febriyani, T. Nopiansyah and L.K Nuswantara. 2017. Pengaruh pemberian minyak jagung dan suplementasi urea pada ransum terhadap profil cairan rumen (KcBK, KcBO, pH, N-NH₃ dan total mikroba rumen) (*Effect of corn oil and urea supplementation in ration on rumen fluid profile* (DMD, OMD, pH, N-NH₃ and total rumen microbes). *Jurnal Ilmu Ternak (J. of Animal Science)* 17 (1): 1-9.
- Rahmadi, D., Sunarso, J. Achmadi, E. Pangestu, A. Muktiani, M. Christiyanto, Surono, and Surahmanto. 2010. Ruminologi Dasar (*Basic Ruminology*). Diponegoro University. Semarang.
- Riswandi, Muhakka and M. Lehan. 2015. Evaluasi nilai pencernaan secara *in vitro* ransum ternak Sapi Bali yang disuplementasi dengan probiotik bioplus (*Evaluation of in vitro digestibility of Bali cattle ration supplemented with bioplus probiotics*). *J. of Sriwijaya Animal Science* 4 (1): 35-46.
- Riswandi, L. Priyanto, A. Imsya, Meilia, and Nopiyanti. 2017. Pencernaan *in vitro* ransum berbasis rumput Kumpai (*Hymenachne acutigluma*) fermentasi disuplementasi legume berbeda (*In vitro digestion of fermented Kumpai (Hymenachne acutigluma) grass-based ration supplemented with different legume*). *Jurnal Peternakan Sriwijaya (J. of Veterinary)* 18 (2): 303-311.
- Sandi, S., A. I. M. Ali and A. A. Akbar. 2015. Uji *in vitro* wafer ransum komplit dengan bahan perekat yang berbeda (*In vitro test of complete wafer ration with different adhesives*). *Jurnal Peternakan Sriwijaya (J. of Sriwijaya Animal Science)*. 4 (2) pp. 7-16.
- Sunarso. 2003. Pakan ruminansia dalam sistem integrasi ternak-pertanian (Pidato Pengukuhan Guru Besar Universitas Diponegoro 10 September 2003) (*Ruminants Feed in the livestock-farming integration system (Speech for Professor Inauguration of Diponegoro University 10 September 2003)*). Diponegoro University Publishing Board Semarang, Semarang.
- Sutardi, T. 2001. Revitalisasi peternakan sapi perah melalui penggunaan ransum berbasis limbah perkebunan dan suplemen mineral organik (*Revitalizing dairy farming through the use of plantation waste-based ration and organic mineral supplements*). RUT Research VIII.1. Ministry of Research and Technology with the Indonesian Institute of Sciences. Bogor.
- Tilley, J. M. A. and R. A. Terry. 1963. A two stage technique for the *in vitro* digestion of forage crop. *Journal of the British Grassland Society* 18: 104-111.
- Trisnadewi, A.A.A.S., I.G.L.O. Cakra, I. W. Wirawan, I.M. Mudita, and N. L. G. Sumardani. 2014. Substitusi gamal (*Gliricidia sepium*) dengan kaliandra (*Calliandra calothyrsus*) pada ransum terhadap pencernaan *in-vitro* (*In vitro digestion of Gamal (Gliricidia sepium) substitution with calliandra (Calliandra calothyrsus)-based ration*). *Pastura: J. of Tropical Forage Science* 3 (2): 106-109.
- Usman. 2013. Pemberian pakan serat sisa tanaman pertanian (jerami kacang tanah, jerami jagung, pucuk tebu) terhadap evolusi pH, N-NH₃ dan VFA di dalam rumen sapi (*Provision of feed from agricultural residual fibers (peanut straw, corn straw, sugar cane shoots) for the evolution of pH, N-NH₃ and VFA in the cattle rumen*). *Agripet* (13) 2: 53-58.
- Wiyatna. M. F., E. Gurnadi, and K. Mudikdjo. 2012. Produktivitas Sapi Peranakan Ongole pada Peternakan Rakyat di Kabupaten Sumedang (*Productivity of Ongole Breeders in Small Scale Livestock in Sumedang Regency*). *Jurnal Ilmu Ternak (J. of Animal Science)* 12 (2).
- Yakin. E.A., N. Ngadiyono, and R. Utomo. 2012. Pengaruh Substitusi Silase Isi Rumen Sapi pada Pakan Basal Rumput dan Konsentrat terhadap Kinerja Sapi Potong (*Effect of Silage Substitution Fill in Cattle Rumen on Basal Grass Feed and Concentrate on Beef Cattle Performance*). *Buletin Peternakan (Animal Science Bulletin)* 36 (3): 174-180.