

## Correction

# Study of Sperm Reproductive Parameters in Mature Zanjani Viper

Malihe Moshiri, M.Sc.<sup>1</sup>, Fatemeh Todehdehghan, Ph.D.<sup>2\*</sup>, Abdolhossein Shiravi, Ph.D.<sup>1</sup>

1. Department of Basic Science, Islamic Azad University, Damghan Branch, Damghan, Iran  
2. Department of Venomous Animals and Antivenin Production, Vaccine and Serum Research Institute, Karaj, Iran

\*Corresponding Addresses: P.O. Box: 31975/148, Department of Venomous Animals and Antivenin Production, Vaccine and Serum Research Institute, Karaj, Iran  
Email: F.Todehdehghan@rvsri.ac.ir

In this article which was published in Cell J, Vol 16, No 2, Summer 2014, on pages 111-116, the title of table 3 was published incorrectly as: "Sperms concentration, vitality, survival and morphology in Zanjani viper, (mean  $\pm$  SD)". The correct one is "Sperm motility rate in three different parts of vas deferens duct in Zanjani viper".

Table 3: Sperm motility rate in three different parts of vas deferens duct in Zanjani viper

Motile sperm in first region of right duct (%)**	50.62 (0-63)	Left duct (%)	51.14 (48-54)
Grade A	2.06 (0-13)		1.42 (0-7)
Grade B	1.93 (0-4)		2.143 (0-4)
Grade C	46.62 (0-54)		47.57 (43-51)
Grade D (immotile sperm)	49.37 (0-75)		48.86 (46-52)
Motile sperm in middle region of right duct (%)**	54 (0-66)	Left duct (%)	50 (48-52)
Grade A	4.58 (0-13)		0.38 (0-3)
Grade B	3.29 (0-8)		2.00 (0-3)
Grade C	46.11 (0-53)		47.63 (44-49)
Grade D (immotile sperm)	46 (0-57)		50 (48-52)
Motile sperm in final region of right duct (%)**	55 (0-85)	Left duct (%)	49.37 (44-54)
Grade A	4.16 (0-27)		0.00
Grade B	3.83 (0-13)		2.75 (0-8)
Grade C	47 (37-57)		46.62 (42-54)
Grade D (immotile sperm)	45 (15-53)		50.62 (46-56)

\*\*; There is no significant difference in motility rate among three different parts of ducts at  $p > 0.05$ .