

## Metabolic rate and evaporative water loss in the silky starling (*Sturnus sericeus*)

Huan-Huan BAO<sup>1</sup>, Qing-Jian LIANG<sup>2</sup>, Hong-Lei ZHU<sup>2</sup>, Xiao-Qiu ZHOU<sup>2</sup>, Wei-Hong ZHENG<sup>2</sup>, Jin-Song LIU<sup>2,\*</sup>

1. College of Chemistry and Life Sciences, Zhejiang Normal University, Jinhua 321004, China

2. College of Life and Environmental Sciences, Wenzhou University, Wenzhou 325035, China

**Abstract:** To better understand the physiological characteristics of the silky starling (*Sturnus sericeus*), its body temperature ( $T_b$ ), basal metabolic rate ( $BMR$ ), evaporative water loss ( $EWL$ ) and thermal conductance ( $C$ ) elicited by different ambient temperatures ( $T_a$ ) (5–30 °C) were determined in the present study. Our results showed that they have a high  $T_b$  (41.6±0.1 °C), a wide thermal neutral zone ( $TNZ$ ) (20–27.5 °C) and a relatively low  $BMR$  within the  $TNZ$  (3.37±0.17 mL O<sub>2</sub>/g·h). The  $EWL$  was nearly stable below the  $TNZ$  (0.91±0.07 mg H<sub>2</sub>O/g·h) but increased remarkably within and above the  $TNZ$ . The  $C$  was constant below the  $TNZ$ , with a minimum value of 0.14±0.01 mL O<sub>2</sub>/g·h·°C. These findings indicate that the  $BMR$ ,  $T_b$  and  $EWL$  of the silky starling were all affected by  $T_a$ , especially when  $T_a$  was below 20 °C and the  $EWL$  plays an important role in thermal regulation.

**Keywords:** Silky starling (*Sturnus sericeus*); Basal metabolic rate; Body temperature; Evaporative water loss

Endotherms rely primarily on energy metabolism to maintain a constant body temperature. For birds, keeping an optimal energy balance is a key survival strategy, and is primarily achieved by adjusting morphology, physiology and behavior in response to the energy requirements of the environment (Bozinovic, 1992; Weathers, 1997; Lovegrove, 2003). The energy metabolism of birds is affected by many environmental and physiological factors, including body mass, food quality/quantity, especially temperature, which significantly affects the metabolic heat production and thermoregulation (McNab, 2009).

The basal metabolic rate ( $BMR$ ) is the minimum rate of heat production needed to maintain normal physiological mechanisms, and is the minimum energy required by basic metabolic functions necessary to keep animals awake (McKechnie & Wolf, 2004).  $BMR$  is an important parameter in both inter-specific and intra-specific comparisons of energy metabolism, reflecting both energy consumption levels in individuals or species and adaptations of a species to their environments (Burton & Weathers, 2003; McKechnie et al, 2006).

Thermoregulation is conducted by balancing heat production and heat dissipation. The evaporative water loss ( $EWL$ ) is the main way animals dissipate heat and includes both cutaneous ( $CWL$ ) and respiratory water loss ( $RWL$ ) (Dawson, 1982; Tieleman & Williams, 2002).  $EWL$  differs in different temperatures and habitat conditions, e.g. the  $EWL$  of desert animals is lower than that of those living in wet areas (Williams, 1996; Tieleman & Williams, 1999; Tieleman et al, 2003a; Tieleman et al, 2002; Williams et al, 2012).  $EWL$  has an important role in maintaining thermal balance, especially for animals living in hot and dry environments. Consequently, the ability to reduce  $EWL$  has an important adaptive significance (Tieleman et al, 1999; Tieleman & Williams, 2002;

Received: 12 November 2013; Accepted: 20 January 2014

Foundation items: This study was financially supported the National Natural Science Foundation of China (31070366), the Natural Science Foundation (LY13C030005) in Zhejiang Province and the Zhejiang Province 'Xinmiao' Project.

\*Corresponding author, Email: ljs@wzu.edu.cn

Tieleman et al, 2003b). So far, the research on *EWL* has been mainly focusing on the desert rodents (Tieleman & Williams, 1999; Tieleman et al, 2003b; Bozinovic & Gallardo, 2006; Zhu et al, 2008b), especially the relationship of their *EWL* and the ambient temperature ( $T_a$ ) (MacMillen et al, 1977; Donald, 1992; Williams & Tieleman, 2000; Zhu et al, 2008a). However, the publications on the *EWL* in birds are little (Xia et al, 2013).

The silky starling (*Sturnus sericeus*; Passeriformes, Sturnidae) is a summer resident of vast areas of southern and southeastern China that migrates to North Vietnam and the Philippines in winter (MacKinnon & Phillipps, 2000). Their preferred habitats are broadleaf and coniferous-broadleaf mixed forests, but they can also be found in orchards and farmland. The silky starling primarily feeds on insects, fruits and seeds. Studies have found that the silky starling has a high body temperature ( $T_b$ ), a high thermal conductance ( $C$ ), as well as a low *BMR* with a relatively wide thermal neutral zone (*TNZ*) (Zhang et al, 2006).

To investigate the mechanisms of small birds adapting to a broad range of temperature, in this study, the effects of  $T_a$  on the metabolic rate, body temperature and *EWL* of the silky starling have been evaluated. We hypothesized that the warm-adapted bird species, e.g. the silky starling, will reduce their energy expenditure by decreasing *BMR* but increasing  $C$  in response to warm ambient temperatures. Furthermore, because of their mesic habitats, their *EWL* will be higher than the metabolic water production (*MWP*).

## MATERIALS AND METHODS

### Animals

Ten silky starlings were captured in Wenzhou city (N27°29', E120°51'), Zhejiang Province, China, during June of 2008 and were then transported to the laboratory and singly caged in enclosures (50 cm×30 cm×20 cm, length×width×height) under natural photoperiod and temperature of 28 °C. Their mean body mass at capture was 69.01±0.42 g (61.0–77.4 g). Food and water were supplied *ad libitum*. After three weeks acclimation, subjects were exposed to different  $T_a$  ranging from 5 °C to 30 °C.

### Metabolic rates

The *BMR* of silky starlings was expressed as

oxygen consumption per hour per gram of body mass (mL O<sub>2</sub>/g·h), and it was measured by using an open-circuit respirometry system (AEI technologies S-3A/I, USA). The volume of the metabolic chamber was 3.6 L and the water in it was absorbed by silica gel. The experimental temperatures were set at 5, 10, 15, 20, 22.5, 25, 27.5 and 30 °C, respectively. The chamber temperature was controlled within ±0.5 °C by a SHP-250 artificial climate box. Dry CO<sub>2</sub>-free air was pumped through the chamber at 300 mL/min using a flow control system (AEI technologies R-1, USA) calibrated with a general purpose thermal mass flow-meter (TSI 4100 Series, USA) (Xia et al, 2013). All measurements were conducted between 2000h and 2400h. Animals were under fasting 4 h before being put into the metabolic chamber. Metabolic measurements commenced after birds had acclimatized to the chamber for 1 h. To calculate *BMR*, 10 continuous stable minimum recordings were taken over a 1 h period. Body mass and temperature were measured before and after each *BMR* measurement session.

### Evaporative water loss (EWL)

A 'U' tube containing silica gel was placed behind the respiratory chamber and weighed to the nearest 0.1 mg. Any water lost by birds in the experimental chamber would be absorbed by the silica gel and thus could be measured by reweighing the tube at the end of each 30 min experimental period. A 30 min session without bird served as the control. Differences between treatments and control were taken as measurements of *EWL*.

### Thermal conductance (C)

The thermal conductance ( $C$ ) (mL O<sub>2</sub>/g·h·°C) was calculated according to Aschoff (1981):

$$C = MR / (T_b - T_a) \quad (1)$$

where,  $MR$  is the metabolic rate (mL O<sub>2</sub>/g·h),  $T_b$  is the body temperature (°C) and  $T_a$  is the ambient temperature (°C).

The dry thermal conductance ( $C_{dry}$ ) was calculated according to Williams (1999):

$$C_{dry} = (BMR - EWL) / (T_b - T_a) \quad (2)$$

where, oxygen consumption was converted to energy expenditure using 20.09 J/mL O<sub>2</sub> consumed (Schmidt-Nielsen, 1997), the *EWL* converted to energy expenditure using 2.43 J/mg H<sub>2</sub>O consumed (Burton & Weathers, 2003).

### Metabolic water production/evaporative water loss (MWP/EWL)

*EWL* and *MWP* could be used to evaluate the efficiency of water regulation. *MWP* was estimated from oxygen consumption values, assuming that in average, 1 mL of  $O_2$  yields 0.62 mg of metabolic water (Williams, 1999).

### Data Analysis

Data were analyzed via the SPSS statistical package for Windows 18.0. The effects of  $T_a$  on *BMR*, *C* and *EWL* were determined by ANOVA and linear regression analysis. Graphs were generated via Origin 8.0. All results are expressed as mean  $\pm$  SE, with  $P < 0.05$  being considered statistically significant.

## RESULTS

### Body mass and body temperature ( $T_b$ )

Body mass and  $T_b$  were stable over the range of experimental  $T_a$ , and no significant fluctuation in  $T_b$  at  $T_a$  was detected ( $P > 0.05$ ). Mean values of body mass and  $T_b$  were  $69.01 \pm 0.42$  g and  $41.63 \pm 0.08$  °C, respectively.

### Basal metabolic rate (BMR)

*BMR* and  $T_a$  were significantly correlated at  $T_a$  below 20 °C or above 27.5 °C, which could be described by the following equations, respectively (Figure 1):

$$BMR \text{ (mL } O_2/\text{g}\cdot\text{h)} = 5.75(\pm 0.29) - 0.12(\pm 0.02)T_a \quad (3)$$

$(R^2 = 0.963, P < 0.001, n = 40)$

$$BMR \text{ (mL } O_2/\text{g}\cdot\text{h)} = -1.64(\pm 0.25) + 0.19(\pm 0.09)T_a \quad (4)$$

$(R^2 = 0.159, P < 0.05, n = 20)$

*BMR* was remained stable at  $20 \text{ }^\circ\text{C} < T_a < 27.5 \text{ }^\circ\text{C}$ , but was significantly lower at  $T_a = 15 \text{ }^\circ\text{C}$  or  $T_a = 30 \text{ }^\circ\text{C}$ , which suggests that the thermal neutral zone (*TNZ*) of silky starlings was ranged from 20 °C (lower critical temperature) to 27.5 °C (upper critical temperature). The mean value of *BMR* within the *TNZ* was  $3.36 \pm 0.09$  mL  $O_2/\text{g}\cdot\text{h}$ .

### Thermal conductance (C)

Below the *TNZ*, *C* was not significantly correlated with  $T_a$  ( $P > 0.05$ ), averaging at  $0.15 \pm 0.01$  mL  $O_2/\text{g}\cdot\text{h}\cdot^\circ\text{C}$  (Figure 2). Within and above the *TNZ*, *C* increased significantly with  $T_a$  in a linear fashion as described by the following equation:

$$C \text{ (mL } O_2/\text{g}\cdot\text{h}\cdot^\circ\text{C)} = -0.19(\pm 0.03) + 0.02(\pm 0.00)T_a \quad (5)$$

$(R^2 = 0.846, P < 0.001, n = 50)$

### Evaporative water loss (EWL)

*EWL* was stable below the *TNZ* ( $0.91 \pm 0.07$  mg




Figure 1 Metabolic rate of the silky starling as a function of ambient temperature ( $T_a$ ) ( $n=10$ )




Figure 2 Thermal conductance (*C*) of the silky starling as a function of ambient temperature ( $T_a$ ) ( $n=10$ )

$H_2O/\text{g}\cdot\text{h}$ ,  $P > 0.05$ ), but, increased significantly within and above the *TNZ* (Figure 3), which could be described by the following equation:

$$EWL \text{ (mg } H_2O/\text{g}\cdot\text{h)} = -3.38 (\pm 0.33) + 0.26 (\pm 0.01) T_a \quad (6)$$

$(R^2 = 0.961, P < 0.001, n = 50)$

$C_{dry}$  increased with  $T_a$  (Figure 4). At  $20 \text{ }^\circ\text{C} < T_a < 30 \text{ }^\circ\text{C}$ , their relationship could be described by the following equation:

$$C_{dry} \text{ (mL } O_2/\text{g}\cdot\text{h}\cdot^\circ\text{C)} = -0.24 (\pm 0.05) + 0.02 (\pm 0.00) T_a \quad (7)$$

$(R^2 = 0.923, P < 0.001, n = 40)$

The ratio of *EWL/BMR* was positively correlated with  $T_a$  (Figure 5) as described by the following equation:

$$EWL/BMR \text{ (%) = (mg } H_2O/\text{mL } O_2) = -0.73 (\pm 0.18) + 0.07 (\pm 0.01) T_a \quad (8)$$

$(R^2 = 0.970, P < 0.001, n = 50)$

The ratio of *MWP/EWL* was negatively correlated with  $T_a$  (Figure 6). Above the *TNZ*, their relationship could be described by the following equation:

$$MWP/EWL \text{ (%) = } 2.01 (\pm 0.16) - 0.05 (\pm 0.01) T_a \quad (9)$$

$(R^2 = 0.966, P < 0.001, n = 50)$




Figure 3 Evaporative water loss ( $EWL$ ) of the silky starling as a function of ambient temperature ( $T_a$ ) ( $n=10$ )




Figure 6  $MWP/EWL$  of the silky starling as a function of ambient temperature ( $T_a$ ) ( $n=10$ )




Figure 4 Dry thermal conductance ( $C_{dry}$ ) of the silky starling as a function of ambient temperature ( $T_a$ ) ( $n=10$ )




Figure 7  $EHL/HP$  of the silky starling as a function of ambient temperature ( $T_a$ ) ( $n=10$ )




Figure 5  $EWL/BMR$  of the silky starling as a function of ambient temperature ( $T_a$ ) ( $n=10$ )

The percentage of evaporative heat loss to total heat production ( $EHL/HP$ ) was increased with  $T_a$  (Figure 7). Above the  $TNZ$ , their relationship could be described by the following equation:

$$EHL/HP (\%) = -9.01 (\pm 2.07) - 0.80 (\pm 0.08) T_a \quad (R^2 = 0.974, P < 0.01, n = 50) \quad (10)$$

## DISCUSSION

Given the potential conditions posed by global warming or climate change, exploring the physiological characteristics of birds' adaptation to the changeable environments may be of previously unconsidered significance. In this study, we found that the silky starlings had a reduced  $EWL$  at  $T_a > 20$  °C, which was beneficial for them to acclimate to the seasonally hot weather.

### Body temperature

Birds primarily maintain a constant body temperature by altering their heat production and heat dissipation mechanisms (McNab, 1983; Silva, 2006). The body temperature of birds is generally higher than that of mammals because most birds have higher physiological maintenance costs (McNab, 2009). Similarly, small birds' body temperatures are slightly higher than those of

large birds for the same reason (McNab, 2009). Although birds are homeotherms, their  $T_b$  can vary in response to changes in their environment, physiology and behavior, including diurnal rhythm and seasonal changes (Liu et al, 2005; Zheng et al, 2008), e.g. the iridescent hummingbird (*Calypte anna*) (Donald, 1992) and Chinese bulbul (*Pycnonotus sinensis*) (Zhou et al, 2010) are often experience hypothermia in cold weather. Birds reduce their body temperature by reducing metabolic heat production. Low body temperature can decrease energy consumption at night and in winter (McKechnie & Lovegrove, 2002). Our results show that silky starlings maintain their thermostasis by increasing metabolic heat production and lowering thermal conductance at  $T_a$  fell below the *TNZ*, however, increasing metabolic heat production at  $T_a$  within and above the *TNZ*. At high  $T_a$ , to shed excess heat, silky starlings increase their evaporative water loss by increasing thermal conductance and slightly increasing  $T_b$ . This phenomenon has great adaptive significance because during hot summer, the bigger the difference is between  $T_b$  and  $T_a$ , the better the excess heat could be dissipated.

#### Basal metabolic rate and thermal conductance

BMR provides the energy required to maintain the basic activities of life (Dawson, 2003), and it has become a main criterion for assessing both inter- and intra-specific differences in energy metabolism. The BMR of birds, especially in passerine, is higher than that of mammals (Lasiewski & Dawson, 1967; Aschoff and Pohl, 1970; Kendeigh et al, 1977; McNab, 2009). Birds adapt to the environment by changing their BMR (Liu et al, 2005; McNab, 2009). Climate is one of the most important factors determining energy consumption. Therefore, BMR directly reflects the cold tolerance of animals exposed to low temperatures. Birds living in temperate climates, such as the red-billed leiothrix (*Leiothrix lutea*), yellow-browed bunting (*Emberiz achrysophrys*), waxwing (*Bombycilla garrulous*) and black-faced buntings (*Emberiza spodocephala*), are more cold tolerant than tropical birds and have higher BMR because in cold conditions, more energy are required to maintain metabolic process and body temperature (Liu et al, 2005; Li et al, 2005; Wiersma et al, 2007).

In this study, we found that at  $T_a$  below the *TNZ*, the *BMR* of the silky starling increased to produce more heat to maintain basic life activities, a constant  $T_b$  and a balanced energy budget. Within the *TNZ*,  $T_b$  and *BMR*

were relatively stable. Above the *TNZ*, *BMR* slightly increased, presumably to maintain homeostasis. So, we hypothesize that the low *BMR* is an adaptation to the hot and humid environment.

Animal thermal conductance mainly depends on the ratio of their surface area to volume and is also affected by  $T_a$ . Small birds have a relatively large surface area and poor thermal insulation resulting in relatively high thermal conductance (Aschoff, 1981; Bartholomew et al, 1983; Schmidt-Nielsen, 1997). To maintain thermostasis, thermal conductance minimizes at low temperature to retain body heat, and increases at high temperature to dissipate excess heat. The thermal conductance of silky starlings and Chinese bulbuls is 100% and 126% of the value predicted from body size, respectively (Zhang et al, 2006; Lin et al, 2010). Similarly, in summer, the thermal conductance of the waxwing and black-faced bunting is 153% and 157% of the value predicted from body size, respectively (Li et al, 2005). The difference between actual and predicted thermal conductance indicates that in hot conditions, the capacity of birds to dissipate heat enhances. In this study, we found that the thermal conductance of silky starlings increased with the increasing of  $T_a$  above the *TNZ*, suggesting that silky starlings maintain a constant  $T_b$  by shedding excess heat. Below the *TNZ*, thermal conductance minimized at  $0.149 \pm 0.028$  mL  $O_2/g \cdot h \cdot ^\circ C$  (126% of the value predicted from their body size), suggesting that high thermal conductance plays an important role in the adaptations of silky starlings to different climates.

#### Thermal neutral zone

An animal's *TNZ* is the range of environmental temperatures within which temperature regulation can be achieved simply by controlling heat loss, without either metabolic thermogenesis or evaporative cooling. Within the *TNZ*, metabolic rate therefore is unaffected by ambient temperature (Schmidt, 1997). The *TNZ* is itself, however, affected by environmental conditions. In cold and dry climates, such as the arctic, a wide *TNZ* and a low critical temperature in small birds are important to reduce energy consumption and water evaporation, e.g. the *TNZs* of the common redpoll (*Carpodacus roseus*), brambling (*Fringilla montifringilla*), pallas's rosy finch (*Acanthis flammea*), waxwing and black-faced bunting are 25–28 °C, 25–30 °C, 22.5–27.5 °C, 18–27 °C and 20–26 °C respectively (Liu et al, 2004; Li et al, 2005). Conversely, birds that live in hot and humid habitats tend

to have a high thermal conductance, narrow *TNZ* and higher critical temperature, e.g., the *TNZs* of the red-billed leiothrix (*Leiothrix lutea*) and Dunn's Lark (*Eremalauda dunnii*) are 30.0–32.5 °C and 31.5–43.6 °C, respectively (Liu et al, 2005; Tieleman et al, 2002). In this study, a relatively broad *TNZ* in the silky starling was found (20–27.5 °C), which is beneficial for them to acclimate to a wide range of temperatures and to decrease energy expenditure in cold weather.

### Evaporative water loss

An animal's metabolism includes both substance and energy metabolism, and water metabolism plays an important role between these (Bozinovic & Gallardo, 2006). Birds excrete excess water through pulmonary, respiratory and skin surface evaporative water loss. *EWL* is affected by  $T_a$  and humidity, e.g. in four different lark species, Tieleman & Williams (2002) found that their *EWLs* were low at low  $T_a$ , but increased rapidly at  $T_a > 35$  °C; in 12 different lark species, Tieleman et al (2003a) found that their *EWLs* were negatively correlated with the aridity gradient of habitats. Our results showed that

because body temperature regulation was not necessary in the balanced heat budget below the *TNZ* ( $MWP/EWL=1$ ,  $T_a=20$  °C), *EWL* was irrelevant with  $T_a$ . Above the *TNZ* ( $T_a > 20$  °C), birds need more water to maintain the balance of water loss in hot season, so the *EWL* increased but  $MWP/EWL$  decreased with  $T_a$ .  $EHL/HP$  increased with  $T_a$  at  $20$  °C  $< T_a < 30$  °C, and maximized (15.05%) at  $T_a=30$  °C. We presume that this was due to the increased heat production and therefore unbalanced heat budget in high temperature (above the *TNZ*). In our study, silky starlings dissipated excess heat and maintained a stable  $T_b$  by increasing their thermal conductance and *EWL*, suggesting that *EWL* plays an important role in the thermoregulation of small birds in hot climates.

In conclusion, our results indicate that silky starlings have a low *BMR*, a high body temperature, a high thermal conductance, a high *EWL*, as well as a relatively wide *TNZ*, and among which, *EWL* plays an important role in the metabolism and body temperature regulation. We presume that these physiological characteristics in birds are outputs of multiple-mechanism adaptations.

### References

- Aschoff J. 1981. Thermal conductance in mammals and birds: Its dependence on body size and circadian phase. *Comparative Biochemistry and Physiology Part A*[0], **69**(4): 611-619.
- Aschoff J, Pohl H. 1970. Der ruheumsatz von vögeln als function der tageszeit and der körpergröße. *Journal of Ornithology*, **111**(1): 38-47.
- Bartholomew GA, Vleck CM, Bucher TL. 1983. Energy metabolism and nocturnal hypothermia in two tropical passerine frugivores. *Manacus vitellinus* and *Pipra mentalis*. *Physiological Zoology*, **56**(3): 370-379.
- Bozinovic F. 1992. Rate of basal metabolism of grazing rodents from different habitats. *Journal of Mammalogy*, **73**(2): 379-384.
- Bozinovic F, Gallardo P. 2006. The water economy of South American desert rodents: From integrative to molecular physiological ecology. *Comparative Biochemistry and Physiology Part C*, **142**(3-4): 163-172.
- Burton CT, Weathers WW. 2003. Energetics and thermoregulation of the Gouldian finch *Erythrura gouldiae*. *The Emu*, **103**: 1-10.
- Dawson WR. 1982. Evaporative losses of water by birds. *Comparative Biochemistry and Physiology Part A*[0], **71**(4): 495-509.
- Donald RP. 1992. Effect of temperature and humidity on evaporative water loss in Anna's hummingbird (*CMypte anna*). *Journal of Comparative Physiology B*, **162**(1): 74-84.
- Dawson WR. 2003. Plasticity in avian responses to thermal challenges— an essay in honor of Jacob Marder. *Israel Journal of Zoology*, **49**: 95-109.
- Kendeigh SC, Dol'nik VR, Gavrillo VM. 1977. Avian energetic. In: Pinowski J, Kendeigh S C. Granivorous Birds in Ecosystems. Cambridge University Press, 127-204.
- Lovegrove BG. 2003. The influence of climate on the basal metabolic rate of small mammals: A slow fast metabolic continuum. *Journal of Comparative Physiology B*, **173**(2): 87-112.
- Lasiewski RC, Dawson WR. 1967. A re-examination of the relation between standard metabolic rate and body weight in birds. *The Condor*, **69**(1): 13-23.
- Liu JS, Wang DH, Sun RY. 2005. Climatic adaptations in metabolism of four species of small birds in China. *Acta Zoologica Sinica*, **51**(1): 24-30.
- Lin L, Wang LH, Liu JS. 2010. Metabolism and thermoregulation in crested mynas (*Acridotheres cristatellus*). *Chinese Journal of Zoology*, **45**(5): 47-53.
- Li M, Liu JS, Han HL, Zhang HJ, Fang H. 2005. Metabolism and thermoregulation in waxwings (*Bombycilla garrulous*) and black-faced buntings (*Emberiza spodocephala*). *Zoological Research*, **26**(3): 287-293.
- Liu JS, Wang DH, Wang Y, Chen MH, Song CG, Sun RY. 2004. Energetics and thermoregulation of the *Carpodacus roseus* *Fringilla montifringilla* and *Acanthis flammea*. *Acta Zoologica Sinica*, **50**(3): 357-363.
- MacKinnon J, Phillipps K. 2000. A field guide to the birds of China.

- London: Oxford University Press, 312-340.
- McKechnie AE, Wolf BO. 2004. The allometry of avian basal metabolic rate: Good predictions need good data. *Physiological and Biochemical Zoology*, **77**(3): 502-521.
- McKechnie AE, Freckleton RP, Jetz W. 2006. Phenotypic plasticity in the scaling of avian basal metabolic rate. *Proceedings of the Royal Society of London B*, **273**(1589): 931-937.
- McNab BK. 1983. Energetics, body size, and the limits of endothermy. *Journal of Zoology*, **199**(1): 1-29.
- McNab BK. 2009. Ecological factors affect the level and scaling of avian BMR. *Comparative Biochemistry and Physiology Part A*, **152**(1): 22-45.
- McKechnie AE, Lovegrove BG. 2002. Avian facultative hypothermic responses: A review. *The Condor*, **104**(4): 705-724.
- Macmillen RE, Whittow GC, Christophe EA, Ebisu RJ. 1977. Oxygen consumption, evaporative water loss, and body temperature in the sooty tern. *The Auk*, **94**(1): 72-79.
- Silva JE. 2006. Thermogenic mechanisms and their hormonal regulation. *Physiological Reviews*, **86**(2): 435-464.
- Schmidt-Nielsen K. 1997. *Animal Physiology: Adaptation and Environment*. 5<sup>th</sup> ed. London: Cambridge University Press, 169-214.
- Tieleman BI, Williams JB. 1999. The role of hyperthermia in the water economy of desert birds. *Physiological and Biochemical Zoology*, **72**(1): 87-100.
- Tieleman BI, Williams JB, Michaeli E, Pinshow B. 1999. The role of the nasal passages in the water economy of crested larks and desert larks. *Physiological and Biochemical Zoology*, **72**(2): 219-226.
- Tieleman BI, Williams JB. 2002. Cutaneous and respiratory water loss in larks from arid and mesic environments. *Physiological and Biochemical Zoology*, **75**(6): 590-599.
- Tieleman BI, Williams JB, Buschur ME. 2002. Physiological adjustments to arid and mesic environments in larks (Alaudidae). *Physiological and Biochemical Zoology*, **75**(3): 305-313.
- Tieleman BI, Williams JB, Bloomer P. 2003a. Adaptation of metabolism and evaporative water loss along an aridity gradient. *Proceedings of the Royal Society of London B*, **270**: 207-214.
- Tieleman BI, Williams JB, Buschur ME, Brown CR. 2003b. Phenotypic variation of larks along an aridity gradient: are desert birds more flexible? *Ecology*, **84**(7): 1800-1815.
- Weathers WW. 1997. Energetics and thermoregulation by small passerines of the humid, low land tropics. *The Auk*, **114**(3): 341-353.
- Williams JB. 1996. A phylogenetic perspective of evaporative water loss in birds. *The Auk*, **113**(2): 457-472.
- Williams JB, Muñoz-Garcia A, Champagne A. 2012. Climate change and cutaneous water loss of birds. *The Journal of Experimental Biology*, **215**(Pt7): 1053-1060.
- Williams JB, Tieleman BI. 2000. Flexibility in basal metabolic rate and evaporative water loss among hoopoe larks exposed to different environmental temperatures. *The Journal of Experimental Biology*, **203**(Pt 20): 3153-3159.
- Williams JB. 1999. Heat production and evaporative water loss of Dune larks from the Namib desert. *The Condor*, **101**(2): 432-438.
- Wiersma P, Muñoz-Garcia A, Walker A, Williams JB. 2007. Tropical birds have a slow pace of life. *Proceedings of the National Academy of the Sciences of the United States of America*, **104**(22): 9340-9345.
- Xia SS, Yu AW, Zhao LD, Zhang HY, Zheng WH, Liu JS. 2013. Metabolic thermogenesis and evaporative water loss in the Huamei *Garrulax canorus*. *Journal of Thermal Biology*, **38**(8): 576-581.
- Zhu WL, Jia T, Lian X, Wang ZK. 2008a. Evaporative water loss and energy metabolism in two small mammals, voles (*Eothenomys miletus*) and mice (*Apodemus chevrieri*), in Hengduan Mountains region. *Journal of Thermal Biology*, **33**(6): 324-331.
- Zhu WL, Yang YH, Jia T, Lian X, Wang ZK, Gong ZD, Guo XG. 2008b. Evaporative water loss and body temperature regulation in *Eothenomys miletus* and *Apodemus chevrieri*. *Acta Theriologica Sinica*, **28**(1): 65-74.
- Zhou W, Wang YP, Chen DH, Liu JS. 2010. Diurnal rhythms of Chinese bulbul (*Pycnonotus sinensis*) body temperature, body mass, and energy metabolism. *Chinese Journal of Ecology*, **29**(12): 2395-2400.
- Zheng WH, Liu JS, Jiang XH, et al. 2008. Seasonal variation on metabolism and thermoregulation in Chinese bulbul. *Journal of Thermal Biology*, **33**(6): 315-319.
- Zhang YP, Liu JS, Hu XJ, et al. 2006. Metabolism and thermoregulation in two species of passerines from southeastern China in summer. *Acta Zoologica Sinica*, **52**(4): 641-647.