




Contents lists available at ScienceDirect

Asian Pacific Journal of Tropical Biomedicine

journal homepage: www.elsevier.com/locate/apjtb

Document heading doi:10.12980/APJTB.4.2014C742 © 2014 by the Asian Pacific Journal of Tropical Biomedicine. All rights reserved.

Bartonella species in small mammals and their potential vectors in AsiaTawisa Jiyipong^{1,4,5,6}, Sathaporn Jittapalpong^{2,6}, Serge Morand^{3,7}, Jean–Marc Rolain^{1*}¹Research Unit on Infectious and Emerging Tropical Diseases (URMITE), CNRS–IRD–INSERM UMR 7278, IHU Méditerranée Infection, Faculty of Medicine and Pharmacy, Aix–Marseille–University, Marseille, France²Department of Parasitology, Faculty of Veterinary Medicine, Kasetsart University, Bangkok, Thailand³Institut Sciences of Evolution, UMR 5554 CNRS–IRD–UM2, CC65, Université de Montpellier 2, F–34095, Montpellier, France⁴Center for Agricultural Biotechnology (AG–BIO/PEDRO–CHE), Kasetsart University, Kamphaeng Saen Campus, Nakhon Pathom, 73140, Thailand⁵Center of Excellence on Agricultural Biotechnology (AG–BIO/PERDO–CHE), Bangkok 10900, Thailand⁶Center of Advanced Studies for Agriculture and Food, KU Institute for Advanced Studies, Kasetsart University, Bangkok 10900, Thailand (CASAF, NRU–KU, Thailand)⁷Walai Rukhvej Botanical Research Institute, Mahasarakham University, Mahasarakham, Thailand

PEER REVIEW

Peer reviewer

Dr. Hossein Ghalehnoei, University of Medical Science, Tehran, Iran.
 Tel: +989385518026, E–mail: hossein.ghalehnoei58@gmail.com
 Co–reviewers: Misael Chinchilla, San José, Costa Rica,

Comments

This is a good research about the presence of various species of *Bartonella* and its role at causing different infections. The consideration of authors to various reservoirs of this bacteria in animals that a little have been already surveyed is interesting. Diagnostic tests referred in this article are very useful.

Details on Page 764

ABSTRACT

In this article, authors review the current knowledge of *Bartonella* infection in small mammals including rodents, insectivores, bats and exotic small mammal pets and their vectors in Asia. Species of *Bartonella* are Gram–negative intracellular bacteria that infect erythrocytes of various mammalian and non–mammalian animals and mainly transmitted by blood sucking arthropod vectors. The genus *Bartonella* includes several species of important human diseases with severe clinical signs. Several new *Bartonella* species were isolated from rodents and other small mammals, and from human patients in Asia. *Bartonella* species are identified using standard polymerase chain reaction amplification and a sequencing targeting two housekeeping genes (*gltA* and *rpoB*) and the internal transcribed spacer fragment. Authors also discuss the implications in term of potential emerging zoonotic diseases.

KEYWORDS

Bartonella, Small mammals, Rodents, Shrews, Asia**1. Introduction**

Member of the genus *Bartonella* are fastidious, hemotropic, Gram–negative and aerobic bacilli bacteria belonging to the class Alphaproteobacteria. Several species have been implicated in causing human diseases, ranging with short–term fever to severe endocarditis. Five species are known to be causative of significant human diseases:

Bartonella bacilliformis (*B. bacilliformis*) is the causative agent of Oroya fever and verruga peruana (Peruvian wart); *Bartonella quintana* (*B. quintana*) is the causative agent of trench fever; *Bartonella henselae* (*B. henselae*) is the causative agent of cat–scratch disease[1–3]. Additionally, several species have been described as coincident zoonotic infection agents including *Bartonella alsatica* (*B. alsatica*) [4], *Bartonella koehlerae* (*B. koehlerae*)[5], *Bartonella vinsonii*

*Corresponding author: Jean–Marc Rolain, Research unit on infectious and emerging tropical diseases (URMITE) CNRS–IRD–INSERM UMR 7278, IHU Méditerranée Infection, Faculty of Medicine and Pharmacy, Aix–Marseille–University, Marseille, France.

E–mail: jean-marc.rolain@univ-amu.fr

Foundation Project: Supported by the French ANR CERoPath project (number ANR 07 BDIV 012) and the French ANR BiodivHealthSEA project (number ANR 11 CPEL 002).

Article history:

Received 28 Dec 2013

Received in revised form 20 Feb, 2nd revised form 3 Apr, 3rd revised form 20 Apr 2014

Accepted 20 May 2014

Available online 14 Sep 2014

subsp. *berkhoffii* (*B. vinsonii* subsp. *berkhoffii*)[6,7], *Bartonella tamiae* (*B. tamiae*)[8], *Bartonella rochalimae* (*B. rochalimae*) [9], *Bartonella washoensis* (*B. washoensis*)[10] and *Candidatus Bartonella mayotimonensis*[11].

Bartonella species parasitize the erythrocytes, endothelial cells, monocytes and dendritic cells of mammals[12,13]. Additionally, a *Bartonella* species has been described in loggerhead sea turtle[14]. Bacteria of the genus *Bartonella* are widespread in domestic and wild animals and are transmitted by a great variety of bloodsucking arthropods including fleas, mites, sand flies and ticks[15,16]. Since 1990s, numerous studies have demonstrated that several mammals such as cats, dogs, rabbits and rodents are potential reservoir hosts of *Bartonella*. Until now, new host species such as insectivores, bats and exotic pets are continually added in the reservoir hosts' list. This review provides an update of reservoir host species focusing on small mammals, their potential vectors and case reports of *Bartonella* infection in Asia, a hotspot of emerging infectious diseases[17,18]. Hosts such as cats, dogs, and rabbits have been excluded from this review since several review articles have been already written on these reservoir hosts[19,20].

2. Taxonomy and bacteriology of *Bartonella*

Bacteria of the genus *Bartonella* belong to the family Bartonellaceae, order Rhizobiales, class Alphaproteobacteria, and phylum Proteobacteria. *Bartonella* are closely related with *Brucella* species and *Agrobacterium tumefaciens*. *Bartonella* genus includes a large diversity of *Bartonella* species, which were identified from different host species. Since the development of more efficient molecular tools for detection, genetic criteria and species identification are greatly improved[21], and the description of new *Bartonella* species has rapidly increased over the last 10 years and is still continually growing. The genus currently contains more than 30 species and 3 subspecies including *B. alsatica*, *Bartonella australis*, *B. bacilliformis*, *Bartonella birtlesii* (*B. birtlesii*), *Bartonella bovis* (*B. bovis*), *Bartonella capreoli* (*B. capreoli*), *Bartonella chomelii*, *Bartonella coopersplainsensis* (*B. coopersplainsensis*), *Bartonella clarridgeiae* (*B. clarridgeiae*), *Bartonella doshiae* (*B. doshiae*), *Bartonella durdenii*, *Bartonella grahamii* (*B. grahamii*), *B. henselae*, *Bartonella japonica* (*B. japonica*), *B. koehlerae*, *Bartonella melophagi*, *Bartonella phoceensis* (*B. phoceensis*), *Bartonella queenslandensis* (*B. queenslandensis*), *Bartonella quintana* (*B. quintana*), *Bartonella rattaaustraliansi* (*B. rattaaustraliansi*), *Bartonella rattimassiliensis* (*B. rattimassiliensis*), *B. rochalimae*, *Bartonella schoenbuchensis*, *Bartonella silvatica* (*B. silvatica*), *Bartonella silvicola*, *B. tamiae*, *Bartonella taylorii* (*B. taylorii*), *Bartonella tribocorum* (*B. tribocorum*), *Bartonella vinsonii* subsp. *arupensis* (*B. vinsonii* subsp. *arupensis*), *B. vinsonii* subsp. *berkhoffii*, *Bartonella vinsonii* subsp. *vinsonii* (*B. vinsonii* subsp. *vinsonii*), *B. washoensis*, *Candidatus Bartonella antechini*, *Candidatus Bartonella mayotimonensis*, *Candidatus Bartonella thailandensis* (Table 1). Of these, several species may cause either asymptomatic

or mild diseases or severe diseases. Bacteria from this genus are fastidious to grow *in vitro*. The culture on blood agar requires 7 to 45 d in primary isolation. Microscopically, the *Bartonella* species are Gram-negative bacilli. Several *Bartonella* species are flagellate, *B. bacilliformis*, *B. bovis* and *B. clarridgeiae*[22,23].

Table 1

List of the currently known described species of *Bartonella*.

<i>Bartonella</i> species	First detection	Country	Reference
<i>B. alsatica</i>	Wild rabbit	France	[59]
<i>B. australis</i>	Gray kangaroo	Australia	[60]
<i>B. bacilliformis</i>	Human	Europe	[61]
<i>B. birtlesii</i>	Mouse	German	[62]
<i>B. bovis</i>	Cow	France	[62]
<i>B. capreoli</i>	Deer	France	[63]
<i>B. chomelii</i>	Domestic cattle	France	[64]
<i>B. clarridgeiae</i>	Cat	USA	[65]
<i>B. coopersplainsensis</i>	Rat	Australia	[66]
<i>B. doshiae</i>	Field vole	United Kingdom	[67]
<i>B. durdenii</i>	Squirrel	USA	unpublished data
<i>B. elizabethae</i>	Human	USA	[68]
<i>B. grahamii</i>	Bank vole	United Kingdom	[67]
<i>B. henselae</i>	Cat	USA	[69]
<i>B. japonica</i>	Mouse	Japan	[70]
<i>B. koehlerae</i>	Cat	USA	[71]
<i>B. melophagi</i>	Sheep		unpublished data
<i>B. merieuxii</i>	Canid	Irak	[72]
<i>B. peromysci</i>	Mouse		[67]
<i>B. phoceensis</i>	Wild rat	France	[73]
<i>B. queenslandensis</i>	Rat	Australia	[66]
<i>B. quintana</i>	Human		[74]
<i>B. rattaaustraliansi</i>	Rat	Australia	[66]
<i>B. rattimassiliensis</i>	Rat	France	[73]
<i>B. rochalimae</i>	Human	USA	[9]
<i>B. schoenbuchensis</i>	Deer	German	[75]
<i>B. silvatica</i>	Mouse	Japan	[70]
<i>B. silvicola</i>	Bat		unpublished data
<i>B. talpae</i>	Mole		[59]
<i>B. tamiae</i>	Human	Thailand	[6]
<i>B. taylorii</i>	Mouse	United Kingdom	[67]
<i>B. tribocorum</i>	Rat	France	[76]
<i>B. vinsonii</i> subsp. <i>arupensis</i>	Cattle		[77,78]
<i>B. vinsonii</i> subsp. <i>berkhoffii</i>	Dog		[79]
<i>B. vinsonii</i> subsp. <i>vinsonii</i>	Vole		[80]
<i>B. volans</i>	Squirrel	USA	unpublished data
<i>B. washoensis</i>	Human	USA	unpublished data

3. *Bartonella* genomes

The genome sizes of *Bartonella* species range from 1.5 to 2.5 Mb. Several *Bartonella* species including *B. bacilliformis*, *B. birtlesii*, *B. clarridgeiae*, *B. grahamii*, *B. henselae*, *B. quintana*, *B. rattimassiliensis*, *B. rattaaustraliansi* and *B. tribocorum* were completely sequenced and two species (*B. grahamii* and *B. tribocorum*) contain plasmid. The first genomes have been described for *B. henselae* and *B. quintana* with 1.9 and 1.6 Mb, respectively[24], followed by the sequencing of the genomes of *B. bacilliformis* (1.4 Mb) (the Institute for Genomic Research, unpublished), *B. clarridgeiae* (1.5 Mb)[25], *B. grahamii* (2.3 Mb)[26], *B. tribocorum* (2.6 Mb)[27], *B. rattimassiliensis* (2.0 Mb)[28], *B. rattaaustraliansi* (2.1 Mb)[29], *B. birtlesii* (1.8 Mb)[30] and *B.*

quintana (1.6 Mb)^[31] among others. The guanine–cytosine content of *Bartonella* species range from 38.5 mol% for *B. quintana* to 41.1 mol% for *B. vinsonii*^[2]. Additionally, some species such as *B. bacilliformis*, *B. henselae*, *B. quintana* and *B. vinsonii* subsp. *berkhoffii* have been shown to contain phage^[32]. The phage particles from *B. bacilliformis* and *B. vinsonii* subsp. *berkhoffii* were tailed, whereas those from *B. henselae* and *B. quintana* lacked tails, but all contained 14 kb linear, double–stranded DNA, packaged in a round to head.

To date, *Bartonella* species are identified using standard polymerase chain reaction (PCR) amplification and a sequencing targeting two housekeeping genes (*gltA* and *rpoB*) and the internal transcribed spacer fragment^[2].

4. Hosts and reservoirs

Bartonella species may show either low specificity with some species infecting several different host species, while some other species show high specificity by infecting a single host. For example, *B. bovis* generally infects only one ruminant species and is seldom associated with other animals. The prevalence of *Bartonella* in both wild and domestic mammals has been studied in many different countries. To date, several animal species have been reported as a potential reservoir hosts, such as cats, dogs, rabbits, ruminants, monkeys, marsupials, marine mammals,

bats, insectivores and rodents^[19,33].

4.1. Rodents

Wild rodents are known to be important reservoir hosts of various pathogens, which are causative of human illnesses including Bartonellosis. Of the current species of the genus *Bartonella*, fifteen species including *B. birtlesii*, *Bartonella elizabethae* (*B. elizabethae*), *B. coopersplainsensis*, *B. doshaiae*, *B. grahamii*, *B. japonica*, *B. phoceensis*, *B. queenslandensis*, *B. rattaaustraliani*, *B. rattimassiliensis*, *B. silvatica*, *B. taylorii*, *B. tribocorum*, *B. vinsonii* subsp. *arupensis*, *B. vinsonii* subsp. *vinsonii*, *B. washoensis* have been isolated from various rodent species^[2]. Of these, *B. birtlesii*, *B. elizabethae*, *B. grahamii*, and *B. washoensis* are causative agent of human illnesses.

In Asia, the occurrence of *Bartonella* infection in rodent populations have been reported in several countries including China, Japan, Taiwan, Lao PDR, Cambodia, Thailand, Indonesia, Bangladesh and Israel. *Bartonella* is present in most surveyed rodent populations, with an overall prevalence in the sites investigated ranging from 6% in Korea^[34] to 47% in China^[35]. Table 2 summarizes *Bartonella* species, host species and geographic distribution of *Bartonella* in Asia. *Bartonella* infections are highly prevalent in rodents in China, Korea, Japan, Russia and Taiwan ranging from 8.6% to 82.3%. Several *Rattus* species have been found as highly infected including the house rat *Rattus tanezumi* (*R. tanezumi*). Several species of the

Table 2

Bartonella infection and host species in East Asia.

Country	<i>Bartonella</i> infection	Diagnostic test		Host species and prevalence of each species	Type of animal				Reference	
		PCR	Culture		Rodent	Shrew	Bat	Exotic pet		
China	<i>B. elizabethae</i>	+	+	<i>Rattus tanezumi</i> subsp. <i>flavipectus</i> , 41.4% (24/58)	+	–	–	–	[35]	
	<i>B. grahamii</i>	+	+	<i>Rattus norvegicus</i> , 42.9% (3/7)	+	–	–	–	[35]	
	<i>B. tribocorum</i>		+	+	<i>Apodemus chevrieri</i> , 62.5% (20/32)	+	–	–	–	[35]
					<i>Apodemus draco</i> , 33.3% (2/6)	+	–	–	–	[35]
					<i>Apodemus latronum</i> , 71.4% (5/7)	+	–	–	–	[35]
					<i>Eothenomys miletus</i> , 18.8% (3/16)	+	–	–	–	[35]
					<i>R. rattus</i> , 28.6% (2/7)	+	–	–	–	[81]
	<i>B. grahamii</i>	+	–	<i>Rattus losea</i> , 31% (14/45)	+	–	–	–	[81]	
	<i>Bartonella</i> unknown species		+	–	<i>Rattus norvegicus</i> , 30% (7/23)	+	–	–	–	[81]
					<i>Suncus murinus</i> , 16% (7/43)	+	–	–	–	[81]
					<i>Niviventer confucianus</i> , 4% (1/28)	+	–	–	–	[81]
					<i>Apodemus agrarius</i> , 46.8% (90/192)	+	–	–	–	[81]
					<i>Apodemus peninsulae</i> , 28.6% (2/7)	+	–	–	–	[81]
					<i>Eothenomys melanogaster</i> , 24% (9/37)	+	–	–	–	[81]
				<i>Microtus fortis</i> , 22% (2/9)	+	–	–	–	[81]	
				<i>Apodemus speciosus</i> , 51.2% (116/224)	+	–	–	–	[82]	
Japan	<i>B. grahamii</i>	+	+	<i>Apodemus argenteus</i> , 37.1% (13/35)	+	–	–	–	[82]	
	<i>B. phoceensis</i>		+	<i>R. rattus</i> , 3.7% (11/297)	+	–	–	–	[82]	
			+	<i>R. rattus</i> , 1.0% (3/297)	+	–	–	–	[82]	
	<i>B. rattimassiliensis</i>		+	<i>Apodemus speciosus</i> , 8.9% (2/224)	+	–	–	–	[82]	
			+	<i>Myodes rufocanus</i> subsp. <i>bedfordiae</i> , 23.5% (4/17)	+	–	–	–	[82]	
	<i>B. taylorii</i>		+	<i>Apodemus speciosus</i> , 1.8% (4/224)	+	–	–	–	[82]	
			+	<i>Apodemus argenteus</i> , 5.7% (2/35)	+	–	–	–	[82]	
			+	<i>R. rattus</i> , 2.0% (6/297)	+	–	–	–	[82]	
	Unknown species		+	<i>Apodemus speciosus</i> , 8.0% (18/224)	+	–	–	–	[82]	
			+	<i>Apodemus argenteus</i> , 11.4% (4/35)	+	–	–	–	[82]	
<i>B. clarridgeiae</i>	+	+	<i>Spermophilus dauricus</i> , 40% (4/10)	–	–	–	+	[48]		
<i>B. elizabethae</i>	+	+	<i>Sciurus vulgaris</i> subsp. <i>orientis</i> , 20% (2/10)	–	–	–	+	[48]		
<i>B. grahamii</i>	+	+	<i>Tamias sibiricus</i> , 41.4% (12/29)	–	–	–	+	[48]		

continued Table 2

	<i>B. washoensis</i>	+	+	<i>Pteromys volans</i> , 50% (5/10)	-	-	-	+	[48]
				<i>Callosciurus notatus</i> , 63.3% (19/30)	-	-	-	+	[48]
				<i>Tamiasciurus hudsonicus</i> , 16.7% (3/18)	-	-	-	+	[48]
				<i>Glaucomys volans</i> , 60% (6/10)	-	-	-	+	[48]
				<i>Pachyuromys duprasi</i> , 72.2% (13/18)	-	-	-	+	[48]
				<i>Acomys cahirinus</i> , 9.7% (3/31)	-	-	-	+	[48]
				<i>Acomys russatus</i> , 61.5% (8/13)	-	-	-	+	[48]
				<i>Psammomys obesus</i> , 60% (6/10)	-	-	-	+	[48]
				<i>Sekeetamys calurus</i> , 100% (10/10)	-	-	-	+	[48]
				<i>Gerbillus pyramidum</i> , 90% (9/10)	-	-	-	+	[48]
				<i>Jaculus orientalis</i> , 81.3% (13/16)	-	-	-	+	[48]
				<i>Jaculus jaculus</i> , 75% (6/8)	-	-	-	+	[48]
Korea	<i>B. elizabethae</i>	+	-	<i>Apodemus agrarius</i> , 14.2% (53/373)	+	-	-	-	[43]
				<i>Eothenomys melanogaster</i> , 11.1% (1/9)	+	-	-	-	[43]
				<i>Crocidura lasiura</i> , 12.1% (4/33)	-	+	-	-	[43]
Taiwan	<i>B. elizabethae</i>	+	+	<i>Apodemus agrarius</i> , 6.7% (24/358)	+	-	-	-	[34]
	<i>B. elizabethae</i>	+	-	<i>R. rattus</i> , 33.3% (1/3)	+	-	-	-	[73]
	<i>B. tribocorum</i>	+	-	<i>Rattus norvegicus</i> , 9.4% (5/53)	+	-	-	-	[73]
	<i>B. elizabethae</i>	+	+	<i>R. tanezumi</i> (<i>R. rattus</i>), 20% (1/5)	+	-	-	-	[42]
	<i>B. queenslandensis</i>	+	+	<i>Rattus norvegicus</i> , 36% (22/61)	+	-	-	-	[42]
	<i>B. phoceensis</i>	+	+	<i>Sciurus murinus</i> , 20% (4/20)	-	+	-	-	[42]
	<i>B. rattimassiliensis</i>	+	+						[42]
	<i>B. tribocorum</i>	+	+						[42]
	<i>B. elizabethae</i>	+	+	<i>R. rattus</i> , 10% (1/10)	+	-	-	-	[74]
	<i>B. grahamii</i>	+	+	<i>Rattus norvegicus</i> , 52.7% (89/169)	+	-	-	-	[74]
	<i>B. phoceensis</i>	+	+	<i>Rattus losea</i> , 66.7% (2/3)	+	-	-	-	[74]
	<i>B. rattimassiliensis</i>	+	+	<i>Sattus murinus</i> , 28.6% (36/126)	-	-	-	-	[74]
	<i>B. tribocorum</i>	+	+						[74]
	<i>B. elizabethae</i>	+	+	<i>Miniopterus schreibersii</i> , 42.9% (6/14)	-	-	+	-	[47]
	<i>B. grahamii</i>	+	+	<i>Rattus coxinga niviventer</i> , 100% (1/1)	+	-	-	-	[47]
	<i>B. rattimassiliensis</i>	+	+	<i>Crocidura attenuata tanakae</i> , 77.8% (7/9)	-	+	-	-	[47]
	<i>B. tribocorum</i>	+	+						[47]
	unknown species	+	+						[47]
Western	<i>B. grahamii</i>	+	-	<i>Myodes rutilus</i> , 14.1% (11/78)	+	-	-	-	[85]
Siberia	<i>B. taylorii</i>	+	-	<i>Myodes glareolus</i> , 16% (4/25)	+	-	-	-	[85]
Russia	<i>Bartonella</i> unknown species	+	-						[85]

genera *Apodemus*, *Eothenomys*, *Mus* and *Myodes* have also been shown to host *Bartonella*. A high diversity of *Bartonella* species was observed from rodents in this region including *B. elizabethae*, *B. grahamii*, *B. queenslandensis*, *B. phoceensis*, *B. rattimassiliensis*, *B. taylorii*, *B. tribocorum* and unknown species.

In Southeast Asia, *Bartonella* species have been investigated from rodents in Indonesia, Lao PDR, Cambodia and Thailand^[36–40] (Table 3). Again, the genus *Rattus* appears to be the more prevalent rodent genus followed by the genera *Bandicota* and *Mus*. In South and West Asia, studies of *Bartonella* in rodents are more limited (Table 4).

Altogether, it appears that *B. coopersplainsensis* is mainly found in *Bandicota* species [*Bandicota indica* (*B. indica*) and *Bandicota savilei* (*B. savilei*)] and *R. tanezumi*; *B. elizabethae* in *Apodemus agrarius* and *Rattus* species [*Rattus rattus* (*R. rattus*) and *R. tanezumi*]; *B. grahamii* in *Rattus norvegicus*; *B. queenslandensis* in *Mus* species [*Mus cookii* (*M. cookii*), *B. phoceensis* in *R. tanezumi*, *B. rattimassiliensis* in *M. cookii* and *R. tanezumi*; *B. rochalimae* in *Mus cervicolor*; *B. silvatica* in *Clethrionomys rufocanus*; *B. taylorii* in *C. rufocanus*; *B. tribocorum* in *Apodemus* species (*A. chevrieri*, *A. latronum*) and in *Rattus* species (*R. tanezumi*, *Rattus norvegicus*); *B. washoensis* in *Pachyuromys duprasi*.

4.2. Shrews

Several shrew species live in the domestic areas or peri-domestic environments such as agricultural areas. Isolation of *Bartonella* from shrews has been carried out in Asia and North America. In Asia, most *Bartonella* infections in shrews have been described in East Asia (Korea and Taiwan) and South Asia (Bangladesh and Nepal)^[41–43]. In Southeast Asia, the prevalence of *Bartonella* in shrews has been reported in *Sattus murinus* in Indonesia^[37] (Table 3). Potential new species of *Bartonella* has been recently observed in *Sattus murinus* in Cambodia^[36].

Finally, it appears that *Sattus murinus* is a main host for *B. elizabethae*, *B. phoceensis*, *B. rattimassiliensis* and *B. tribocorum*, although *Crocidura attenuata* is found infected with high prevalence by *B. rattimassiliensis* and *B. tribocorum*.

4.3. Bats

The knowledge of *Bartonella* in bats is still scarce, with only few studies published concerning bats from Asia^[44]. Studies on bats and their arthropod vectors have

Table 3*Bartonella* infection and host species in Southeast Asia.

Country	<i>Bartonella</i> infection	Diagnostic test		Host species and prevalence of each species	Type of animal				Reference	
		PCR	Culture		Rodent	Shrew	Bat	Exotic pet		
Cambodia	<i>B. coopersplainsensis</i>	+	+	<i>B. savilei</i> , 9.5% (7/74)	+	-	-	-	[36]	
	<i>B. elizabethae</i>	+	+	<i>Berylmys berdmorei</i> , 9.1% (1/11)	+	-	-	-	[36]	
	<i>B. queenslandensis</i>	+	+	<i>Maxomys surifer</i> , 1.1% (1/92)	+	-	-	-	[36]	
	<i>B. rattimassiliensis</i>	+	+	<i>Niviventer fulvescens</i> , 10% (1/10)	+	-	-	-	[36]	
	<i>B. tribocorum</i>	+	+	<i>Rattus argentiventer</i> , 9.5% (4/42)	+	-	-	-	[36]	
	Unknown new species	+	+	<i>Rattus exulans</i> , 3.5% (4/115)	+	-	-	-	[36]	
			+	+	<i>R. tanzumi phylogenetic R3</i> , 24.8%(29/117)	+	-	-	-	[36]
Indonesia	<i>B. phoceensis</i>	+	-	<i>R. tanezumi (R. rattus)</i> , 10.3% (8/78)	+	-	-	-	[37]	
	<i>B. rattimassiliensis</i>	+	-	<i>Rattus norvegicus</i> , 2% (1/49)	+	-	-	-	[37]	
				<i>S. murinus</i> , 4.8% (4/83)	-	+	-	-	[37]	
Laos	<i>B. elizabethae</i>	+	-	<i>R. tanezumi (R. rattus)</i> , 16% (61/381)	+	-	-	-	[38]	
	<i>B. phoceensis</i>	+	-	<i>Rattus exulans</i> , 21.5% (17/79)	+	-	-	-	[38]	
	<i>B. tribocorum</i>	+	-	<i>Cannomys badius</i> , 50% (1/2)	+	-	-	-	[38]	
				<i>Mus cervicolor</i> , 1.4% (1/71)	+	-	-	-	[38]	
				<i>Mus caroli</i> , 33.3% (1/3)	+	-	-	-	[38]	
	<i>B. coopersplainsensis</i>	+	+	<i>B. savilei</i> , 7.1% (1/14)	+	-	-	-	[36]	
	<i>B. elizabethae</i>	+	+	<i>Berylmys berdmorei</i> , 20% (1/5)	+	-	-	-	[36]	
	<i>B. queenslandensis</i>	+	+	<i>Mus caroli</i> , 4.8% (1/23)	+	-	-	-	[36]	
	<i>B. rattimassiliensis</i>	+	+	<i>M. cookii</i> , 30.9% (17/55)	+	-	-	-	[36]	
	<i>B. tribocorum</i>	+	+	<i>Rattus andamanensis</i> , 40% (2/5)	+	-	-	-	[36]	
	Unknown new species	+	+	<i>R. tanezumi</i> , 21.8% (17/78)	+	-	-	-	[36]	
	Thailand	<i>B. queenslandensis</i>	+	-	<i>B. savilei</i> , 6.1% (2/33)	+	-	-	-	[39]
		<i>B. coopersplainsensis</i>	+	-	<i>R. tanezumi (R. rattus)</i> , 8.2% (4/49)	+	-	-	-	[39]
		<i>B. phoceensis</i>	+	-	<i>Rattus exulans</i> , 1.5% (1/69)	+	-	-	-	[39]
		<i>B. rochalimae</i>	+	-	<i>Mus cervicolor</i> , 50% (1/2)	+	-	-	-	[39]
<i>Candidatus Bartonella thailandensis</i>		+	+	<i>Berylmys berdmorei</i> , 25% (5/20)	+	-	-	-	[39]	
<i>B. elizabethae</i>		+	+	<i>Maxomys surifer</i> , 24.3% (8/33)	+	-	-	-	[39]	
<i>B. grahamii</i>		+	+	<i>B. indica</i> , 8.1% (12/147)	+	-	-	-	[40]	
				<i>R. tanezumi (R. rattus)</i> , 12.5% (3/24)	+	-	-	-	[40]	
				<i>Rattus losea</i> , 18% (2/11)	+	-	-	-	[40]	
<i>B. coopersplainsensis</i>		+	+	<i>B. indica</i> , 32.6% (15/46)	+	-	-	-	[86]	
<i>B. elizabethae</i>		+	+	<i>B. savilei</i> , 57.1% (4/7)	+	-	-	-	[86]	
<i>B. phoceensis</i>		+	+	<i>R. tanezumi (R. rattus)</i> , 65.2% (88/135)	+	-	-	-	[86]	
<i>B. rattimassiliensis</i>		+	+	<i>Rattus exulans</i> , 3.2% (3/95)	+	-	-	-	[86]	
<i>B. tribocorum</i>		+	+	<i>Rattus losea</i> , 33.3% (1/3)	+	-	-	-	[86]	
				<i>R. norvegicus</i> , 86.4% (19/22)	+	-	-	-	[86]	
				<i>Rattus argentiventer</i> , 66.7% (2/3)	+	-	-	-	[86]	
				<i>Rattus remotus</i> , 50% (1/2)	+	-	-	-	[86]	
				<i>Mus cervicolor</i> , 42.9% (3/7)	+	-	-	-	[86]	
				<i>Berylmys berdmorei</i> , 100% (1/1)	+	-	-	-	[86]	
<i>B. coopersplainsensis</i>		+	+	<i>B. indica</i> , 7.7% (5/65)	+	-	-	-	[36]	
<i>B. elizabethae</i>		+	+	<i>Berylmys berdmorei</i> , 13.3% (2/15)	+	-	-	-	[36]	
<i>B. phoceensis</i>		+	+	<i>Mus caroli</i> , 8% (2/26)	+	-	-	-	[36]	
<i>B. queenslandensis</i>		+	+	<i>M. cookii</i> , 17.1% (6/35)	+	-	-	-	[36]	
<i>B. rattimassiliensis</i>		+	+	<i>Mus cervicolor</i> , 20.8% (10/48)	+	-	-	-	[36]	
<i>B. tribocorum</i>		+	+	<i>Mus sp.</i> , 75% (3/4)	+	-	-	-	[36]	
				<i>Niviventer fulvescens</i> , 12.5% (1/8)	+	-	-	-	[36]	
				<i>Rattus exulans</i> , 4.2% (6/142)	+	-	-	-	[36]	
			<i>R. tanezumi</i> , 31.6% (12/38)	+	-	-	-	[36]		
			<i>R. tanezumi R3</i> , 20.7% (6/29)	+	-	-	-	[36]		

demonstrated that bats were infected with unknown *Bartonella* species[45]. Another study found new putative species in bats from Kenya[46]. In Asia, reports are even scarcer. *Bartonella* infection in bat has been described in Taiwan[47]. Six blood samples of *Miniopterus schreibersii* were positive with new putative *Bartonella* spp. with high prevalence (42.9%, Table 2).

4.4. Exotic pets

Exotic pets are wildlife that are traded around the world, with important illegal trafficking, and have been (and still being) imported into many countries. The illegal trade without quarantine may cause the dissemination of infectious diseases. However, there are little data on

Table 4

Bartonella infection and host species in South and West Asia.

Country	Bartonella infection	Diagnostic test		Host species	Type of animal				Reference
		PCR	Culture		Rodent	Shrew	Bat	Exotic pet	
Bangladesh	<i>B. elizabethae</i>	+	+	<i>Bandicota bengalensis</i> , 63.2% (48/76)	+	–	–	–	[37]
	<i>B. tribocorum</i>	+	+	<i>R. rattus</i> , 32.3% (32/99)	+	–	–	–	[37]
				<i>Sciurus murinus</i> , 42.9% (6/14)	–	+	–	–	[37]
Nepal	<i>B. elizabethae</i>	+	–	<i>B. bengalensis</i> , 26.3% (10/38)	+	–	–	–	[87]
	<i>B. queenslandensis</i>	+	–	<i>R. rattus brunneusculus</i> , 43.3% (39/90)	+	–	–	–	[87]
	<i>B. phoceensis</i>	+	–	<i>Sciurus murinus</i> , 64.1% (59/92)	–	+	–	–	[87]
	<i>B. rochalimae</i>	+	–						[87]
	<i>B. rattimassiliensis</i>	+	–						[87]
	<i>B. tribocorum</i>	+	–						[87]
Israel	<i>B. elizabethae</i>	+	+	<i>R. rattus</i> , 16% (10/62)	+	–	–	–	[88]
	<i>B. tribocorum</i>	+	+						[88]
	<i>B. elizabethae</i>	+	–	<i>Apodemus cahirinus</i> , 25% (1/4)	+	–	–	–	[89]
	<i>B. tribocorum</i>	+	–	<i>R. rattus</i> , 24% (19/79)	+	–	–	–	[89]

Table 5

Species of Bartonella detected on arthropod vectors from rodents and shrews in Asia.

Country	Bartonella infection	Diagnostic test	Vector species	Type of vector	Host reservoirs	Ref
Afghanistan	<i>B. elizabethae</i>	PCR	<i>Xenopsylla cheopis</i>	flea	<i>Cricetulus migratorius</i>	[90]
	<i>B. dosiae</i>	PCR	<i>Xenopsylla cheopis</i>	flea	<i>Meriones libicus</i> (gerbil)	[90]
	<i>B. koehlerae</i>	PCR	<i>Xenopsylla cheopis</i>	flea	<i>Mus</i> spp.	[90]
	<i>B. quintana</i>	PCR	<i>Xenopsylla cheopis</i>	flea	<i>Rattus</i> spp.	[90]
	<i>B. taylorii</i>	PCR	<i>Xenopsylla cheopis</i>	flea		[90]
China	<i>B. tribocorum</i>	PCR	<i>Xenopsylla cheopis</i>	flea	<i>Rattus tanezumi</i> subsp. <i>flavipectus</i>	[91]
		PCR	<i>Ctenophthalmus lushuiensis</i>	flea	<i>Eothenomys</i> spp.	[91]
Korea	<i>Bartonella</i> spp.	PCR	<i>Haemaphysalis longicornis</i>	tick	Wild rodents	[34]
		PCR	<i>Haemaphysalis flava</i>	tick	Unidentified species	[34]
		PCR	<i>Ixodes nipponensis</i>	tick		[34]
		PCR	<i>Ixodes turdus</i>	tick		[34]
		PCR	<i>Ixodes persulcatus</i>	tick		[34]
		PCR	<i>Ixodes</i> spp.	tick		[34]
Israel	<i>Bartonella</i> sp.	PCR	<i>Ctenophthalmus congeneroides</i>	flea	<i>Apodemus agrarius</i> , <i>Crocidura lasiura</i> , <i>Microtus fortis</i> , <i>Myodes regulus</i> , <i>Micromys minutus</i> , <i>Mus musculus</i>	[92]
		PCR	<i>Stenoponia sidimi</i>	flea		[92]
		PCR	<i>Xenopsylla cheopis</i>	flea	<i>R. rattus</i>	[93]
		PCR	<i>Leptopsylla algira</i>	flea	<i>Mus musculus</i>	[93]
		PCR	<i>Xenopsylla ramesis</i>	flea	<i>Meriones tristrami</i>	[93]
		PCR	<i>Synosternus cleopatrae</i>	flea	<i>Gerbillus andersoni allenbyi</i>	[93]
Taiwan	<i>Bartonella</i> spp.	PCR			<i>Gerbillus pyramidum</i>	[93]
		PCR			<i>Meriones sacramenti</i>	[93]
		PCR	<i>Laelaps</i> spp.	mite	<i>Apodemus cahirinus</i>	[42]
		PCR	<i>Laelaps echianinus</i>	mite	<i>Rattus norvegicus</i>	[42]
		PCR	<i>Xenopsylla cheopis</i>	flea	<i>R. rattus</i>	[42]
		PCR	<i>Polyplax spinulosa</i>	lice	<i>Sciurus murinus</i>	[42]
Thailand	<i>Bartonella</i> BNFrs	PCR	<i>Dermacentor</i> spp.	tick		[42]
		PCR	<i>Ixodes</i> spp.	tick		[42]
		PCR	<i>Nosopsyllus fasciatus</i>	flea	<i>Rattus surifer</i>	[94]
		PCR	<i>Leptotrombidium</i> spp.	mite	<i>Rattus argentiventer</i>	[49]
		PCR	<i>Schoengastia</i> spp.	mite	<i>R. rattus</i>	[49]
		PCR	<i>Blankarttia</i> spp.	mite	<i>B. indica</i>	[49]
	<i>B. tamiae</i>	PCR	<i>Haemaphysalis</i> spp.	tick	<i>B. savilei</i>	[49]
		PCR			<i>Mus cervicolor</i>	[49]
		PCR				[49]

prevalence of infection with *Bartonella* spp. from exotic small mammal pets in Asia (Table 2). Among them, a study reported that several exotic species in Japan were infected with *Bartonella* spp. with quite high prevalence[48].

4.5. Vectors

Several blood-sucking arthropods including fleas, mites,

sandflies and ticks have been reported as potential vectors transmitting *Bartonella* among animals and between animals and humans.

Of the several blood-sucking arthropods, fleas are the key vectors of *Bartonella* infection. Several studies have suggested that fleas, especially *Xenopsylla cheopis*, are potential vectors in rodents (Table 5). Ticks, mites and lice are also potential vectors for *Bartonella* infections in

Table 6Case reports of *Bartonella* infection in Asia.

Country	<i>Bartonella</i> infection	Diagnostic test	Clinical sign	Patient	Ref
India	<i>B. quintana</i>	Culture, serology	endocarditis	endocarditis patients	[95]
Japan	<i>B. henselae</i>	Culture, PCR	endocarditis	77– year–old, male	[96]
	<i>B. quintana</i>	Serology	fever, headache, anorexia, history of body lice and fleas infestation	homeless people, male and female	[97]
Jordan	<i>B. quintana</i>	Culture, PCR	endocarditis	70– year–old, female	[98]
	<i>B. quintana</i>	Culture, PCR	endocarditis	66–year–old, male	[99]
	<i>B. quintana</i>	Serology, PCR	None (Seroprevalence among Jordanian children)	None	[100]
	<i>B. henselae</i>	Serology, PCR		None	[100]
Korea	<i>B. henselae</i>	Gram staining, PCR	fever, enlargement of axillary lymph nodes	8–year–old, girl	[101]
	<i>B. henselae</i>	Serology	cervical lymphadenitis	25– year–old, female	[101]
Nepal	<i>B. elizabethae</i>	Serology	headache, fever, myalgia	ranging from 9 to 70 years, both male and female	[53]
	<i>B. quintana</i>	Serology	headache, fever, myalgia		[53]
	<i>B. tamiae</i>	Serology	headache, fever, myalgia		[53]
	<i>B. vinsonii arupensis</i>	Serology	headache, fever, myalgia		[53]
	<i>B. elizabethae</i>	Serology	febrile patients, fever > 38 °C for < 2 weeks	febrile patients, ≥7 years old	[52]
	<i>B. henselae</i>	Serology	non–febrile patients, (unidentified)	non–febrile patients, ≥14 years old	[52]
	<i>B. quintana</i>	Serology			[52]
	<i>B. henselae</i>	Culture, PCR	fever, pharyngitis, meningitis	febrile patients	[50]
	<i>B. henselae</i>	Culture, PCR	having symptomatic HIV infection	40 year–old	[102]
	<i>B. tamiae</i>	Culture, PCR	fatigue, myalgia, fever, headache	38 year–old	[8]
Thailand	<i>B. tamiae</i>	Culture, PCR	pterygium in each eye	41 year–old	[8]
	<i>B. tamiae</i>	Culture, PCR	fever, fatigue, myalgia, a headache	12 year–old	[8]
	<i>B. vinsonii</i>	Culture, PCR	headache, myalgias, dizziness, and fatigue	11 to 56 year–old	[59]

rodents. Epidemiological surveys of *Bartonella* infection in ticks, mites and lice have been done in Korea, Taiwan and Thailand[16,34,42,49].

6. Zoonotic diseases and case reports in Asia

Bartonella species are important zoonotic bacteria with increasing array of emerging infections in humans and animals. Bartonellosis has been described in Asia. Possible routes of transmission are contacts with infected animals or blood–sucking arthropod via their feces. Four *Bartonella* species including *B. henselae*, *B. quintana*, *B. elizabethae* and *B. tamiae* have been reported as causative agents of several human diseases in Asia (Table 6). However, few cases of human infections with *Bartonella* bacteria of rodent origins have been reported. The first case of *Bartonella* endocarditis in Thailand was found in a 57–year–old male poultry farmer in Khon Kaen province, where the source of pathogen was a cat. *B. tamiae* was isolated from three febrile patients in Thailand[8]. In another report, several farmers were supposed to be exposed to infected rodents at home or fields[50]. Recently, four patients were found infected by *B. vinsonii* in Northern and Northeastern Thailand, with dogs and rats supposed to be reservoirs[51]. Additional serological studies evidenced exposure to rodent *Bartonella* spp. in Thailand but also in Nepal[52,53]. However, to date, only *B. elizabethae* was found in rodent surveys, where the most common rodents were murid from *Apodemus* spp. and *Rattus* spp. (and particularly the commensal *R. rattus* and *R. tanezumii*).

7. Perspectives

The diversity of *Bartonella* species is far from being known as the recent study of Jiyipong *et al.* emphasized on three new putative species[36]. Indeed, Southeast Asia is a hotspot of mammal species[54] and a diversification centre for several rodent families. Two–thirds of living rodent species belongs to the family Muridae[55], which also represents most of the rodents found in Southeast Asia with 35 species[56]. It appears that less than 50% of these murid species have been investigated for the presence of *Bartonella* species (Table 3). The knowledge is even poorer for shrews and bats. This emphasizes the need to increase the screening, detection and characterization of *Bartonella* in small mammals[57].

Since *Bartonella* species may be transmitted to humans by ticks, fleas and lice, surveys of *Bartonella* distribution within arthropods are needed. In particular, the population changes of these vectors of *Bartonella* in relation to the impacts of habitat and land–use changes should be better investigated[57].

Rodents, shrews and bats live in a wide range of habitats that are frequented by humans[58], which warrants further investigations on the transmission ecology of *Bartonella* in order to improve prevention of *Bartonella* infections. Identification of risky habitats for human transmission is needed to develop a surveillance strategy, which could be done only after the improvement of our knowledge on the diversity of *Bartonella* in small mammals and in their arthropod vectors.

This review by giving up–to–date list of reservoirs will help develop a strategy of reservoirs, vectors and habitats

prone to be sources of outbreaks and/or emerging *Bartonella* infections in humans.

Conflict of interest statement

We declare that we have no conflict of interest.

Acknowledgements

The study was supported by the French ANR CERoPath project (number ANR 07 BDIV 012) (www.ceropath.org), the French ANR BiodivHealthSEA project (number ANR 11 CPEL 002) (www.biodivhealthsea.org), the Infectiopole Sud and the Center of Excellence on Agricultural Biotechnology (Science and Technology Postgraduate Education and Research Development Office, Office of Higher Education Commission, Ministry of Education) (AG-BIO/PERDO-CHE). Part of this research was also funded by the Center of Advanced Studies for Agriculture and Food, Institute for Advanced Studies, Kasetsart University. We thank the editor and a reviewer for helpful comments.

Comments

Background

Bartonella is an important bacteria that can make various infections in different animals specially humans. This bacteria has various hosts and it is transmitted by several vectors. Species of *Bartonella* are Gram-negative intracellular bacteria that infects erythrocytes of various mammalian and non-mammalian animals and mainly transmitted by blood-sucking arthropod vectors.

Research frontiers

This is a review article about *Bartonella* infection in small mammals including rodents, insectivores, bats and exotic small mammal pets and their vectors in Asia. This article dicusses the present species of *Bartonella* in animals cited.

Related reports

This is a review article and it is based on various reports that are already reported. Various articles mentioned role of different species of *Bartonella* in making infections in animals (mammals and non-mammals). *Bartonella* in different animals have been recognized.

Innovations and breakthroughs

This review article investigates various species of *Bartonella* in animals that already a little have been studied and can offer good information about the role of *Bartonella* at causing infection in animals cited.

Applications

This review article surveys prevalence of various species of *Bartonella* in different areas of Asia. Informations offered in this article can be useful for more studies in future.

Peer review

This is a good research about the presence of various species of *Bartonella* and its role at causing different infections. The consideration of authors to various reservoirs of this bacteria in animals that a little have been already surveyed is interesting. Diagnostic tests referred in this article are very useful.

References

- [1] Boulouis HJ, Chang CC, Henn JB, Kasten RW, Chomel BB. Factors associated with the rapid emergence of zoonotic *Bartonella* infections. *Vet Res* 2005; **36**: 383–410.
- [2] Saisongkorh W, Rolain JM, Suputtamongkol Y, Raoult D. Emerging *Bartonella* in humans and animals in Asia and Australia. *J Med Assoc Thai* 2009; **92**: 707–731.
- [3] Angelakis E, Billeter SA, Breitschwerdt EB, Chomel BB, Raoult D. Potential for tick-borne bartonellosis. *Emerg Infect Dis* 2010; **16**: 385–391.
- [4] Raoult D, Roblot F, Rolain JM, Besnier JM, Loulergue J, Bastides F, et al. First isolation of *Bartonella alsatica* from a valve of a patient with endocarditis. *J Clin Microbiol* 2006; **44**: 278–279.
- [5] Avidor B, Graidy M, Efrat G, Leibowitz C, Shapira G, Schattner A, et al. *Bartonella koehlerae*, a new cat-associated agent of culture-negative human endocarditis. *J Clin Microbiol* 2004; **42**: 3462–3468.
- [6] Roux V, Eykyn SJ, Wyllie S, Raoult D. *Bartonella vinsonii* subsp. *berkhoffii* as an agent of afebrile blood culture-negative endocarditis in a human. *J Clin Microbiol* 2000; **38**: 1698–1700.
- [7] Olarte L, Ampofo K, Thorell EA, Sanderson S, Doby E, Pavia AT, et al. *Bartonella vinsonii* endocarditis in an adolescent with congenital heart disease. *Pediatr Infect Dis J* 2012; **31**: 531–534.
- [8] Kosoy M, Morway C, Sheff KW, Bai Y, Colborn J, Chalcraft L, et al. *Bartonella tamiae* sp. nov., a newly recognized pathogen isolated from three human patients from Thailand. *J Clin Microbiol* 2008; **46**: 772–775.
- [9] Ereemeeva ME, Gerns HL, Lydy SL, Goo JS, Ryan ET, Mathew SS, et al. Bacteremia, fever, and splenomegaly caused by a newly recognized *Bartonella* species. *N Engl J Med* 2007; **356**: 2381–2387.
- [10] Kosoy M, Murray M, Gilmore RD Jr, Bai Y, Gage KL. *Bartonella* strains from ground squirrels are identical to *Bartonella washoensis* isolated from a human patient. *J Clin Microbiol* 2003; **41**: 645–650.
- [11] Lin EY, Tsigrelis C, Baddour LM, Lepidi H, Rolain JM, Patel R, et al. *Candidatus Bartonella mayotimonensis* and endocarditis. *Emerg Infect Dis* 2010; **16**: 500–503.
- [12] Dehio C. *Bartonella* interactions with endothelial cells and erythrocytes. *Trends Microbiol* 2001; **9**: 279–285.

- [13] Dehio M, Quebatte M, Foser S, Certa U. The transcriptional response of human endothelial cells to infection with *Bartonella henselae* is dominated by genes controlling innate immune responses, cell cycle, and vascular remodelling. *Thromb Haemost* 2005; **94**: 347–361.
- [14] Valentine KH, Harms CA, Cadenas MB, Birkenheuer AJ, Marr HS, Braun–McNeill J, et al. *Bartonella* DNA in loggerhead sea turtles. *Emerg Infect Dis* 2007; **13**: 949–950.
- [15] Tsai YL, Chang CC, Chuang ST, Chomel BB. *Bartonella* species and their ectoparasites: selective host adaptation or strain selection between the vector and the mammalian host? *Comp Immunol Microbiol Infect Dis* 2011; **34**: 299–314.
- [16] Billeter SA, Levy MG, Chomel BB, Breitschwerdt EB. Vector transmission of *Bartonella* species with emphasis on the potential for tick transmission. *Med Vet Entomol* 2008; **22**: 1–15.
- [17] Jones KE, Patel NG, Levy MA, Storeygard A, Balk D, Gittleman JL, et al. Global trends in emerging infectious diseases. *Nature* 2008; **451**: 990–993
- [18] Coker RJ, Hunter BM, Rudge JW, Liverani M, Hanvoravongchai P. Emerging infectious diseases in Southeast Asia: regional challenges to control. *Lancet* 2011; **377**: 599–609.
- [19] Chomel BB, Boulouis HJ, Maruyama S, Breitschwerdt EB. *Bartonella* spp. in pets and effect on human health. *Emerg Infect Dis* 2006; **12**: 389–394.
- [20] Breitschwerdt EB, Kordick DL. *Bartonella* infection in animals: carriership, reservoir potential, pathogenicity, and zoonotic potential for human infection. *Clin Microbiol Rev* 2000; **13**: 428–438.
- [21] La Scola B, Zeaiter Z, Khamis A, Raoult D. Gene–sequence–based criteria for species definition in bacteriology: the *Bartonella* paradigm. *Trends Microbiol* 2003; **11**: 318–321.
- [22] Benson LA, Kar S, McLaughlin G, Ihler GM. Entry of *Bartonella bacilliformis* into erythrocytes. *Infect Immun* 1986; **54**: 347–353.
- [23] Rolain JM, Chanet V, Laurichesse H, Lepidi H, Beytout J, Raoult D. Cat scratch disease with lymphadenitis, vertebral osteomyelitis, and spleen abscesses. *Ann N Y Acad Sci* 2003; **990**: 397–403.
- [24] Alsmark CM, Frank AC, Karlberg EO, Legault BA, Ardell DH, Canbäck B, et al. The louse–borne human pathogen *Bartonella quintana* is a genomic derivative of the zoonotic agent *Bartonella henselae*. *Proc Natl Acad Sci USA* 2004; **101**: 9716–9721.
- [25] Engel P, Salzburger W, Liesch M, Chang CC, Maruyama S, Lanz C, et al. Parallel evolution of a type IV secretion system in radiating lineages of the host–restricted bacterial pathogen *Bartonella*. *PLoS Genet* 2011; **7**: e1001296.
- [26] Berglund EC, Frank AC, Calteau A, Pettersson OV, Granberg F, Eriksson AS, et al. Run–off replication of host–adaptability genes is associated with gene transfer agents in the genome of mouse–infecting *Bartonella grahamii*. *PLoS Genet* 2009; **5**: e1000546.
- [27] Saenz HL, Engel P, Stoeckli MC, Lanz C, Raddatz G, Vayssier–Taussat M, et al. Genomic analysis of *Bartonella* identifies type IV secretion systems as host adaptability factors. *Nat Genet* 2007; **39**: 1469–1476.
- [28] Merhej V, Croce O, Robert C, Rolain JM, Raoult D. Genome sequence of *Bartonella rattimassiliensis*, a bacterium isolated from European *Rattus norvegicus*. *J Bacteriol* 2012; **194**: 7013.
- [29] Merhej V, Croce O, Robert C, Rolain JM, Raoult D. Genome sequence of *Bartonella rattaaustraliani*, a bacterium isolated from an Australian rat. *J Bacteriol* 2012; **194**: 7012.
- [30] Rolain JM, Vayssier–Taussat M, Gimenez G, Robert C, Fournier PE, Raoult D. Genome sequence of *Bartonella birtlesii*, a bacterium isolated from small rodents of the genus *Apodemus*. *J Bacteriol* 2012; **194**: 4779.
- [31] Li H, Tong YG, Huang Y, Bai JY, Yang H, Liu W, et al. Complete genome sequence of *Bartonella quintana*, a bacterium isolated from rhesus macaques. *J Bacteriol* 2012; **194**: 6347.
- [32] Chenoweth MR, Somerville GA, Krause DC, O’Reilly KL, Gherardini FC. Growth characteristics of *Bartonella henselae* in a novel liquid medium: primary isolation, growth–phase–dependent phage induction, and metabolic studies. *Appl Environ Microbiol* 2004; **70**: 656–663.
- [33] Kosoy M, Hayman DT, Chan KS. *Bartonella* bacteria in nature: where does population variability end and a species start? *Infect Genet Evol* 2012; **12**: 894–904.
- [34] Chae JS, Yu DH, Shringi S, Klein TA, Kim HC, Chong ST, et al. Microbial pathogens in ticks, rodents and a shrew in northern Gyeonggi–do near the DMZ, Korea. *J Vet Sci* 2008; **9**: 285–293.
- [35] Bai Y, Kosoy MY, Maupin GO, Tsuchiya KR, Gage KL. Genetic and ecologic characteristics of *Bartonella* communities in rodents in southern China. *Am J Trop Med Hyg* 2002; **66**: 622–627.
- [36] Jiyipong T, Jittapalapong S, Morand S, Raoult D, Rolain JM. Prevalence and genetic diversity of *Bartonella* spp. in small mammals from Southeastern Asia. *Appl Environ Microbiol* 2012; **78**: 8463–8466.
- [37] Winoto IL, Goethert H, Ibrahim IN, Yuniherlina I, Stoops C, Susanti I, et al. *Bartonella* species in rodents and shrews in the greater Jakarta area. *Southeast Asian J Trop Med Public Health* 2005; **36**: 1523–1529.
- [38] Angelakis E, Khamphoukeo K, Grice D, Newton PN, Roux V, Aplin K, et al. Molecular detection of *Bartonella* species in rodents from the Lao PDR. *Clin Microbiol Infect* 2009; **15**: 95–97.
- [39] Saisongkroh W, Wootta W, Sawanpanyalert P, Raoult D, Rolain JM. “*Candidatus* *Bartonella thailandensis*”: a new genotype of *Bartonella* identified from rodents. *Vet Microbiol* 2009; **139**: 197–201.
- [40] Castle KT, Kosoy M, Lerdthusnee K, Phelan L, Bai Y, Gage KL, et al. Prevalence and diversity of *Bartonella* in rodents of northern Thailand: a comparison with *Bartonella* in rodents from southern China. *Am J Trop Med Hyg* 2004; **70**: 429–433.
- [41] Bai Y, Montgomery SP, Sheff KW, Chowdhury MA, Breiman RF, Kabeya H, et al. *Bartonella* strains in small mammals from Dhaka, Bangladesh, related to *Bartonella* in America and Europe. *Am J Trop Med Hyg* 2007; **77**: 567–570.
- [42] Tsai YL, Chuang ST, Chang CC, Kass PH, Chomel BB. *Bartonella* species in small mammals and their ectoparasites in Taiwan. *Am J Trop Med Hyg* 2010; **83**: 917–923.
- [43] Kim CM, Kim JY, Yi YH, Lee MJ, Cho MR, Shah DH, et al. Detection of *Bartonella* species from ticks, mites and small mammals in Korea. *J Vet Sci* 2005; **6**: 327–334.
- [44] Bai Y, Kosoy M, Recuenco S, Alvarez D, Moran D, Turmelle A, et al. *Bartonella* spp. in bats, Guatemala. *Emerg Infect Dis* 2011; **17**:

- 1269–1272.
- [45] Bai Y, Recuenco S, Turmelle A, Osikowicz LM, Gómez J, Rupprecht C, et al. Prevalence and diversity of *Bartonella* spp. in bats in Peru. *Am J Trop Med Hyg* 2012; **87**: 518–523.
- [46] Kosoy M, Bai Y, Lynch T, Kuzmin IV, Niezgodna M, Franka R, et al. *Bartonella* spp. in bats, Kenya. *Emerg Infect Dis* 2010; **16**: 1875–1881.
- [47] Lin JW, Hsu YM, Chomel BB, Lin LK, Pei JC, Wu SH, et al. Identification of novel *Bartonella* spp. in bats and evidence of Asian gray shrew as a new potential reservoir of *Bartonella*. *Vet Microbiol* 2012; **156**: 119–126.
- [48] Inoue K, Maruyama S, Kabeya H, Hagiya K, Izumi Y, Une Y, et al. Exotic small mammals as potential reservoirs of zoonotic *Bartonella* spp. *Emerg Infect Dis* 2009; doi: 10.3201/eid1504.081223.
- [49] Kabeya H, Colborn JM, Bai Y, Lerdthusnee K, Richardson JH, Maruyama S, et al. Detection of *Bartonella tamiae* DNA in ectoparasites from rodents in Thailand and their sequence similarity with bacterial cultures from Thai patients. *Vector Borne Zoonotic Dis* 2010; **10**: 429–434.
- [50] Kosoy M, Bai Y, Sheff K, Morway C, Baggett H, Maloney SA, et al. Identification of *Bartonella* infections in febrile human patients from Thailand and their potential animal reservoirs. *Am J Trop Med Hyg* 2010; **82**: 1140–1145.
- [51] Bai Y, Kosoy MY, Diaz MH, Winchell J, Baggett H, Maloney SA, et al. *Bartonella vinsonii* subsp. *arupensis* in humans, Thailand. *Emerg Infect Dis* 2012; doi: 10.3201/eid1806.111750.
- [52] Bhengsi S, Baggett HC, Peruski LF, Morway C, Bai Y, Fisk TL, et al. *Bartonella* seroprevalence in rural Thailand. *Southeast Asian J Trop Med Public Health* 2011; **42**: 687–692.
- [53] Myint KS, Gibbons RV, Iverson J, Shrestha SK, Pavlin JA, Mongkolsirichaikul D, et al. Serological response to *Bartonella* species in febrile patients from Nepal. *Trans R Soc Trop Med Hyg* 2011; **105**: 740–742.
- [54] Schipper J, Chanson JS, Chiozza F, Cox NA, Hoffmann M, Katariya V, et al. The status of the world's land and marine mammals: diversity, threat, and knowledge. *Science* 2008; **322**: 225–230.
- [55] Wilson DE, Reeder DA. *Mammal species of the world: a taxonomic and geographic reference*. 3rd ed. Maryland: Johns Hopkins University Press; 2005.
- [56] Aplin KP, Brown PR, Jacob J, Krebs CJ, Singleton GR. *Field methods for rodent studies in Asia and the Indo-Pacific*. Canberra: ACIAR; 2003.
- [57] Bordes F, Herbretreau V, Dupuy S, Chaval Y, Tran A, Morand S. The diversity of microparasites of rodents: a comparative analysis that helps in identifying rodent-borne rich habitats in Southeast Asia. *Infect Ecol Epidemiol* 2013; doi: 10.3402/iee.v3i0.20178.
- [58] Boulouis HJ, Chang CC, Henn JB, Kssten RW, Chomel BB. Factors associated with the rapid emergence of zoonotic *Bartonella* infections. *Vet Res* 2005; **36**: 383–410.
- [59] Heller R, Kubina M, Mariet P, Riegel P, Delacour G, Dehio C, et al. *Bartonella alsatica* sp. nov., a new *Bartonella* species isolated from the blood of wild rabbits. *Int J Syst Bacteriol* 1999; **49**: 283–288.
- [60] Fournier PE, Taylor C, Rolain JM, Barrassi L, Smith G, Raoult D, et al. *Bartonella australis* sp. nov. from kangaroos, Australia. *Emerg Infect Dis* 2007; **13**: 1961–1963.
- [61] Birtles RJ, Harrison TG, Fry NK, Saunders NA, Taylor AG. Taxonomic considerations of *Bartonella bacilliformis* based on phylogenetic and phenotypic characteristics. *FEMS Microbiol Lett* 1991; **67**: 187–191.
- [62] Bermond D, Heller R, Barrat F, Delacour G, Dehio C, Alliot A, et al. *Bartonella birtlesii* sp. nov., isolated from small mammals (*Apodemus* spp.). *Int J Syst Evol Microbiol* 2000; **50**: 1973–1979.
- [63] Bermond D, Boulouis HJ, Heller R, Van Laere G, Monteil H, Chomel BB, et al. *Bartonella bovis* Bermond et al. sp. nov. and *Bartonella capreoli* sp. nov., isolated from European ruminants. *Int J Syst Evol Microbiol* 2002; **52**: 383–390.
- [64] Maillard R, Riegel P, Barrat F, Bouillin C, Thibault D, Gandoin C, et al. *Bartonella chomelii* sp. nov., isolated from French domestic cattle (*Bos taurus*). *Int J Syst Evol Microbiol* 2004; **54**: 215–220.
- [65] Lawson PA, Clarridge JE, Collins MD. Description of *Bartonella clarridgeiae* sp. nov. isolated from the cat of a patient with *Bartonella henselae* septicemia. *Med Microbiol Lett* 1996; **5**: 64–73.
- [66] Gundi VA, Taylor C, Raoult D, La Scola B. *Bartonella rattaaustraliani* sp. nov., *Bartonella queenslandensis* sp. nov. and *Bartonella cooperplainsensis* sp. nov., identified in Australian rats. *Int J Syst Evol Microbiol* 2009; **59**: 2956–2961.
- [67] Birtles RJ, Harrison TG, Saunders NA, Molyneux DH. Proposals to unify the genera *Grahamella* and *Bartonella*, with descriptions of *Bartonella talpae* comb. nov., *Bartonella peromysci* comb. nov., and three new species, *Bartonella grahamii* sp. nov., *Bartonella taylorii* sp. nov., and *Bartonella doshiae* sp. nov. *Int J Syst Bacteriol* 1995; **45**: 1–8.
- [68] Daly JS, Worthington MG, Brenner DJ, Moss CW, Hollis DG, Weyant RS, et al. *Rochalimaea elizabethae* sp. nov. isolated from a patient with endocarditis. *J Clin Microbiol* 1993; **31**: 872–881.
- [69] Regnery RL, Anderson BE, Clarridge JE 3rd, Rodriguez-Barradas MC, Jones DC, Carr JH. Characterization of a novel *Rochalimaea* species, *R. henselae* sp. nov., isolated from blood of a febrile, human immunodeficiency virus-positive patient. *J Clin Microbiol* 1992; **30**: 265–274.
- [70] Inoue K, Kabeya H, Shiratori H, Ueda K, Kosoy MY, Chomel BB, et al. *Bartonella japonica* sp. nov. and *Bartonella silvatica* sp. nov., isolated from *Apodemus* mice. *Int J Syst Evol Microbiol* 2010; **60**: 759–763.
- [71] Droz S, Chi B, Horn E, Steigerwalt AG, Whitney AM, Brenner DJ. *Bartonella koehlerae* sp. nov., isolated from cats. *J Clin Microbiol* 1999; **37**: 1117–1122.
- [72] Chomel BB, McMillan-Cole AC, Kasten RW, Stuckey MJ, Sato S, Maruyama S, et al. Candidatus *Bartonella merieuxii*, a potential new zoonotic *Bartonella* species in canids from Iraq. *PLoS Negl Trop Dis* 2012; **6**: e1843.
- [73] Gundi VA, Davoust B, Khamis A, Boni M, Raoult D, La Scola B. Isolation of *Bartonella rattimassiliensis* sp. nov. and *Bartonella phoceensis* sp. nov. from European *Rattus norvegicus*. *J Clin Microbiol* 2004; **42**: 3816–3818.
- [74] Brenner DJ, O'Connor SP, Winkler HH, Steigerwalt AG. Proposals to unify the genera *Bartonella* and *Rochalimaea*, with descriptions of *Bartonella quintana* comb. nov., *Bartonella vinsonii* comb. nov., *Bartonella henselae* comb. nov. and *Bartonella elizabethae* comb. nov., and to remove the family

- Bartonellaceae* from the order Rickettsiales. *Int J Syst Bacteriol* 1993; **43**: 777–786.
- [75] Dehio C, Lanz C, Pohl R, Behrens P, Bermond D, Piémont Y, et al. *Bartonella schoenbuchii* sp. nov., isolated from the blood of wild roe deer. *Int J Syst Evol Microbiol* 2001; **51**: 1557–1565.
- [76] Heller R, Riegel P, Hansmann Y, Delacour G, Bermond D, Dehio C, et al. *Bartonella tribocorum* sp. nov., a new *Bartonella* species isolated from the blood of wild rats. *Int J Syst Bacteriol* 1998; **48**: 1333–1339.
- [77] Welch DF, Carroll KC, Hofmeister EK, Persing DH, Robison DA, Steigerwalt AG, et al. Isolation of a new subspecies, *Bartonella vinsonii* subsp. arupensis, from a cattle rancher: identity with isolates found in conjunction with *Borrelia burgdorferi* and *Babesia microti* among naturally infected mice. *J Clin Microbiol* 1999; **37**: 2598–2601.
- [78] Hofmeister EK, Kolbert CP, Abdulkarim AS, Magera JM, Hopkins MK, Uhl JR, et al. Cosegregation of a novel *Bartonella* species with *Borrelia burgdorferi* and *Babesia microti* in *Peromyscus leucopus*. *J Infect Dis* 1998; **177**: 409–416.
- [79] Breitschwerdt EB, Kordick DL, Malarkey DE, Keene B, Hadfield TL, Wilson K. Endocarditis in a dog due to infection with a novel *Bartonella* subspecies. *J Clin Microbiol* 1995; **33**: 154–160.
- [80] Kordick DL, Swaminathan B, Greene CE, Wilson KH, Whitney AM, O'Connor S, et al. *Bartonella vinsonii* subsp. *berkhoffii* subsp. nov., isolated from dogs; *Bartonella vinsonii* subsp. *vinsonii*; and emended description of *Bartonella vinsonii*. *Int J Syst Bacteriol* 1996; **46**: 704–709.
- [81] Liu Q, Sun J, Lu L, Fu G, Ding G, Song X, et al. Detection of *Bartonella* species in small mammals from Zhejiang Province, China. *J Wildl Dis* 2010; **46**: 179–185.
- [82] Inoue K, Maruyama S, Kabeya H, Yamada N, Ohashi N, Sato Y, et al. Prevalence and genetic diversity of *Bartonella* species isolated from wild rodents in Japan. *Appl Environ Microbiol* 2008; **74**: 5086–5092.
- [83] Lin JW, Chen CY, Chen WC, Chomel BB, Chang CC. Isolation of *Bartonella* species from rodents in Taiwan including a strain closely related to '*Bartonella rochalimae*' from *Rattus norvegicus*. *J Med Microbiol* 2008; **57**: 1496–1501.
- [84] Hsieh JW, Tung KC, Chen WC, Lin JW, Chien LJ, Hsu YM, et al. Epidemiology of *Bartonella* infection in rodents and shrews in Taiwan. *Zoonoses Public Health* 2010; **57**: 439–446.
- [85] Samoylenko I, Raoult D, Malkova M, Tancev A, Yakimenko V, Rudakov N, et al. Detection of alpha-proteobacteria in rodents in a steppe-forest zone of Western Siberia. *Clin Microbiol Infect* 2009; **15**: 127–129.
- [86] Bai Y, Kosoy MY, Lerdthusnee K, Peruski LF, Richardson JH. Prevalence and genetic heterogeneity of *Bartonella* strains cultured from rodents from 17 provinces in Thailand. *Am J Trop Med Hyg* 2009; **81**: 811–816.
- [87] Gundi VA, Kosoy MY, Myint KS, Shrestha SK, Shrestha MP, Pavlin JA, et al. Prevalence and genetic diversity of *Bartonella* species detected in different tissues of small mammals in Nepal. *Appl Environ Microbiol* 2010; **76**: 8247–8254.
- [88] Harrus S, Bar-Gal GK, Golan A, Elazari-Volcani R, Kosoy MY, Morick D, et al. Isolation and genetic characterization of a *Bartonella* strain closely related to *Bartonella tribocorum* and *Bartonella elizabethae* in Israeli commensal rats. *Am J Trop Med Hyg* 2009; **81**: 55–58.
- [89] Morick D, Baneth G, Avidor B, Kosoy MY, Mumcuoglu KY, Mintz D, et al. Detection of *Bartonella* spp. in wild rodents in Israel using HRM real-time PCR. *Vet Microbiol* 2009; **139**: 293–297.
- [90] Marie JL, Fournier PE, Rolain JM, Briolant S, Davoust B, Raoult D. Molecular detection of *Bartonella quintana*, *B. elizabethae*, *B. koehlerae*, *B. doshiae*, *B. taylorii*, and *Rickettsia felis* in rodent fleas collected in Kabul, Afghanistan. *Am J Trop Med Hyg* 2006; **74**: 436–439.
- [91] Li DM, Liu QY, Yu DZ, Zhang JZ, Gong ZD, Song XP. Phylogenetic analysis of *Bartonella* detected in rodent fleas in Yunnan, China. *J Wildl Dis* 2007; **43**: 609–617.
- [92] Kim BJ, Kim SJ, Kang JG, Ko S, Won S, Kim H, et al. First report for the seasonal and annual prevalence of flea-borne *Bartonella* from rodents and soricomorphs in the Republic of Korea. *Vector Borne Zoonotic Dis* 2013; **13**: 457–467.
- [93] Morick D, Krasnov BR, Khokhlova IS, Shenbrot GI, Kosoy MY, Harrus S. *Bartonella* genotypes in fleas (Insecta: Siphonaptera) collected from rodents in the Negev desert, Israel. *Appl Environ Microbiol* 2010; **76**: 6864–6869.
- [94] Parola P, Sanogo OY, Lerdthusnee K, Zeaiter Z, Chauvancy G, Gonzalez JP, et al. Identification of *Rickettsia* spp. and *Bartonella* spp. in from the Thai-Myanmar border. *Ann N Y Acad Sci* 2003; **990**: 173–181.
- [95] Balakrishnan N, Menon T, Fournier PE, Raoult D. *Bartonella quintana* and *Coxiella burnetii* as causes of endocarditis, India. *Emerg Infect Dis* 2008; **14**: 1168–1169.
- [96] Tsuneoka H, Yanagihara M, Otani S, Katayama Y, Fujinami H, Nagafuji H, et al. A first Japanese case of *Bartonella henselae*-induced endocarditis diagnosed by prolonged culture of a specimen from the excised valve. *Diagn Microbiol Infect Dis* 2010; **68**: 174–176.
- [97] Seki N, Sasaki T, Sawabe K, Sasaki T, Matsuoka M, Arakawa Y, et al. Epidemiological studies on *Bartonella quintana* infections among homeless people in Tokyo, Japan. *Jpn J Infect Dis* 2006; **59**: 31–35.
- [98] Yamada Y, Ohkusu K, Yanagihara M, Tsuneoka H, Ezaki T, Tsuboi J, et al. Prosthetic valve endocarditis caused by *Bartonella quintana* in a patient during immunosuppressive therapies for collagen vascular diseases. *Diagn Microbiol Infect Dis* 2011; **70**: 395–398.
- [99] Yoda M, Hata M, Sezai A, Unosawa S, Furukawa N, Minami K. First report of *Bartonella quintana* endocarditis in Japan. *Circ J* 2008; **72**: 1022–1024.
- [100] Al-Majali AM, Al-Qudah KM. Seroprevalence of *Bartonella henselae* and *Bartonella quintana* infections in children from Central and Northern Jordan. *Saudi Med J* 2004; **25**: 1664–1669.
- [101] Yoon HJ, Lee WC, Choi YS, Cho S, Song YG, Choi JY, et al. Cervical lymphadenitis in a patient coinfecting with *Toxoplasma gondii* and *Bartonella henselae*. *Vector Borne Zoonotic Dis* 2010; **10**: 415–419.
- [102] Paitoonpong L, Chitsomkasem A, Chantrakoopungool S, Kanjanahareutai S, Tribuddharat C, Srifuengfung S. *Bartonella henselae*: first reported isolate in a human in Thailand. *Southeast Asian J Trop Med Public Health* 2008; **39**: 123–129.