

Accepted Manuscript

A numerical method for solving singular fractional Lane-Emden type equations

A. Kazemi Nasab, Z. Pashazadeh Atabakan, A.I. Ismail, Rabha W. Ibrahim

PII: S1018-3647(16)30349-4

DOI: <http://dx.doi.org/10.1016/j.jksus.2016.10.001>

Reference: JKUS 413

To appear in: *Journal of King Saud University - Science*

Received Date: 12 July 2016

Accepted Date: 2 October 2016



Please cite this article as: A. Kazemi Nasab, Z. Pashazadeh Atabakan, A.I. Ismail, R.W. Ibrahim, A numerical method for solving singular fractional Lane-Emden type equations, *Journal of King Saud University - Science* (2016), doi: <http://dx.doi.org/10.1016/j.jksus.2016.10.001>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

ACCEPTED MANUSCRIPT

A numerical method for solving singular fractional Lane-Emden type equations

A. Kazemi Nasab^{a,1,*}, Z. Pashazadeh Atabakan^a, A. I. Ismail^a, Rabha W. Ibrahim^b

^a*School of Mathematical Sciences, Universiti Sains Malaysia, 11800 Penang, Malaysia*

^b*Faculty of Computer Science and Information Technology, University Malaya, 50603, Malaysia*

Abstract

In this paper, a hybrid numerical method combining Chebyshev wavelets and a finite difference approach is developed to obtain solutions of singular fractional Lane-Emden type equations. The properties of the Chebyshev wavelets and finite difference approaches are used to convert the problem under consideration into a system of algebraic equations which can be conveniently solved by suitable algorithms. A comparison of results obtained using the present strategy and those reported using other techniques seems to indicate the precision and computational effectiveness of the proposed hybrid method.

Keywords: Chebyshev wavelets, finite difference method, Fractional Lane-Emden, Singular problems, Initial and boundary value problems.

1. Introduction

Certain phenomena in physics and engineering sciences can best be mathematically modelled using differential equations. The Lane-Emden equation is an ordinary differential equation which arises in mathematical physics. In astrophysics, the Lane-Emden equation is a dimensionless form of Poisson's equation for the gravitational potential of simple models of a star (Momoniat, 2006). Due to the singularity behavior at the origin, it is numerically challenging to solve the Lane-Emden problem, as well as other various linear and nonlinear singular initial value problems in quantum mechanics and astrophysics. This paper deals

*Corresponding author:

Email address: a.kazeminasab@usm.edu.my (A. Kazemi Nasab)

with the numerical solution for the singular fractional Lane-Emden equation using Chebyshev wavelet finite difference method.

A given function can be represented in terms of a set of basis functions localized both in location and scale using wavelets. Advantages of wavelet analysis include the capacity of carrying out local analysis to uncover signal features such as trend, breaking points, and discontinuities which other analysis methods might fail to detect (Misiti et al., 2000). The multiresolution analysis of wavelet transform makes it a powerful tool to represent correctly a variety of functions and operators. Wavelet analysis results in sparse matrices which in turn enables fast computation. Such advantageous aspects of wavelet analysis make it a powerful analytical technique to deal with differential equations in which the solution might become distorted due to singularities or sharp transitions.

There has been much research on wavelet based method for the integer order Lane-Emden equations in recent years. Solving Lane-Emden equations using Chebyshev wavelet operational matrix was proposed in (Pandey et al., 2012). Kaur et al. (2013) obtained the solutions for the generalized Lane-Emden equations using Haar wavelet approximation. More recently, Youssri et al. (2015) used ultraspherical wavelets for solving such equations. Kazemi Nasab et al. (2015) proposed Chebyshev wavelet finite difference method for solving singular nonlinear Lane-Emden equations arising in astrophysics. Laguerre wavelets was employed in (Zhou and Xu, 2016).

Approximate solution methods for fractional differential equations developed over the last decade include homotopy analysis method (Abdulaziz et al., 2008), variational iteration method (Wu and Lee, (2010); Odibat and Momani, 2006), finite difference method for fractional partial differential equations (Zhang, 2009), Adomian decomposition method (Daf-tardar and Jafari, 2007; Momani and Odibat, 2007), fractional differential transform method (Odibat et al., 2008), generalized block pulse operational matrix method (Li and Sun, 2011).

El-Ajou et al. (2010) employed homotopy analysis method to find symbolic approximate solutions for linear and nonlinear differential equations of fractional order. In particular for fractional Lane-Emden equations, the least square method was used in (Mechee and Senu, 2012). Dávila et al. (2014) classified solutions of finite Morse index of the fractional Lane-Emden equation. Two hybrid methods were proposed to solve two-dimensional linear time-fractional partial differential equations in (Jacobs and Harley, 2014). More recently, Modified differential transform was proposed in (Marasi et al., 2015) to obtain numerical solutions of the nonlinear and singular Lane-Emden equations of fractional order. Akgül et al. (2015) suggested the reproducing kernel method to study the fractional order Lane-Emden differential equations. The operational matrices of fractional order integration for the Chebyshev wavelet, Legendre wavelet, and Haar wavelet were introduced in (Li, 2010; Jafari et al., 2011; Li and Zhau, 2010; Yuanlu, 2010) to solve the differential equations of fractional order. Bhrawy et al. (2016) introduced a space-time Legendre spectral tau method for the two-sided space-time Caputo fractional diffusion-wave equation.

Kazemi Nasab et al. (2013) proposed Chebyshev wavelet finite difference method for solving fractional differential equations. It is a hybrid method based on Chebyshev wavelet analysis and a finite difference method which converts a given problem to a system of algebraic equations. Ibrahim (2012) studied the general dual formal of fractional order as follows:

$$D^\beta \left(D^\alpha + \frac{a}{x} \right) u(x) + g(x) = h(x), \quad (1.1)$$

The objective of this study is to investigate the effectiveness of Chebyshev wavelet finite difference method when it is applied to a new generalized dual form of the equation (1.1).

In this paper we consider the singular fractional Lane-Emden equations formulated as

$$D^\alpha u(x) + \frac{L}{x^{\alpha-\beta}} D^\beta u(x) + g(x, u(x)) = h(x), \quad (1.2)$$

subject to the following types of initial or boundary conditions:

$$\begin{aligned}
\text{I} & : u(a) = A_1, \quad u'(a) = B_1, \\
\text{II} & : u(a) = A_2, \quad u(b) = B_2, \\
\text{III} & : u'(a) = A_3, \quad u(b) = B_3, \\
\text{IV} & : u(a) = A_4, \quad u'(b) = B_4, \\
\text{V} & : a_1u(a) + a_2u'(a) = A_5, \quad b_1u(b) + b_2u'(b) = B_5, \quad (1.3)
\end{aligned}$$

where $0 < x \leq 1$, $L \geq 0$, $1 < \alpha \leq 2$, $0 < \beta \leq 1$, A_i and B_i are constants, $g(x, u(x))$ is a continuous real valued function and $h(x)$ is in $C[0, 1]$. We employ Chebyshev wavelet finite difference (CWFD) method for solving different types of fractional Lane-Emden equations.

2. Preliminaries

2.1. Fractional calculus

Fractional calculus is the theory of derivatives and integrals of any arbitrary real or complex order that its high significance in a variety of disciplines, including mathematical, physical and engineering sciences is undeniable. It is indeed a generalized conception of integer-order differentiation and n -fold integration. The main benefit of fractional derivatives in comparison with integer-order one is that they provide us with a superb tool to describe general properties of various materials and process specifically dynamical processes in a fractal medium while standard integer-order cases are fail to do so. The benefits of fractional derivatives is evident in representing physical systems more accurately, in demonstrating mechanical and electrical properties of real materials, and also in the depiction of properties of gasses, fluids and rocks, and in numerous different fields (Hilfer, 2000; Lewandowski and Chorazyczewski, 2010; Mainardi, 2012). These equations incorporate the effects of memory and can account for subdiffusion (due to crowded systems) and superdiffusion (due to active transport processes). Thus, taking into account the memory effects are better described

within the fractional derivatives, the fractional Lane-Emden equations extract hidden aspects for the complex phenomena they describe in various field of applied mathematics, mathematical physics, and astrophysics. An advantage of the fractional case is that it converges to the integer case. Therefore, it has the same benefits (Ibrahim, 2012; Akgül et al., 2015). There have been many books and articles in the area of fractional calculus and its application in astrophysics (Miller and Ross, 1993; Samko et al., 1993; Podlubny, 1999; Saxena et al., 2004; Kilbas et al., 2006; Stanislavsky, 2009; Diethelm, 2010; Chaurasia and pandey, 2010; El-Nabulsi, 2011; El-Nabulsi, 2012; Sharma et al., 2015; El-Nabulsi, 2016a; El-Nabulsi, 2016b).

The calculus of variations is a branch of mathematical analysis dealing with extrema of functionals that is of remarkable significance in a variety of disciplines, including science, engineering, and mathematics (Bliss, 1963; Weinstock, 1974). The problems relating to find minimum of a functional such as, Lagrangian, strain, potential, and total energy, actually occur in engineering and science that give the laws governing the systems behavior (Agrawal, 2002). It has been indicated in many textbooks and monographs that all fundamental equations of modern physics can be obtained from corresponding Lagrangians (Lanczos, 1986; Doughty, 1990; Burgess, 2002; Basdevant, 2007). Some basic equations of mathematical physics such as Bessel, Legendre, Laguerre, Hermite, Chebyshev, Jacobi, hypergeometric and confluent hypergeometric equations, and the Lane-Emden equation admit the Lagrangian description. Musielak (2008) suggested some methods to obtain standard and non-standard Lagrangians and identify classes of equations of motion that admit a Lagrangian description equation. Fractional calculus of variations dates back to year 1996 when Riewe successfully described dissipative systems from potentials represented by fractional derivatives (Riewe, 1996; 1997). The advantage of fractional calculus of variations in comparison with the standard one is its ability to describe dissipative classical and quantum dynamical systems where the standard calculus of variations cannot illustrate them well (El-Nabulsi, 2013). The fractional action-like variational approach is a subclass of the fractional calculus of variations was

introduced in (El-Nabulsi, 2005a, 2005b, 2008). El-Nabulsi, (2013) generalized the fractional action-like variational approach for the case of non-standard power-law Lagrangians which can be used to derive the generalized Lane-Emden-type equations. The fractional Chandrasekhar or Lane-Emden non-linear differential equation was also obtained in (El-Nabulsi, 2011). The interested reader may refer to a number of references to get more information on fractional calculus of variations (Hartley and Lorenzo, 1998; Agarwal, 2006; Baleanu et al., 2009; El-Nabulsi, 2011b; Malinowska and Torres, 2012; Odziejewicz et al. 2012; Odziejewicz et al. 2013; Almeida et al. 2015; El-Nabulsi, 2016c).

There are several definitions of fractional integral and derivatives including Riemann-Liouville, Caputo, Weyl, Hadamard, Marchaud, Riesz, Grunwald-Letnikov, and Erdelyi-Kober (Miller and Ross, 1993; Samko et al., 1993; Hilfer, 2000). The most frequently used definition of fractional derivatives is of Riemann-Liouville and Caputo. None of these definitions has satisfied the whole prerequisite required. There is no need a function to be continuous at the origin and differentiable to have Riemann-Liouville fractional derivative. nevertheless, the derivative of a constant is not zero. It also has certain weaknesses to model real-world phenomena with fractional differential equations. Moreover, if an arbitrary function such as exponential and Mittag-Leffler functions, is zero at the origin, its fractional derivation has a singularity at the origin. Furthermore, the Riemann-Liouville method cannot incorporate the nonzero initial condition at lower limit. These disadvantages reduce the field of application of the Riemann-Liouville fractional derivative. The Caputo derivative is very useful when dealing with real-world problem because, it allows the utilization of initial and boundary conditions involving integer order derivatives, which have clear physical interpretations. in addition, the derivative of a constant is zero. In contrast to the Riemann-Liouville fractional derivative, it is not necessary to define the fractional order initial conditions (Podlubny, 1999) when using Caputo's definition. however, with the Caputo fractional derivative, the function needs to be differentiable which diminishes the field of utilization of it (Atangana and Secer, 2013). In this study, we use the Caputo fractional derivative D^α .

Definition 1

A real function $u(x), x > 0$, is said to be in the space $C_\mu, \mu \in \mathbb{R}$ if there exists a real number $p > \mu$ such that $u(x) = x^p u_1(x)$, where $u_1(x) \in C[0, \infty)$, and it is said to be in the space C_μ^n iff $u^n(x) \in C_\mu, n \in \mathbb{N}$.

Definition 2

The Riemann-Liouville fractional integration of order $\alpha \geq 0$ of a function $u \in C_\mu, \mu \geq -1$ is defined as (Podlubny, 1999)

$$\begin{cases} (I^\alpha u)(x) = 1/\Gamma(\alpha) \int_0^x (x - \tau)^{\alpha-1} u(\tau) d\tau, \\ (I^0 u)(x) = u(x), \end{cases} \quad (2.1)$$

Definition 3

The fractional derivative of $u(x)$, in the Caputo sense is defined as (Podlubny, 1999)

$$(D^\alpha u)(x) = (I^{n-\alpha} u^{(n)})(x) = 1/\Gamma(n - \alpha) \int_0^x (x - \tau)^{n-\alpha-1} u^{(n)}(\tau) d\tau, \quad (2.2)$$

for $n - 1 < \alpha \leq n, n \in \mathbb{N}, x > 0, u \in C_{-1}^n$.

2.2. Properties of Chebyshev wavelets

Wavelets are a collection of functions created from dilation and translation of a single function $\psi \in L^2(\mathbb{R})$ called the mother wavelet. If we continuously vary the dilation parameter \mathbf{r} and the translation parameter \mathbf{s} , we then obtain the following family of continuous wavelets (Daubechies, 1993):

$$\psi_{r,s}(x) = |r|^{-\frac{1}{2}} \psi\left(\frac{x-s}{r}\right), \quad r, s \in \mathbb{R}, r \neq 0. \quad (2.3)$$

For some very special choices of $\psi \in L^2(\mathbb{R})$ and $r = 2^{-a}, s = 2^{-ab}$, the family of functions $\{2^{a/2} \psi(2^a x - b) : a, b \in \mathbb{Z}\}$ constitutes an orthonormal basis for the Hilbert space $L^2(\mathbb{R})$. Chebyshev wavelets in turn are defined as:

$$\psi_{n,m}(t) = \begin{cases} 2^{\frac{k}{2}} p_m T_m(2^k x - 2n + 1), & \frac{n-1}{2^{k-1}} \leq x < \frac{n}{2^{k-1}}, \\ 0, & \text{otherwise} \end{cases} \quad (2.4)$$

where

$$p_m = \begin{cases} \frac{1}{\sqrt{\pi}}, & m = 0, \\ \sqrt{\frac{2}{\pi}}, & m \geq 1 \end{cases} \quad (2.5)$$

$m = 0, 1, \dots, M$ is degree of Chebyshev polynomials of the first kind, $n = 1, \dots, 2^{k-1}$; k any positive integer, and x denotes the time. The Chebyshev wavelets $\psi_{n,m}(x)$ construct an orthonormal basis for $L^2_{w_n}[0, 1]$ with respect to the weight function $w_n(x) = w(2^k x - 2n + 1)$ in which $w(x) = \frac{1}{\sqrt{1-x^2}}$ (Babolian and Fattahzadeh, 2007).

In view of orthonormality of the Chebyshev wavelets, a function $u \in L^2_{w_n}[0, 1)$ may be expanded as

$$u(x) = \sum_{n=1}^{\infty} \sum_{m=0}^{\infty} c_{n,m} \psi_{n,m}(x), \quad (2.6)$$

where $c_{n,m} = (u(x), \psi_{n,m}(x))_{w_n}$, in which (\cdot, \cdot) denotes the inner product in $L^2_{w_n}[0, 1)$. If the series in (2.6) is truncated, we then obtain

$$u(x) \approx \sum_{n=1}^{2^{k-1}} \sum_{m=0}^M c_{n,m} \psi_{n,m}(x) = C^T \Psi(x), \quad (2.7)$$

where C and $\Psi(x)$ are $2^k(M+1) \times 1$ matrices given by

$$\begin{aligned} C &= [c_{1,0}, \dots, c_{1,M}, c_{2,0}, \dots, c_{2,M}, \dots, c_{2^{k-1},1}, \dots, c_{2^{k-1},M}]^T, \\ \Psi(x) &= [\psi_{1,0}, \dots, \psi_{1,M}, \psi_{2,0}, \dots, \psi_{2,M}, \dots, \psi_{2^{k-1},1}, \dots, \psi_{2^{k-1},M}]^T. \end{aligned} \quad (2.8)$$

3. Chebyshev wavelet finite difference method

Clenshaw (1960) suggested employing Chebyshev polynomials as a basis to approximate a function u as

$$(P_M u)(x) \approx \sum_{m=0}^M u_m T_m(x), \quad (3.1)$$

$$u_m = \frac{2}{M} \sum_{k=0}^M u(x_k) T_m(x_k) = \frac{2}{M} \sum_{k=0}^M u(x_k) \cos\left(\frac{mk\pi}{M}\right),$$

where the summation symbol with double primes denotes a sum with both the first and last terms halved. Furthermore, the well known Chebyshev-Gauss-Lobatto interpolated points

x_m are defined as

$$x_m = \cos\left(\frac{m\pi}{M}\right), \quad m = 0, 1, 2, \dots, M. \quad (3.2)$$

Elbarbary and El-Kady (2003) obtained the derivatives of the function u at the points $x_m, m = 0, 1, \dots, M$ as

$$u^{(n)}(x_m) \approx \sum_{j=0}^M d_{m,j}^{(n)} u(x_j), \quad n = 1, 2 \quad (3.3)$$

where

$$\begin{aligned} d_{m,j}^{(1)} &= \frac{4\gamma_j}{M} \sum_{k=1}^M \sum_{\substack{l=0 \\ (k+l) \text{ odd}}}^{k-1} \frac{k\gamma_k}{c_l} T_k(x_j) T_l(x_m), \\ &= \frac{4\gamma_j}{M} \sum_{k=1}^M \sum_{\substack{l=0 \\ (k+l) \text{ odd}}}^{k-1} \frac{k\gamma_k}{c_l} \cos\left(\frac{kj\pi}{M}\right) \cos\left(\frac{lm\pi}{M}\right), \end{aligned} \quad (3.4)$$

$$\begin{aligned} d_{m,j}^{(2)} &= \frac{2\gamma_j}{M} \sum_{k=2}^M \sum_{\substack{l=0 \\ (k+l) \text{ even}}}^{k-2} \frac{k(k^2 - l^2)\gamma_k}{c_l} T_k(x_j) T_l(x_m), \\ &= \frac{2\gamma_j}{M} \sum_{k=2}^M \sum_{\substack{l=0 \\ (k+l) \text{ even}}}^{k-2} \frac{k(k^2 - l^2)\gamma_k}{c_l} \cos\left(\frac{kj\pi}{M}\right) \cos\left(\frac{lm\pi}{M}\right), \end{aligned} \quad (3.5)$$

with $\gamma_0 = \gamma_M = \frac{1}{2}, \gamma_j = 1$ for $j = 1, 2, \dots, M-1$.

Let us define $x_{nm}, n = 1, 2, \dots, 2^{k-1}, m = 0, 1, \dots, M$, as the corresponding Chebyshev-Gauss-Lobatto collocation points at the n th subinterval $\left[\frac{n-1}{2^{k-1}}, \frac{n}{2^{k-1}}\right)$ such that,

$$x_{nm} = \frac{1}{2^k} (x_m + 2n - 1). \quad (3.6)$$

A function u can be approximated in terms of Chebyshev wavelet basis functions as follows

$$(P_M u)(x) \approx \sum_{n=1}^{2^{k-1}} \sum_{m=0}^M c_{nm} \psi_{nm}(x), \quad (3.7)$$

where $c_{nm}, n = 1, 2, \dots, 2^{k-1}, m = 0, 1, \dots, M$, are the expansion coefficients of the function $u(x)$ at the subinterval $\left[\frac{n-1}{2^{k-1}}, \frac{n}{2^{k-1}}\right)$ and $\psi_{n,m}(x), n = 1, 2, \dots, 2^{k-1}, m = 0, 1, \dots, M$, are defined in Eq. (2.4). On account of Eq. (3.1), the coefficients c_{nm} are obtained as

$$c_{nm} = \frac{1}{2^{k/2} p_m} \cdot \frac{2}{M} \sum_{p=0}^M u(x_{np}) \cos\left(\frac{mp\pi}{M}\right) = \frac{1}{2^k p_m^2} \cdot \frac{2}{M} \sum_{p=0}^M u(x_{np}) \psi_{nm}(x_{np}). \quad (3.8)$$

From Eqs. (3.3)-(3.5), the derivatives of the function u at the points $x_{nm}, n = 1, 2, \dots, 2^{k-1}, m = 0, 1, \dots, M$, can be obtained as

$$u^{(r)}(x_{nm}) = \sum_{j=0}^M d_{n,m,j}^{(r)} u(x_{nj}), \quad r = 1, 2 \quad (3.9)$$

where

$$\begin{aligned} d_{n,m,j}^{(1)} &= \frac{4\gamma_j}{M} \sum_{k=1}^M \sum_{\substack{l=0 \\ (k+l) \text{ odd}}}^{k-1} \frac{k\gamma_k}{c_l p_k p_l} \psi_{nk}(x_{nj}) \psi_{nl}(x_{nm}), \\ &= \frac{4\gamma_j}{M} \sum_{k=1}^M \sum_{\substack{l=0 \\ (k+l) \text{ odd}}}^{k-1} \frac{2^k k \gamma_k}{c_l} \cos\left(\frac{kj\pi}{M}\right) \cos\left(\frac{lm\pi}{M}\right), \end{aligned} \quad (3.10)$$

$$\begin{aligned} d_{n,m,j}^{(2)} &= \frac{2\gamma_j}{M} \sum_{k=2}^M \sum_{\substack{l=0 \\ (k+l) \text{ even}}}^{k-2} \frac{2^k k(k^2 - l^2) \gamma_k}{c_l p_k p_l} \psi_{nk}(x_{nj}) \psi_{nl}(x_{nm}), \\ &= \frac{2\gamma_j}{M} \sum_{k=2}^M \sum_{\substack{l=0 \\ (k+l) \text{ even}}}^{k-2} \frac{4^k k(k^2 - l^2) \gamma_k}{c_l} \cos\left(\frac{kj\pi}{M}\right) \cos\left(\frac{lm\pi}{M}\right). \end{aligned}$$

4. Characterization of the proposed method

In this section, the CWFD method is used to solve the fractional Lane-Emden equations introduced in Eqs. (1.2)-(1.3). To do so, the interval $[0, 1]$ is divided into 2^{k-1} subintervals $I_n = [\frac{n-1}{2^{k-1}}, \frac{n}{2^{k-1}}]$, $n = 1, 2, \dots, 2^{k-1}$. It is also assumed shifted Chebyshev-Gauss-Lobatto points be the collocation points on the n th subinterval I_n which are defined as

$$x_{ns} = \frac{1}{2^k} (x_s + 2n - 1), \quad s = 1, 2, \dots, M - l. \quad (4.1)$$

$u(x)$ is approximated using Eq. (3.7) and then collocate Eq. (1.2) at the collocation points $x_{ns}, n = 1, 2, \dots, 2^{k-1}, s = 1, \dots, M - l$ to obtain

$$D^\alpha u(x_{ns}) + \frac{L}{x_{ns}^{\alpha-\beta}} D^\beta u(x_{ns}) + g(x, u(x_{ns})) = h(x_{ns}). \quad (4.2)$$

Moreover, we rewrite $D^\alpha u(x_{ns}), D^\beta u(x_{ns})$ in the Caputo sense by virtue of Eq. (2.2) as

$$\begin{aligned}
 D^\alpha u(x_{ns}) &= \frac{1}{\Gamma(n-\alpha)} \int_0^{x_{ns}} (x_{ns} - \tau)^{n-\alpha-1} u^{(n)}(\tau) d\tau \quad n = [\alpha] + 1, \\
 &= \frac{1}{\Gamma(n-\alpha)} \int_0^{\frac{n-1}{2^{k-1}}} (x_{ns} - \tau)^{n-\alpha-1} u^{(n)}(\tau) d\tau \\
 &+ \frac{1}{\Gamma(n-\alpha)} \int_{\frac{n-1}{2^{k-1}}}^{x_{ns}} (x_{ns} - \tau)^{n-\alpha-1} u^{(n)}(\tau) d\tau. \tag{4.3}
 \end{aligned}$$

In view of the Clenshaw-Curtis quadrature formula (Davis and Rabinowitz, 1984), the first integral can be calculated as

$$\begin{aligned}
 &\frac{1}{\Gamma(n-\alpha)} \int_0^{\frac{n-1}{2^{k-1}}} (x_{ns} - \tau)^{n-\alpha-1} u^{(n)}(\tau) d\tau = \\
 &\frac{1}{\Gamma(n-\alpha)} \sum_{r_1=1}^{n-1} \int_{\frac{r_1-1}{2^{k-1}}}^{\frac{r_1}{2^{k-1}}} (x_{ns} - \tau)^{n-\alpha-1} u^{(n)}(\tau) d\tau = \\
 &\frac{1}{\Gamma(n-\alpha)} \cdot \frac{1}{2^k} \sum_{r_1=1}^{n-1} \sum_{r_2=0}^M w_{r_2} (x_{ns} - x_{r_1 r_2})^{n-\alpha-1} u^{(n)}(x_{r_1 r_2}), \tag{4.4}
 \end{aligned}$$

where

$$w_r = \frac{2}{N\gamma_r} \left[1 - \sum_{k=1}^{[M/2]} \frac{2}{\gamma_{2k}(4k^2 - 1)} \cos \frac{2kr\pi}{M} \right], \quad r = 1, 2, \dots, M-1, \tag{4.5}$$

in which $\gamma_0 = \gamma_M = 2$ and $\gamma_r = 1$, for $r = 1, 2, \dots, M-1$. The second singular integral in Eq. (4.3) is evaluated with the help of integration by parts and Eq. (4.1) as following:

$$\begin{aligned}
 &\frac{1}{\Gamma(n-\alpha)} \int_{\frac{n-1}{2^{k-1}}}^{x_{ns}} (x_{ns} - \tau)^{n-\alpha-1} u^{(n)}(\tau) d\tau = \\
 &\frac{(x_{ns} - \frac{n-1}{2^{k-1}})^{n-\alpha}}{(n-\alpha)\Gamma(n-\alpha)} u^{(n)}\left(\frac{n-1}{2^{k-1}}\right) + \frac{(2^{k-1}x_{ns} - n + 1)}{2^k(n-\alpha)\Gamma(n-\alpha)} \\
 &\sum_{s_1=0}^M (x_{ns} - (x_{ns} - \frac{n-1}{2^{k-1}}))(2^{k-1}x_{ns_1} - n + 1) - \frac{n-1}{2^{k-1}})^{n-\alpha} \\
 &u^{(n+1)}\left(\left(x_{ns} - \frac{n-1}{2^{k-1}}\right)(2^{k-1}x_{ns_1} - n + 1) + \frac{n-1}{2^{k-1}}\right). \tag{4.6}
 \end{aligned}$$

Substitute Eqs. (4.4) and (4.6) into Eq. (4.1) to obtain $2^{k-1}(M-1)$ equations. Furthermore,

replacing conditions in Eq. (1.3) with Eqs. (3.7) and/or (3.9) results in obtaining 2 more equations. Moreover, continuity condition on the approximate solution and its first derivative at the interface between subintervals should be imposed to have other $(2^k - 2)$ equations.

$$u^{(r)}(x_{n+1,M}) = u^{(r)}(x_{n,0}), \quad r = 0, 1 \quad n = 1, 2, \dots, 2^{k-1} - 1. \quad (4.7)$$

From the above procedures carried out, we will have a system for a total of $2^{k-1}(M + 1)$ algebraic equations, which can be solved for the $u(x_{ns})$. As a consequence, we get the solution $u(x)$ of Eqs. (1.2)-(1.3) using Eqs. (3.7)-(3.8).

The convergence of the method has been shown in (Kazemi et. al, 2015).

5. Illustrative examples

In this section, fractional Lane-Emden problems of different types are solved to verify the applicability and accuracy of the proposed method in solving singular fractional Lane-Emden problems.

Example 5.1 As a first example for illustration, consider the following nonlinear Lane-Emden type equation:

$$y''(x) + \frac{2}{x}y'(x) + \sinh(y(x)) = 0, \quad x \geq 0 \quad (5.1)$$

subject to the boundary conditions

$$y(0) = 1, \quad y'(0) = 0.$$

Wazwaz (2001) employed Adomian decomposition method to obtain a series solution as follows

$$y(x) \cong 1 - \frac{(e^2-1)x^2}{12e} + \frac{1}{480} \frac{(e^4-1)x^4}{e^2} - \frac{1}{30240} \frac{(2e^6+3e^2-3e^4-2)x^6}{e^3} + \frac{1}{26127360} \frac{(61e^8-104e^6+104e^2-61)x^8}{e^4}.$$

We apply the introduced method with $M = 10, k = 5$ for solving Eq. (5.1). In Table 1, we compare the $y(x)$ values obtained by the present method and those reported in (Wazwaz,

2001; Parand, 2010). The results are in well agreement with those obtained by Adomian decomposition method and more accurate than those acquired by the introduced method in (Parand, 2010).

Table 1: Comparison of $y(x)$ values in Example 5.1, for the present method, method in (Parand, 2010) and series solution given in (Wazwaz, 2001).

t	<i>Present method</i>	Error	<i>Hermit function collocation method</i>	<i>Adomian decomposition method</i>
0.0	1.0000000000	0.00×10^0	0.00×10^0	1.0000000000
0.1	0.9980428414	4.48×10^{-11}	7.10×10^{-5}	0.9980428414
0.2	0.9921894348	1.22×10^{-11}	8.64×10^{-5}	0.9921894348
0.5	0.9519611092	7.37×10^{-8}	7.65×10^{-5}	0.9519611019
1.0	0.8182429285	8.74×10^{-6}	5.31×10^{-5}	0.8182516669
1.5	0.6254387635	4.53×10^{-4}	4.03×10^{-4}	0.6258916077
2.0	0.4066228875	7.05×10^{-3}	7.02×10^{-3}	0.4136691039

Example 5.2 Consider the following nonlinear fractional spherical isothermal Lane-Emden type equation:

$$D^\alpha u(x) + \frac{2}{x^{\alpha-\beta}} D^\beta u(x) = e^{-u(x)}, \quad 0 < x \leq 1, \quad (5.2)$$

subject to the boundary conditions

$$u(0) = 1, \quad u'(0) = 0. \quad (5.3)$$

We used the technique introduced in (Abdel-Salam and Nouth, 2015) to obtain the exact solution of this problem when $\alpha = \frac{3}{2}$ and $\beta = \frac{3}{4}$ as

$$u(x) \cong 0.2368456765x^{\frac{3}{2}} - 0.02568610429x^3 + 0.005004915254x^{\frac{9}{2}} - 0.001292847099x^6 + 0.0004026307151x^{\frac{15}{2}} - \dots$$

We employ the CWFD method with different values of M and k for solving this problem.

In Table 2, we compare the absolute errors of $u(x)$ at different points. Fig. 1 shows the absolute errors in solutions obtained by the introduced method in this paper with $M = 18$ and $k = 2, 3$

Table 2: Comparison of absolute errors in Example 5.2 for $\alpha = \frac{3}{2}$ and $\beta = \frac{3}{4}$.

x	$M = 7, k = 2$	$M = 10, k = 2$	<i>Exact solution</i>
0.1	1.3×10^{-4}	4.3×10^{-5}	0.007464188801946358826
0.2	1.3×10^{-4}	3.3×10^{-5}	0.020982133267328805960
0.3	9.8×10^{-5}	3.9×10^{-6}	0.038245501398878235380
0.4	6.4×10^{-5}	3.4×10^{-5}	0.058349988262337799440
0.5	2.8×10^{-5}	3.3×10^{-5}	0.080730040537947506919
0.6	6.3×10^{-4}	1.6×10^{-4}	0.104978568300143096020
0.7	1.0×10^{-3}	2.1×10^{-4}	0.130782221081233564320
0.8	1.4×10^{-3}	1.5×10^{-4}	0.157891884073451679110
0.9	1.9×10^{-3}	8.2×10^{-6}	0.186108051348831675130




Figure 1: The graph of absolute errors between approximate and exact solutions with $M = 18$ and $k = 2$ (Right) $M = 18$ and $k = 3$ for Example 5.2.

Example 5.3 Consider the following nonlinear fractional Lane-Emden type equation:

$$D^\alpha u(x) + \frac{2}{x^{\alpha-\beta}} D^\beta u(x) + \sinh(u(x)) = 0, \quad 0 < x \leq 1, \quad (5.4)$$

subject to the boundary conditions

$$u(0) = 1, \quad u'(0) = 0. \quad (5.5)$$

Wawaz (2001) employed Adomian decomposition method to obtain a series solution of (5.4)-(5.5) when $\alpha = 2$ and $\beta = 1$ as follows

$$u(x) \cong 1 - \frac{(e^2 - 1)x^2}{12e} + \frac{(e^4 - 1)x^4}{480e^2} - \frac{(2e^6 + 3e^2 - 3e^4 - 2)x^6}{30240e^3} + \frac{(61e^8 - 104e^6 + 104e^2 - 61)x^8}{26127360e^4}.$$

We apply the method introduced in Section 4 with $M = 12$ and $k = 2$ and solve this problem with different values of α and β . It can be seen from Fig. 2 that as α and β approach 2 and 1, respectively, the solution of the fractional differential equation approaches that of the integer-order differential equation.




Figure 2: Comparison of $u(x)$ for $M = 12$, $k = 2$ and different values of α and β for Example 5.3

Example 5.4 Consider the following linear singular boundary value problem:

$$D^\alpha u(x) + \frac{1}{x^{\alpha-\beta}} D^\beta u(x) + \frac{1}{1-x} u(x) = 4 \cos x - 5x \sin x + \frac{x^3}{1-x} \cos x, \quad 0 < x < 1, \quad (5.6)$$

$$u(0) = 0, \quad u(1) = \cos(1). \quad (5.7)$$

The exact solution of this example is given in (Zhou and Xu, 2016) as $u(x) = x^2 \cos(x)$ when $\alpha = 2$ and $\beta = 1$. We set $M = 12$ and $k = 2$ for this problem. In Table 3, we report the obtained values of $u(x)$ for different values of α and β . It can be seen from Table 3, the closer the values of α and β are respectively to 2 and 1, the more accurate are the results. For further investigation, we plot the graph of $u(x)$ for different values of α and β with $M = 8$ and $k = 3$ in Fig. 3

Table 3: Obtained absolute errors for $M = 12, k = 2$ and different values of α and β for Example 5.4

x	$\alpha = 1.9, \beta = 0.9$	$\alpha = 1.92, \beta = 0.92$	$\alpha = 1.94, \beta = 0.94$	$\alpha = 1.96, \beta = 0.96$	$\alpha = 1.98, \beta = 0.98$
0.1	3.1×10^{-3}	2.1×10^{-3}	1.2×10^{-3}	5.1×10^{-4}	2.0×10^{-5}
0.3	1.8×10^{-2}	1.4×10^{-2}	9.8×10^{-3}	6.1×10^{-3}	2.8×10^{-3}
0.5	3.2×10^{-2}	2.5×10^{-2}	1.8×10^{-2}	1.2×10^{-2}	5.6×10^{-3}
0.7	3.5×10^{-2}	2.7×10^{-2}	2.0×10^{-2}	1.3×10^{-2}	6.4×10^{-3}
0.9	1.8×10^{-2}	1.4×10^{-2}	1.0×10^{-2}	6.7×10^{-3}	3.4×10^{-3}




Figure 3: The graph of $u(x)$ for different values of α and β with $M = 8, k = 3$ for Example 5.4.

Example 5.5 Consider the following nonlinear fractional Lane-Emden type equation:

$$D^\alpha u(x) + \frac{6}{x^{\alpha-\beta}} D^\beta u(x) + 14u(x) + 4u(x) \ln(u(x)) = 0, \quad 0 < x < 1, \quad (5.8)$$

subject to the boundary conditions

$$u(0) = 1, \quad u(1) = e^{-1}. \quad (5.9)$$

If $\alpha = 2$ and $\beta = 1$, the exact solution of this problem is $u(x) = e^{-x^2}$ (Zhou and Xu, 2016).

In Fig. 4, the graph of $u(x)$ for different values of α and β is plotted. Furthermore, Fig.

5 shows the absolute error between the approximate and exact solution when $\alpha = 2$ and

$\beta = 1$.




Figure 4: The graph of $u(x)$ for different values of α and β with $M = 10$, $k = 3$ for Example 5.5.




Figure 5: Absolute error as $\alpha = 2$ and $\beta = 1$ with $M = 10$, $k = 3$ for Example 5.5.

Example 5.6 Consider the following nonlinear fractional boundary value problem:

$$D^\alpha u(x) + \frac{1}{x^{\alpha-\beta}} D^\beta u(x) + e^{u(x)} = 0, \quad 0 < x < 1, \quad (5.10)$$

subject to the boundary conditions

$$u'(0) = 0, \quad u(1) = 0. \quad (5.11)$$

The exact solution of this problem as $\alpha = 2$ and $\beta = 1$ is given as (Zhou and Xu, 2016)

$$u(x) = 2 \ln \left(\frac{4 - 2\sqrt{2}}{(3 - 2\sqrt{2})x^2 + 1} \right)$$

. In Fig. 6, we plot the graph of $u(x)$ for different values of α and β with $M = 7$, $k = 3$.




Figure 6: The graph of $u(x)$ for different values of α and β with $M = 10$, $k = 3$ for Example 5.6.

Example 5.7 Consider the following linear fractional Lane-Emden type equation:

$$D^\alpha u(x) + \frac{2}{x^{\alpha-\beta}} D^\beta u(x) = \frac{0.76129u(x)}{u(x) + 0.03119}, \quad 0 < x < 1, \quad (5.12)$$

subject to the boundary conditions

$$u'(0) = 0, \quad 5u(1) + u'(1) = 5. \quad (5.13)$$

There is no exact solution for this problem even for integer case. We solve this problem using CWFD with $M = 10$, $k = 2$. In Table 4, the values of $u(x)$ for different fractional values of α and β at different points are tabulated. We make a comparison between the results obtained by the CWFD method, The Laguerre wavelet (LW) method in (Zhou and Xu, 2016), and a numerical method based on the operational matrix of orthonormal Bernoulli polynomial (BP) in (Mohsenyadeh et al., 2015) for integer values of α and β in Table 5. It can be seen from Tables 4 and 5, as α and β are increasing closer to integer values, the results are increasing closer to the ones of integer values. Table 5 shows that our results are very close to the results reported in other two papers and indicates the applicability and accurateness of the proposed method. It can be clearly observed from Tables 4 and 5, as α and β approach 2 and 1, the solution of fractional differential equation approaches to that of the integer-order differential equation.

Table 4: Approximate solutions for different values of α and β with $M = 10, k = 2$ for Example 5.7

x	$\alpha = 1.5, \beta = 0.5$	$\alpha = 1.6, \beta = 0.6$	$\alpha = 1.7, \beta = 0.7$	$\alpha = 1.8, \beta = 0.8$	$\alpha = 1.9, \beta = 0.9$
0.1	0.765457385293227	0.777849130711797	0.790555364088865	0.803503300386760	0.816514521141764
0.2	0.776130801423652	0.786585663532280	0.797659339431068	0.809242556063937	0.821123173757560
0.3	0.789838168401597	0.798441923158554	0.807825369576831	0.817890936326706	0.828429359689670
0.4	0.806115080193247	0.812991206693056	0.820716305422740	0.829224626275748	0.838326176512752
0.5	0.824626305619186	0.829933285174916	0.836100384725762	0.843090763744000	0.850740215946676
0.6	0.845309532918059	0.849128944523233	0.853839766598646	0.859385533350940	0.865614149890049
0.7	0.867713969757805	0.870328122893672	0.873779197287922	0.878013361169784	0.882901208768500
0.8	0.891762974684848	0.893431592884813	0.895829358309583	0.898910454952620	0.902571377092311
0.9	0.917364124571176	0.918348799523956	0.919913436181026	0.922022206810561	0.924598283236026

Table 5: Comparison of approximate solutions $\alpha = 2, \beta = 1$ for Example 5.7

x	<i>CWFD</i>	<i>LW</i>	<i>BP</i>
0.1	0.82970609243381	0.82970609243388	0.82970609243390
0.2	0.83337473359101	0.83337473359108	0.83337473359110
0.3	0.83948991395371	0.83948991395376	0.83948991395381
0.4	0.84805278499607	0.84805278499610	0.84805278499617
0.5	0.85906492716924	0.85906492716925	0.85906492716933
0.6	0.87252831995829	0.87252831995828	0.87252831995828
0.7	0.88844530562320	0.88844530562317	0.88844530562329
0.8	0.90681854806681	0.90681854806678	0.90681854806690
0.9	0.92765098836558	0.92765098836555	0.92765098836568

Example 5.8 Consider the following nonlinear singular fractional two-point BVP:

$$D^\alpha u(x) + \frac{1}{x^{\alpha-\beta}} D^\beta u(x) + \frac{u^2(x)}{x(1-x)} = 4 \arctan x + \frac{1+3x^2}{x(1+x^2)} + \frac{(1+x^2)^2 \arctan^2(x)}{x(1-x)}, \quad 0 < x < 1, \quad (5.14)$$

subject to the boundary conditions

$$u(0) + u'(0) = 1, \quad u(1) + u'(1) = 4.14159265358979. \quad (5.15)$$

The exact solution of this problem if $\alpha = 2$ and $\beta = 1$ is given as $u(x) = (1+x^2) \arctan(x)$ in (Zhou and Xu, 2016). The graphs of $u(x)$ for different values of α and β are plotted in Fig. 7.




Figure 7: The graph of $u(x)$ for different values of α and β for Example 5.8.

6. Conclusion

The singular fractional Lane-Emden type equations as a generalized form of the standard ones subject to the different conditions have been considered. The Chebyshev wavelets finite difference method has been successfully applied to obtain numerical solutions of fractional Lane-Emden type equations with linear and nonlinear terms arising in mathematical physics and astrophysics. It can be clearly seen from figures and tables, as α and β approach the integer values, the solution of fractional differential equation approaches to that of the integer-order differential equation. From obtained results, it can be observed that the proposed method is effective and accurate for solving such type of singular fractional equations. It has also been shown that the accuracy can be enhanced either by expanding the number of subintervals or by expanding the quantity of collocation points in subintervals.

Abdel-Salam, E.A., Nouh, M.I., 2015. Approximate Solution to the Fractional Second Type Lane-Emden Equation. 2015arXiv151209116A.

Abdulaziz, O., Hashim, I., Momani, S., 2008. Solving system of fractional differential equations by homotopy perturbation method. Phys. Lett. A 372 (4), 451-459.

Agrawal, O.P., 2002. Formulation of EulerLagrange equations for fractional variational

problems. *Journal of Mathematical Analysis and Applications*, 272(1), 368–379.

Akgül, A., Inc, M., Karatas, E., Baleanu, D., 2015. Numerical solutions of fractional differential equations of Lane-Emden type by an accurate technique. *Advances in Difference Equations*, 220. 1–12.

Almeida, R., Pooseh, S. and Torres, D.F., 2015. *Computational methods in the fractional calculus of variations*. London: Imperial College Press.

Atangana, A. and Secer, A., 2013, April. A note on fractional order derivatives and table of fractional derivatives of some special functions. In *Abstract and Applied Analysis* (Vol. 2013). Hindawi Publishing Corporation.

Babolian, E., Fattahzadeh, F., 2007. Numerical computation method in solving integral equations by using Chebyshev wavelet operational matrix of integration *Applied Mathematics and Computation*, 188, 1016-1022.

Baleanu, D., Güven, Z.B. and Machado, J.T. eds., 2010. *New trends in nanotechnology and fractional calculus applications* (pp. xii+-531). New York, NY, USA: Springer.

Basdevant, J-L., 2007. *Variational Principles in Physics*, New York: Springer.

Bhrawy, A.H., Zaky, M.A., Van Gorder, R.A., 2016. A space-time Legendre spectral tau method for the two-sided space-time Caputo fractional diffusion-wave equation. *Numerical Algorithms*, 71(1), 151–180.

Bliss, G.A., 1946. *Lectures on the Calculus of Variations*.

Burgess, M., 2002. *Classical Covariant Fields*, Cambridge: Cambridge University Press.

Chaurasia, V.B.L. and Pandey, S.C., 2010. Computable extensions of generalized fractional kinetic equations in astrophysics. *Research in astronomy and astrophysics*, 10(1), p.22.

Clenshaw, C.W., Curtis, A.R., 1960. A method for numerical integration on an automatic computer. *Numerische Mathematik* 2: 197–205.

Daubechies. I., 1992. *Ten Lectures on Wavelets*. vol. 61 of CBMS-NSF Regional Conference Series in Applied Mathematics, SIAM, Philadelphia, Pa, USA.

Dávila, J., Dupaigne, L., Wei, J., 2014. *Trans. On the fractional Lane-Emden equation*,

to appear in *Trans. Amer. Math.*

Daftardar-Gejji, V., Jafari, H., 2007. Solving a multi-order fractional differential equation using Adomian decomposition. *Appl. Math. Comput.* 189, 541–548.

Davis, P.J., Rabinowitz, P., 1984. *Methods of Numerical Integration*(2nd ed.). Computer Science and Applied Mathematics. Orlando, Florida: Academic Press, 1-2.

Diethelm, K. 2010. *The Analysis of Fractional Differential Equations: An Application-Oriented Exposition Using Differential Operator of Caputo Type*. Springer-Verlag Berlin Heidelberg 2010.

Doughty, N.A., 1990. *Lagrangian Interaction*, New York: Addison-Wesley.

El-Ajou, A., Odibat, Z., Momani, S., Alawneh, A., 2010. Construction of Analytical Solutions to Fractional Differential Equations Using Homotopy Analysis Method. *IAENG International Journal of Applied Mathematics*, 40 (2), 43–51 .

Elbarbary, Elsayed M.E., El-Kady, M., 2003. Chebyshev finite difference approximation for the boundary value problems. *Appl. Math. Comput.* 139, 513–523.

Hilfer, R. ed., 2000. *Applications of fractional calculus in physics*. World Scientific, 1–85.

EL Nabulsi, R.A., 2005a. A fractional approach to non-conservative Lagrangian dynamical systems.

EL Nabulsi, R.A., 2005. A fractional action-like variational approach of some classical, quantum and geometrical dynamics.

El-Nabulsi, R.A. and Torres, D.F., 2008. Fractional action-like variational problems, *J. Math. Phys.* 49(5), 053521.

El-Nabulsi, A.R., 2011a. The fractional white dwarf hydrodynamical nonlinear differential equation and emergence of quark stars. *Applied Mathematics and Computation*, 218(6), 2837–2849.

El-Nabulsi, R.A., 2011b. Universal fractional Euler-Lagrange equation from a generalized fractional derivate operator. *Central European Journal of Physics*, 9(1), 250–256.

El-Nabulsi, R.A., 2012. Gravitons in fractional action cosmology. *International Journal*

of Theoretical Physics, 51(12), 3978–3992.

El-Nabulsi, R.A., 2013. Non-standard fractional Lagrangians. *Nonlinear Dynamics*, 74(1-2), 381–394.

El-Nabulsi, R.A., 2016a. A Cosmology Governed by a Fractional Differential Equation and the Generalized Kilbas-Saigo-Mittag-Leffler Function. *International Journal of Theoretical Physics*, 55(2), 625–635.

El-Nabulsi, R.A., 2016b. Implications of the Ornstein-Uhlenbeck-like fractional differential equation in cosmology. *Revista Mexicana de Física*, 62(3), 240–250.

El-Nabulsi, R.A., 2016c. Fractional variational approach with non-standard power-law degenerate Lagrangians and a generalized derivative operator. *Tbilisi Mathematical Journal*, 9(1), 279–293.

Hartley, T.T., Lorenzo, C.F., 1998. A solution to the fundamental linear fractional order differential equation.

Ibrahim RW., 2012. Existence of nonlinear Lane-Emden equation of fractional order, *Miskolc Mathematical Notes*. 13(1), 39–52.

Jacobs, B.A. and Harley, C., 2014, April. Two hybrid methods for solving two-dimensional linear time-fractional partial differential equations. In *Abstract and Applied Analysis* (Vol. 2014). Hindawi Publishing Corporation.

Jafari, H., Yousefi, S.A., Firoozjaee, M.A., Momani, S., Khalique, C.M., 2011. Application of Legendre wavelets for solving fractional differential equations. *Comput.Math. Appl.* 62 (3), 1038–1045.

Kaur, H., Mittal, R.C., Mishra, V., 2013. Haar wavelet approximate solutions for the generalized Lane-Emden equations arising in astrophysics. *Comput. Phys. Commun.* 184, 2169–2177.

Kazemi Nasab, A., Kılıçman, A., Pashazadeh Atabakan, Z., Abbasbandy, S., 2013. Chebyshev Wavelet Finite Difference Method: A New Approach for Solving Initial and Boundary Value Problems of Fractional Order. *Abstr. Appl. Anal.* [Article ID: 916456].

Kazemi Nasab, A., Kılıçman, A., Pashazadeh Atabakan, Z., Leong, W.J., 2015. A

- numerical approach for solving singular nonlinear Lane-Emden type equations arising in astrophysics. *Astronomy*, 34, 178-186.
- Kilbas, A.A., Srivastava, H.M., Trujillo, J.J., 2006. *Theory and Applications of Fractional Differential Equations*, 1st Edition. *Math. Studies*, 204, Elsevier (North-Holland), Amsterdam.
- Lanczos, C., 1986. *The Variational Principles of Mechanics*, New York: Dover.
- Lewandowski, R., Chorazyczewski, B., 2010. Identification of the parameters of the KelvinVoigt and the Maxwell fractional models, used to modeling of viscoelastic dampers. *Computers & structures*, 88(1), 1–17.
- Li, Y.L., 2010. Solving a nonlinear fractional differential equation using Chebyshev wavelets. *Commun. Nonlinear Sci. Numer. Simul.* 15, 2284–2292.
- Li, Y.L., Sun, N., 2011. Numerical solution of fractional differential equations using the generalized block pulse operational matrix. *Comput. Math. Appl.* 62(3), 1046–1054.
- Li, Y.L., Zhao, W.W., 2010. Haar wavelet operational matrix of fractional order integration and its applications in solving the fractional order differential equations. *Appl. Math. Comput.* 216 (8), 2276–2285.
- Mainardi, F., 2012. *Fractional Calculus: Some Basic Problems in Continuum and Statistical Mechanics*. arXiv preprint arXiv:1201.0863.
- Malinowska, A.B. and Torres, D.F., 2012. *Introduction to the fractional calculus of variations* (pp. xvi+-275). London: Imperial College Press.
- Marasi, H.R., Sharifi, N., Piri, H., 2015. Modified differential transform method for singular LaneEmden equations in integer and fractional order. *TWMS J. App. Eng. Math.* 5(1), 124–131.
- Miller, K.S. and Ross, B., 1993. *An introduction to the fractional calculus and fractional differential equations*.
- Misiti, M., Misiti, Y., Oppenheim, G., Poggi, J.-M., 2000. *Wavelet toolbox*, the MathWorks, Inc., Natick, MA.
- Mohsenyazadeh, M., Maleknejad, K., Ezzati, R., 2015. A numerical approach for the

- solution of a class of singular boundary value problems arising in physiology. *Advances in Difference Equations*, 2015 (1), 1–10.
- Momani, S., Odibat, Z., 2007. Numerical approach to differential equations of fractional order. *J Comput Appl Math*, 96–110.
- Momoniat, E., Harley, C., 2006. Approximate implicit solution of a Lane-Emden equation. *New Astronomy*, 11 (7), 520–526.
- Musielak, Z.E., 2008. Standard and non-standard Lagrangians for dissipative dynamical systems with variable coefficients. *Journal of Physics A: Mathematical and Theoretical*, 41(5), p.055205.
- Odibat Z., Momani, S., 2006. Application of variational iteration method to nonlinear differential equations of fractional order. *Int J Nonlinear Sci Numer Simul*, 7, 271–279.
- Odibat Z., Momani S., Erturk, V.S., 2008. Generalized differential transform method: application to differential equations of fractional order. *Appl. Math. Comput.*, 197 (2), 467–477.
- Odziejewicz, T., Malinowska, A.B. and Torres, D.F., 2012. Generalized fractional calculus with applications to the calculus of variations. *Computers & Mathematics with Applications*, 64(10), 3351-3366.
- Odziejewicz, T., Malinowska, A.B. and Torres, D.F., 2013. Fractional calculus of variations of several independent variables. *The European Physical Journal Special Topics*, 222(8), 1813–1826.
- Pandey, Rajesh K., Bhardwaj, A., Kumar, N., 2012. Solution of Lane-Emden type equations using Chebyshev wavelet operational matrix of integration. *Journal of Advanced Research in Scientific Computing*, 4 (1), 1–12.
- Parand, K., Dehghan, M., Rezaeia, A.R.; Ghaderi, S.M., 2010. An approximation algorithm for the solution of the nonlinear Lane-Emden type equations arising in astrophysics using Hermite functions collocation method. *Comput. Phys. Commun.* 181, 1096-1108.
- Podlubny, I., 1999. *An Introduction to Fractional Derivatives, Fractional Differential*

- Equations, Some Methods of Their Solution and Some of Their Applications. Academic Press, San Diego.
- Riewe, F., 1996. Nonconservative lagrangian and hamiltonian mechanics. *Physical Review E*, 53(2), 1890–1899.
- Riewe, F., 1997. Mechanics with fractional derivatives. *Physical Review E*, 55(3), 3581–3592.
- Samko, S., Kilbas, A. and Marichev, O., 1993. *Fractional Integrals and Derivatives; Theory and Applications*. Gordon and Breach. London.
- Saxena, R.K., Mathai, A.M. and Haubold, H.J., 2004. On generalized fractional kinetic equations. *Physica A: Statistical Mechanics and its Applications*, 344(3), 657–664.
- Sharma, M., Ali, M.F. and Jain, R., 2015. Advanced generalized fractional kinetic equation in astrophysics. *Prog. Fract. Differ. Appl*, 1(1), 65–71.
- Stanislavsky, A.A., 2010. Astrophysical Applications of Fractional Calculus. In *Proceedings of the Third UN/ESA/NASA Workshop on the International Heliophysical Year 2007 and Basic Space Science* (pp. 63–78). Springer Berlin Heidelberg.
- Wazwaz, A., 2001. A new algorithm for solving differential equations of Lane-Emden type. *Appl. Math. Comput.* 118, 287–310.
- Weinstock, R., 1974. *Calculus of variations: with applications to physics and engineering*. Courier Corporation.
- Wu, G., Lee, E.W.M., 2010. Fractional variational iteration method and its application. *Phys. Lett. A*, 374, 2506–2509.
- Yousri, Y.H., Abd-Elhameed, W.M., Doha, E.H., 2015. Ultraspherical Wavelets Method for Solving Lane-Emden Type Equations. *Romanian Journal of Physics*, 60 (9).
- Yuanlu, L., 2010. Solving a nonlinear fractional differential equation using Chebyshev wavelets. *Communications in Nonlinear Science and Numerical Simulation*, 15(9), 2284–2292.
- Zhang, Y., 2009. *Finite Difference Method For Fractional Partial Differential Equation*.

Appl. Math. Comput. 215, 524–529.

Zhou, F., Xu, X., 2016. Numerical solutions for the linear and nonlinear singular boundary value problems using Laguerre wavelets. *Advances in Difference Equations*, 2016:17.

ACCEPTED MANUSCRIPT