

## WHY DO PATIENTS LEAVE THE EMERGENCY DEPARTMENT AGAINST MEDICAL ADVICE?

Murat Saritemur, Arzu Denizbasi, Ebru Unal Akoglu, Tuba Cimilli Ozturk, Fatma Sari Dogan

Marmara University, Faculty of Medicine, Department of Emergency Medicine, Istanbul, Turkey

### ABSTRACT

**Introduction:** Leaving against medical advice is a worrying situation which is a problem not only for patients also for physicians. For many different reasons, people may choose to leave the hospital on their own requests. If we know the reasons for leaving, we can find the solution.

**Method:** This prospective study was conducted at an emergency department of a university hospital that is visited by more than 30.000 patients annually. Demographic characteristics such as sex and age, triage level, insurance status of the patients, length of stay in the emergency department and why they were leaving against medical advice were recorded.

**Results:** A total of 321 patients which constituted 1.1% of all admissions to the emergency department during the study period left against medical advice. The main reason was refusing the observation or hospitalization (34.6%; n=111) and the second reason was inadequate health insurance (19.6%; n=63) in all study population (Table 1). Although inadequate health insurance was the second reason in males (36 of 153 patients; 24.0%), the second reason was refusing the intervention or medication in females (52 of 168 patients; 31.1%). The patients who left due to dissatisfaction with health care were predominantly male (72.7%; n=24; p=0.002) whereas who left due to refusing the intervention or medication were predominantly female (88.9%; n=48; p<0.001).

**Conclusion:** Hospitals should endeavor to identify the reasons why patients leave the ED, to document the events in detail and also to solve the problem.

**Key words:** Emergency department, leaving Against Medical Advice

### Corresponding author:

Dr. Murat SARITEMUR  
Department of Emergency Medicine,  
Ataturk University Medical Faculty, 25240, Erzurum, Turkey  
Tel : +90 442 2318399  
Fax : +90 442 2360968  
E-mail: muratsaritemur@gmail.com

**Copyright © 2012- 2014.** Murat SARITEMUR . and al. This is an open access article published under **Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International Public License (CC BY-NC-ND)**. This license allows others to download the articles and share them with others as long as they credit you, but they can't change them in any way or use them commercially.

### INTRODUCTION

Leaving against medical advice (AMA), or known as self-discharge, is a worrying and undesirable situation, which is a problem not only for patients but also for physicians who treat these patients [1]. It means that a patient chooses to leave the hospital before the treating physician recommends discharge. Although discharge should be after the physician's advice, patients admitted to hospital are free to leave whenever they want. This problem has been studied in different hospitals and various patient groups including emergency departments

and the prevalence differs according to the patient groups [1, 2]. Leaving AMA is not uncommon which is estimated at 2% of all discharges from emergency departments going up to 6% in different populations [2-4]. These patients are at significant risks. Patients leaving AMA have higher readmission rates and higher adjusted relative risk of 30-day mortality than patients discharged conventionally [5, 6]. Therefore, treating physicians report distress when patients choose to refuse the treatment and leave AMA [5].

For many different reasons, people may choose to leave the hospital on their own requests. It may be

due to factors related to patients, medical staffs or medical environment. Sometimes, patients may choose to leave AMA when they feel well enough to leave the hospital. However, a conflict between the medical staff and the patient is a more serious cause of self-discharge [2]. If we know the reasons for leaving AMA, we can find solutions.

We aimed to investigate why patients leave the Emergency Department (ED), AMA, factors affecting their decisions, the triage scale of these patients and to discuss what we can do about this population in the ED.

## METHOD

### Study sample and data collection

This study was conducted at an ED of a university hospital in Istanbul that is visited by more than 30,000 patients annually. Patients admitted to our ED and who left AMA before the treatment was completed from June 1, 2007 to May 31, 2008, were enrolled prospectively. A self-constructed questionnaire was used to gather data. Demographic characteristics such as sex and age, triage level according to The Canadian Emergency Department Triage and Acuity Scale (CTAS) [7], insurance status of the patients, length of stay in the ED and why they were leaving the ED AMA were recorded. Patients under 18 years of age, unable to decide by themselves and taken away by their relatives, and patients who did not want to participate in the study were excluded. The Institutional Ethics Committee approved the study protocol.

There were 366 patients who left AMA during the study period, representing 1.1% of all ED admissions at the same period. 45 patients were excluded: six patients were unable to decide and they were taken away by legal relatives and 39 patients refused to participate in the study.

### Definitions

The CTAS was used as the triage method and triage scale was divided into five groups. The number of patients was insufficient for statistical analysis in Level I and II. Therefore, five groups of triage were redesigned to three groups as follows; "emergency" group (EG, Level I-III), "less emergent" group (LEG, level IV) and "non-emergent" group (NEG, level V).

Patients were asked about the primary reasons for leaving AMA and the answers were classified into

five groups including "inadequate health insurance", "dissatisfaction with health care", "longer length of stay", "refusing the observation or hospitalization" and "refusing the intervention or medication".

### Statistical analysis

Statistical Package for Social Sciences software (SPSS 17.0, Chicago, IL, USA) was used for the statistical analysis. Continuous variables were expressed as the mean  $\pm$  standard deviation, whereas categorical variables were presented as percentages. Chi-square ( $\chi^2$ ) test was employed for the comparison of categorical variables. The differences between normally distributed numeric variables were evaluated by Student's t-test or one-way analysis of variance, while non-normally distributed variables were analyzed using Mann-Whitney U test or Kruskal-Wallis variance analysis as appropriate. Statistical significance was assumed for  $p < 0.05$ .

## RESULTS

The final study population included 321 patients representing 1.1% of all patients admitted to the ED during the same period. The mean age was  $48.7 \pm 18.6$  year and 47.7% ( $n=153$ ) of the patients were male. The mean length of stay in the ED was  $150.5 \pm 149.8$  minutes. The main reason of leaving AMA was refusing the observation or hospitalization (34.6%;  $n=111$ ) and the second reason was inadequate health insurance (19.6%;  $n=63$ ) in all study population (Table I).

**Table I: Reasons for leaving AMA**

Reasons for leaving AMA	n (%)
Refusing the observation or hospitalization	111 (34.6)
Inadequate health insurance	63 (19.6)
Longer length of stay	60 (18.7)
Refusing the intervention or medication	54 (16.8)
Dissatisfaction with health care	33 (10.3)

Although inadequate health insurance was the second reason in males (36 of 153 patients; 24.0%), the second reason was refusing the intervention or medication in females (52 of 168 patients; 31.1%). Patients who left due to dissatisfaction with health care were predominantly male (72.7%;  $n=24$ ;  $p=0.002$ ), whereas those who left due to refusing the intervention or medication were predominantly female (88.9%;  $n=48$ ;  $p < 0.001$ ) and the difference was statistically significant (Table II).

**Table II: Reasons for leaving against medical advice (AMA) according to the gender**

Reasons for leaving AMA	Gender		P
	Male; n (%)	Female; n (%)	
Inadequate health insurance	36 (24.0)	27 (16.2)	0.121
Dissatisfaction with health care	24 (15.6)	9 (5.4)	0.002
Longer length of stay	28 (17.5)	32 (18.6)	0.886
Refusing the observation or hospitalization	59 (39.0)	52 (31.1)	0.160
Refusing the intervention or medication	6 (3.9)	48 (28.7)	<0.001
Total	153 (100)	168 (100)	

The triage acuity scales of the patients are showed in table III.


**Table III: The triage scale of the patients who left against medical advice**

Triage scale	(n; %)
Level I	0; 0%
Level II	3; 0.9%
Level III	171; 53.3%
Level IV	96; 29.9%
Level V	51; 15.9%

“Emergent” group (EG, level I-III), “less-emergent” group (LEG, level IV) and “non-emergent” group (NOG, level V)

The mean ages of patients in EG, LEG and NEG were as follows respectively;  $54.0 \pm 18.8$ ,  $46.7 \pm 17.1$  and  $36.1 \pm 12.5$  years and the post-hoc test showed that the difference between each group was

statistically significant ( $p < 0.05$ ). The difference was found to be significant when reasons for leaving were compared with acuity. Of the patients who left due to inadequate health insurance, 57.3% (n=36) were in EG, 38.0% (n=24) in LEG and 4.7% (n=3) in NEG. Of the patients who left due to dissatisfaction with health care, 54.5% (n=18) were in EG, none in LEG and 45.5% (n=15) in NEG. Of the patients who left due to longer length of stay, 36.6% (n=22) were in EG, 46.7% (n=28) in LEG and 16.7% (n=10) in NEG. Of the patients who left due to refusing the observation or hospitalization, 73.2% (n=82) were in EG and only 5.4% (n=6) were in NEG. Of the patients who left due to refusing the intervention or medication, 27.8% (n=15) were in EG, 38.9% (n=21) in LEG and 33.3% (n=18) in NEG (Figure 1).



**Figure 1: Reasons for leaving against medical advice and triage acuity**

The difference between reasons for leaving and age was insignificant ( $p>0.05$ ). The mean waiting time of patients who left due to inadequate health insurance was  $97 \pm 75$  min., due to dissatisfaction with health care was  $133 \pm 151$  min., due to longer length of stay was  $171 \pm 127$  min., due to refusing the observation or hospitalization was  $216 \pm 189$  min., due to refusing the intervention or medication was  $64 \pm 44$  min. post-hoc test showed that the difference between each group according to waiting time was statistically significant. When acuity scale was compared with waiting time, the difference was statistically significant ( $p<0.05$ ). The mean waiting times in EG, LEG and NEG were as follows respectively;  $171 \pm 160$  min.,  $122 \pm 122$  min. and  $130 \pm 148$  min.

## DISCUSSION

We found that the main reasons for leaving AMA were refusing the observation or hospitalization and inadequate health insurance. Although other reasons showed similarity in all study population, the number of patients who left AMA due to refusing the intervention or medication in females was prominent. It is known that the rate of leaving the ED is higher for patients who have less urgent or non urgent conditions; however, studies have shown that patients with potentially emergent, even life threatening conditions are also reported to leave [8, 9]. In our study, more than half of the patients who left the ED were classified as Level I-III according to CTAS and defined as EG. This raises the importance of the subject.

In our study, patients who left the emergency service represented 1.1% of all patients admitted to our ED. Our result is similar to other studies in the literature. However, higher rates were found in other studies [4]. Some studies have revealed a relationship between demographic characteristics and rates for leaving; detecting higher rates for leaving the emergency service especially in male gender and younger age groups [10]. The patients' demographic characteristics have discrepancies among many studies. In our study, females constituted 52.3% of the patients but in the study by Henson et al. [8], males constituted 60% of the patient group. The average age for our patients showed similarity to the other studies in the literature [2, 11, 12]. But the NEG patients were younger than the others.

In most of the other studies, longer length of stay was found to be the dominant reason for leaving the ED [10, 13]. In our study, the rate of patients who left the ED due to longer length of waiting time

seems to be lower than other studies. Waiting period was found to be longer in EG when compared to LEG and NEG. In a previous study performed in our hospital, the waiting period was found to be longer in NEG [12]. This may have been due to the increasing workload of EDs as a result of changes in health system policy in our country. For those patients, since hospital beds were not available, patients had been observed in ED or transferred to a different hospital [14]. So, the waiting time in ED increased.

In the study by Tong and his colleagues, the reasons for leaving the emergency service were classified as patient-related or family-related, physician-related, cultural and social factors [15]. Regarding the waiting period, if it could be shortened, patients' satisfaction would be better, leading to lower rates of leaving the emergency service. Additionally, triage classification of the patients is another factor influencing the waiting period [10]. In the study of Lee et al. [13], 90% of the patients leaving ED were evaluated as "not urgent" or "less urgent". Therefore, it is necessary to make the initial evaluation carefully. In studies performed on patients admitted to emergency services, various criteria were compared to diagnose patients with real emergency conditions, but still a considerable rate of missed diagnoses were observed [16]. For this purpose, various triage evaluation scores have been developed and are currently used in emergency services [7, 17]. Correct identification and faster evaluation of emergency patients will decrease the possible health problems to a minimum. We are not fully aware of the reasons such as refusing the observation, hospitalization, intervention or medication. Therefore, more comprehensive studies are needed.

### Recommendations

The patient leaving the emergency service should not be seen as an incompatible individual. People providing health care services, should endeavor to understand the underlying problems causing refusal of medical treatment. If these problems are identified, the disagreement may also be eliminated. Furthermore, effort spent by both patient and doctor may provide a consensual environment. It was shown that patients were unable to communicate comfortably with the medical care personnel, having difficulty in asking questions or unable to get understandable answers [18]. Therefore, informing the patients and giving appropriate guidance is necessary to eliminate the currently encountered problems. This approach will increase the satisfaction of the patients and their relatives. Patients' satisfaction is a valuable parameter when

evaluating the quality of the services provided in emergency departments [19]. In previous studies, giving explanations to the patient about the emergency service procedures and why the patient should wait were determined to increase patient satisfaction [20]. Taylor et al. pointed out that the improvement of communication skills of emergency service personnel and developing positive attitudes towards their patients, informing the patients in more detail and shortening the waiting period were effective in increasing patient satisfaction in ED [21].

If the patient decides to leave, the specified data-sheet for leaving patients should be dated and signed by the patient. Date and time of admittance and decision to leave should be recorded. If possible, signature of another individual should be taken as a witness [22]. General management of a leaving patient is shown in table IV.

**Table IV: When a patient refuses treatment [22]**

1. Evaluate his/her decision making adequacy.
2. Evaluate patient's point of view on the situation.
3. Try to understand the patient's worries and concerns.
4. Inform about the risks of refusing treatment.
5. If appropriate, call other individuals (eg. family members, social worker).
6. Consider alternative therapies.
7. If possible, provide adequate information regarding the treatment and follow-up.
8. Avoid intimidating and punitive explanations.
9. Document the refusal decision and the result in detail.
10. Think about the telephone follow-up.

Informed consent and informed refusal about treatment are the most fundamental rights of the patients. Emergency service physicians should follow an appropriate route about informed consent and informed refusal of treatment. They should inform the patient about planned medical treatment and interventions and also should act with a persuasive manner about protecting the patient's rights. In the emergency service, many potential obstacles exist about informed consent (For example, inadequacy of patient's decision making ability, language issues, illiteracy, time limitations, communication problems and different expectations). Since general conditions of many emergent patients tend to deteriorate, full attention should be given to them. The adequacy of patient's decision making should be well evaluated, a detailed explanation should be made during informing procedure and the non-compliance of the

patient on planned intervention, its risks and benefits should be abolished.

In conclusion, leaving AMA is a worrying situation and reasons may differ according to the hospitals and patient groups including emergency department patients. Therefore, hospitals should endeavor to identify the reasons why patients leave the ED, to document the events in detail and also to solve the problem.

## REFERENCES

1. Seaborn Moyse H, Osmun WE. Discharges against medical advice: a community hospital's experience. *Canadian journal of rural medicine : the official journal of the Society of Rural Physicians of Canada = Journal canadien de la medecine rurale : le journal officiel de la Societe de medecine rurale du Canada*. 2004 Summer;9(3):148-53. PubMed PMID: 15603687.
2. Ibrahim SA, Kwok CK, Krishnan E. Factors associated with patients who leave acute-care hospitals against medical advice. *American journal of public health*. 2007 Dec;97(12):2204-8. PubMed PMID: 17971552. Pubmed Central PMCID: 2089112.
3. Alfandre DJ. "I'm going home": discharges against medical advice. *Mayo Clinic proceedings*. 2009 Mar;84(3):255-60. PubMed PMID: 19252113. Pubmed Central PMCID: 2664598.
4. Levy F, Mareiniss DP, Iacovelli C. The importance of a proper against-medical-advice (AMA) discharge: how signing out AMA may create significant liability protection for providers. *The Journal of emergency medicine*. 2012 Sep;43(3):516-20. PubMed PMID: 21715123.
5. Alfandre D, Schumann JH. What is wrong with discharges against medical advice (and how to fix them). *JAMA : the journal of the American Medical Association*. 2013 Dec 11;310(22):2393-4. PubMed PMID: 24216713.
6. Southern WN, Nahvi S, Arnsten JH. Increased risk of mortality and readmission among patients discharged against medical advice. *The American journal of medicine*. 2012 Jun;125(6):594-602. PubMed PMID: 22513194. Pubmed Central PMCID: 3372411.
7. Bullard MJ, Unger B, Spence J, Grafstein E, Group CNW. Revisions to the Canadian Emergency Department Triage and Acuity Scale (CTAS) adult guidelines. *Cjem*. 2008 Mar;10(2):136-51. PubMed PMID: 18371252.
8. Henson VL, Vickery DS. Patient self discharge from the emergency department: who is at risk? *Emergency medicine journal : EMJ*. 2005 Jul;22(7):499-501. PubMed PMID: 15983086. Pubmed Central PMCID: 1726841.
9. Ding R, Jung JJ, Kirsch TD, Levy F, McCarthy ML. Uncompleted emergency department care: patients who leave against medical advice. *Academic emergency medicine : official journal of the Society for Academic*

Emergency Medicine. 2007 Oct;14(10):870-6. PubMed PMID: 17766732.

10. Mohsin M, Forero R, Ieraci S, Bauman AE, Young L, Santiano N. A population follow-up study of patients who left an emergency department without being seen by a medical officer. *Emergency medicine journal : EMJ*. 2007 Mar;24(3):175-9. PubMed PMID: 17351221. Pubmed Central PMCID: 2660023.

11. Hwang SW, Li J, Gupta R, Chien V, Martin RE. What happens to patients who leave hospital against medical advice? *CMAJ : Canadian Medical Association journal = journal de l'Association medicale canadienne*. 2003 Feb 18;168(4):417-20. PubMed PMID: 12591781. Pubmed Central PMCID: 143546.

12. Guneyssel O, Onur OE, Saritemur M, Akoglu H, Denizbasi A. General characteristics of patients who leave emergency department against medical advices after being examined. *Turk J Emerg Med*. 2008;8(2):79-83.

13. Lee G, Endacott R, Flett K, Bushnell R. Characteristics of patients who did not wait for treatment in the emergency department: a follow up survey. *Accident and emergency nursing*. 2006 Jan;14(1):56-62. PubMed PMID: 16377190.

14. Emet M, Beyhun NE, Uzkeser M, Aslan S. Descriptive epidemiology of secondary transport from the emergency unit of a state hospital. *Turk J Med Sci* 2008;38(3):263-8.

15. Tong SF, Chen R. A patient who refused medical advice: The doctor and patient should look for a common ground. *Malaysian Family Physician*. 2007;2(3):110-3.

16. Vertesi L. Does the Canadian Emergency Department Triage and Acuity Scale identify non-urgent patients who can be triaged away from the emergency department? *Cjem*. 2004 Sep;6(5):337-42. PubMed PMID: 17381991.

17. Choi YF, Wong TW, Lau CC. Triage rapid initial assessment by doctor (TRIAD) improves waiting time and processing time of the emergency department. *Emergency medicine journal : EMJ*. 2006 Apr;23(4):262-5; discussion -5. PubMed PMID: 16549569. Pubmed Central PMCID: 2579496.

18. Choyce MQ, Maitra AK. Satisfaction with the accident and emergency department--a postal survey of general practitioners' views. *Journal of accident & emergency medicine*. 1996 Jul;13(4):280-2. PubMed PMID: 8832351. Pubmed Central PMCID: 1342732.

19. Al B, Yildirim C, Togun I, Zengin S, Bozkurt S, Kose A, et al. Factors that affect patient satisfaction in emergency department. *JAEM*. 2009;8:39-44.

20. Krishel S, Baraff LJ. Effect of emergency department information on patient satisfaction. *Annals of emergency medicine*. 1993 Mar;22(3):568-72. PubMed PMID: 8442546.

21. Taylor C, Bengner JR. Patient satisfaction in emergency medicine. *Emergency medicine journal : EMJ*. 2004 Sep;21(5):528-32. PubMed PMID: 15333521. Pubmed Central PMCID: 1726409.

22. Strinko JM, Howard CA, Schaeffer SL, Laughlin JA, Berry MA, Turner SN. Reducing risk with

telephone follow-up of patients who leave against medical advice or fail to complete an ED visit. *Journal of emergency nursing: JEN : official publication of the Emergency Department Nurses Association*. 2000 Jun;26(3):223-32. PubMed PMID: 10839849.