

Copyright © 2017 by Academic Publishing House Researcher s.r.o.



Published in the Slovak Republic  
European Journal of Medicine  
Has been issued since 2013.

ISSN: 2308-6513  
E-ISSN: 2310-3434  
2017, 5(2): 31-42

DOI: 10.13187/ejm.2017.2.31  
[www.ejournal5.com](http://www.ejournal5.com)



## Biophysical Research of ZEOLITH detox and ZEOLITH Creme of the Company

Lava Vitae <sup>a</sup>, Ignat Ignatov <sup>a,\*</sup>

<sup>a</sup> Scientific Research Center of Medical Biophysics, Sofia, Bulgaria

### Abstract

We studied the mathematical model of interaction with water of natural mineral and microporous crystalline mineral ZEOLITH detox and ZEOLITH Creme of LavaVitae Company (Austria). In this report are submitted data about the interaction of ZEOLITH detox and ZEOLITH Crème with water, obtained by non-equilibrium (NES) and differential-equilibrium energy spectrum (DNES) of water. The average energy ( $\Delta E_{H...O}$ ) of hydrogen H...O-bonds among individual molecules H<sub>2</sub>O after treatment of ZEOLITH detox with water measured by NES- and DNES-methods is  $\Delta E = -0.0034 \pm 0.0011$  eV for ZEOLITH detox. The average energy ( $\Delta E_{H...O}$ ) of hydrogen H...O-bonds among individual molecules H<sub>2</sub>O after treatment of ZEOLITH detox with water measured by NES- and DNES-methods is  $\Delta E = -0.0034 \pm 0.0011$  eV for ZEOLITH detox. The average energy ( $\Delta E_{H...O}$ ) of hydrogen H...O-bonds among individual molecules H<sub>2</sub>O after treatment of ZEOLITH detox with water measured by NES- and DNES-methods is  $\Delta E = -0.007 \pm 0.0011$  eV for ZEOLITH Creme. These results suggest the restructuring of  $\Delta E_{H...O}$  values among H<sub>2</sub>O molecules with a statistically reliable increase of local extremums in DNES-spectra. The research is performed for ZEOLITH detox with study of pH and oxidative reduction potential (ORP).

**Keywords:** ZEOLITH detox, ZEOLITH Crème, nanostructures, mathematical model, NES, DNES.

### 1. Introduction

The ZEOLITH detox is mineral refers to new generation of natural mineral sorbents (NMS). Zeolites are the aluminosilicate members of the family of microporous solids known as "molecular sieves", named by their ability to selectively sort molecules based primarily on a size exclusion process. Natural zeolites form when volcanic rocks and ash layers react with alkaline groundwater. Zeolites also crystallize in post-depositional environments over periods ranging from thousands to millions of years in shallow marine basins. Naturally occurring zeolites are rarely pure and are contaminated to varying degrees by other minerals, metals, quarts, or other zeolites. For this reason, naturally occurring zeolites are excluded from many important commercial applications where uniformity and purity are essentials.

As natural mineral zeolite has unusually broad scope of application in industry. Adsorption, catalytic, and reduction-oxidation Zeolites is widely used in industry as a desiccant of gases and

\* Corresponding author

E-mail addresses: [mbioph@dir.bg](mailto:mbioph@dir.bg) (I. Ignatov)

liquids, for treatment of drinking and sewage water from heavy metals, ammonia, phosphorus, as catalyst in petrochemical industry for benzene extraction, for production of detergents and for extracting of radionuclides in nuclear reprocessing. It is also used in medicine as nutritional supplements having antioxidant properties. Some authors make qualifications of zeolites as nano materials.

A wide range of properties of zeolite defines the search for new areas of industrial application of these minerals in science and nano technology that contributes to a deeper study the mechanism of interaction of these minerals with water. The company LavaVitae produces ZEOLITH Creme with exceptional results on the skin and skin diseases. This paper deals with evaluating of mathematical models of interaction of ZEOLITH detox and ZEOLITH Creme with water.

## **2. Materials and Methods**

### **2.1. Materials**

The study is performed with samples of ZEOLITH detox ZEOLITH Creme from LavaVitae Company.

**There are valid the following methods for research of zeolite.**

### **2.2. Analytical Methods**

The analytical methods were accredited by the Institute of Geology of Ore Deposits. Petrography, Mineralogy, and Geochemistry (Russian Academy of Sciences). Samples were treated by various methods as ICP-OES, GC, and SEM.

### **2.3. Gas-Chromatography**

Gas-chromatography (GC) is performed at Main Testing Centre of Drinking Water (Moscow, the Russian Federation) on Kristall 4000 LUX M using Chromaton AW-DMCS and Inerton-DMCS columns (stationary phases 5% SE-30 and 5% OV-17), equipped with flame ionization detector (FID) and using helium (He) as a carrier gas.

### **2.4. Inductively Coupled Plasma Optical Emission Spectrometry (ICP-OES)**

The mineral composition is studied by inductively coupled plasma optical emission spectrometry (ICP-OES) on Agilent ICP 710-OES (Agilent Technologies, USA) spectrometer, equipped with plasma atomizer (under argon stream), MegaPixel CCD detector, and 40 MHz free-running, air-cooled RF generator, and Computer-optimized exhale system: the spectral range at 167–785 nm; plasma gas: 0–22.5 l/min in 1.5 l/min; power output: 700–1500 W in 50 W increments.

### **2.5. Transmission Electron Microscopy (TEM)**

The structural studies were carried out with using JSM 35 CF (JEOL Ltd., Korea) device, equipped with X-ray microanalyzer “Tracor Northern TN”, SE detector, thermomolecular pump, and tungsten electron gun (Harpin type W filament, DC heating); working pressure:  $10^{-4}$  Pa ( $10^{-6}$  Torr); magnification: 300.000, resolution: 3.0 nm, accelerating voltage: 1–30 kV; sample size: 60–130 mm.

### **2.6. IR-Spectroscopy**

IR-spectra of water samples, obtained after being contacted 3 days with shungite and zeolite, are registered on Fourier-IR spectrometer Bruker Vertex (“Bruker”, Germany) (a spectral range: average IR – 370–7800  $\text{cm}^{-1}$ ; visible – 2500–8000  $\text{cm}^{-1}$ ; the permission – 0.5  $\text{cm}^{-1}$ ; accuracy of wave number – 0.1  $\text{cm}^{-1}$  on 2000  $\text{cm}^{-1}$ );

**For the research of ZEOLITH detox and ZEOLITH Creme the methods are:**

### **2.7. Non-equilibrium Spectrum (NES) and Differential Non-equilibrium Spectrum (DNES)**

The energy spectrum of water is characterized by a non-equilibrium process of water droplets evaporation, therefore, the term non-equilibrium spectrum (NES) of water is used. The difference  $\Delta f(E) = f(\text{samples of water}) - f(\text{control sample of water})$  – is called the “differential non-equilibrium energy spectrum of water” (DNES).

### **2.8. Measurement of pH and ORP (oxidative-redox potential)**

The research is performed from Georgi Gluhchev with device from Hanna Instruments.

## **3. Results and Discussion**

In comparison with zeolite comprises a microporous crystalline aluminosilicate mineral commonly used as commercial adsorbents, three-dimensional framework of which is formed by linking via the vertices the tetrahedral  $[\text{AlO}_4]^{2-}$  and  $[\text{SiO}_4]^{2-}$  (Panayotova, Velikov, 2002). Each

tetrahedron  $[\text{AlO}_4]^{2-}$  creates a negative charge of the carcasses compensated by cations ( $\text{H}^+$ ,  $\text{Na}^+$ ,  $\text{K}^+$ ,  $\text{Ca}^{2+}$ ,  $\text{NH}_4^+$ , etc.), in most cases, capable of cation exchange in solutions. Tetrahedrons formed the secondary structural units, such as six-membered rings, five-membered rings, truncated octahedra, etc. Zeolites framework comprise interacting channels and cavities forming a porous structure with a pore size of 0.3–1.0 nm. Average crystal size of the zeolites may range from 0.5 to 30  $\mu\text{m}$ .

By the measurement of IR spectra in the range of vibrations in the crystal mineral framework one can obtain the information: a) on the structure of the framework, particularly type lattice ratio  $\text{SiO}_2/\text{Al}_2\text{O}_3$ , nature and location of cations and changes in the structure in the process of the thermal treatment; b) on the nature of the surface of the structural groups, which often serve as adsorption and catalytically active sites.

Other method for obtaining information about the average energy of hydrogen bonds in an aqueous sample is measuring of the spectrum of the water state. It was established experimentally that at evaporation of water droplet the contact angle  $\theta$  decreases discretely to zero, whereas the diameter of the droplet changes insignificantly (Antonov, 1995). By measuring this angle within a regular time intervals a functional dependence  $f(\theta)$  can be determined, which is designated by the spectrum of the water state (Ignatov, 2005; Ignatov, 2012; Ignatov, Mosin, 2013). For practical purposes by registering the spectrum of water state it is possible to obtain information about the averaged energy of hydrogen bonds in an aqueous sample. For this purpose the model of W. Luck was used, which consider water as an associated liquid, consisted of  $\text{O}-\text{H}\dots\text{O}-\text{H}$  groups (Luck et al., 1980). The major part of these groups is designated by the energy of hydrogen bonds ( $-E$ ), while the others are free ( $E = 0$ ). The energy distribution function  $f(E)$  is measured in electron-volts ( $\text{eV}^{-1}$ ) and may be varied under the influence of various external factors on water as temperature and pressure.

For calculation of the function  $f(E)$  experimental dependence between the water surface tension measured by the wetting angle ( $\theta$ ) and the energy of hydrogen bonds ( $E$ ) is established:

$$f(E) = b f(\theta) / 1 - (1 + b E)^2)^{1/2},$$

where  $b = 14.33 \text{ eV}^{-1}$ ;  $\theta = \arcsin(-1 - b E)$

The energy of hydrogen bonds ( $E$ ) measured in electron-volts ( $\text{eV}$ ) is designated by the spectrum of energy distribution. This spectrum is characterized by non-equilibrium process of water droplets evaporation, thus the term “non-equilibrium energy spectrum of water” (NES) is applied.

The difference  $\Delta f(E) = f(\text{samples of water}) - f(\text{control sample of water})$  – is designated the “differential non-equilibrium energy spectrum of water” (DNES).

DNES is calculated in milli-electron volts (0.001  $\text{eV}$  or  $\text{meV}$ ) is a measure of changes in the structure of water as a result of external factors. The cumulative effect of all other factors is the same for the control sample of water and the water sample, which is under the influence of this impact. The research with NES method of water drops received after 3 days stay with zeolite in deionized water may also give valuable information on the possible number of hydrogen bonds as percent of water molecules with different values of distribution of energies. These distributions are basically connected with restructuring of  $\text{H}_2\text{O}$  molecules with the same energies.

### **3.1. Results with spectral analysis of 1% solution of ZEOLITH detox**

The average energy ( $E_{\text{H}\dots\text{O}}$ ) of hydrogen  $\text{H}\dots\text{O}$ -bonds among individual  $\text{H}_2\text{O}$  molecules in 1 % solution of ZEOLITH detox is measured at  $E = -0.1219 \text{ eV}$ . The result for the control sample (deionized water) is  $E = -0.1185 \text{ eV}$ . The results obtained with the NES method are recalculated with the DNES method as a difference of the NES (1 % solution of ZEOLITH detox) minus the NES (control sample with deionized water) equalled the DNES spectrum of 1 % solution of ZEOLITH detox. Thus, the result for 1 % solution of ZEOLITH detox recalculated with the DNES method is  $\Delta E = -0.0034 \pm 0.0011 \text{ eV}$ . The result shows the increasing of the values of the energy of hydrogen bonds in 1 % solution of ZEOLITH detox regarding the deionized water. The result is effect of stimulation on human body. This shows restructuring of water molecules in configurations of clusters, which influence usefully on human health on molecular and cellular level. The effects are describing with mathematical model of 1 % solution of ZEOLITH detox.

### 3.2. Mathematical model of ZEOLITH detox

The research with the NES method of water drops is received with 1 % solution ZEOLITH detox, and deionized water as control sample. The mathematical models of 1 % solution ZEOLITH detox gives the valuable information for the possible number of hydrogen bonds as percent of H<sub>2</sub>O molecules with different values of distribution of energies (Table 1 and Fig. 1). These distributions are basically connected with the restructuring of H<sub>2</sub>O molecules having the same energies.

**Table 1.** The distribution (% , (-E<sub>value</sub>)/(-E<sub>total value</sub>) of H<sub>2</sub>O molecules in 1 % water solution of ZEOLITH detox (product of LavaVitae, Austria) and control deionized water

| -E(eV)<br>x-axis | 1 % water<br>solution<br>ZEOLITH<br>detox<br>(LavaVitae)<br>y-axis<br>(%((-E <sub>value</sub> )* /<br>(-E <sub>total value</sub> )) <sup>**</sup> | Control<br>Sample<br>Deionized<br>water<br>y-axis<br>(%((-<br>E <sub>value</sub> )* / (-<br>E <sub>total value</sub> )) <sup>**</sup> | -E(eV)<br>x-axis | 1 % water<br>solution<br>ZEOLITH<br>detox<br>(LavaVitae)<br>y-axis<br>(%((-E <sub>value</sub> )* /<br>(-E <sub>total<br/>value</sub> )) <sup>**</sup> | Control<br>Sample<br>Deionized water<br>y-axis<br>(%((-E <sub>value</sub> )* / -E <sub>total<br/>value</sub> ) <sup>**</sup> |
|------------------|---|---|------------------|---|--|
| 0.0937           | 0   | 6.7   | 0.1187           | 0   | 15.5   |
| 0.0962           | 0   | 6.7   | 0.1212           | <b>18.9<sup>2</sup></b>   | 0  |
| 0.0987           | 0   | 6.7   | 0.1237           | 0   | 6.7  |
| 0.1012           | 6.0   | 15.5  | 0.1262           | 0   | 6.7  |
| 0.1037           | 12.5  | 6.7   | 0.1287           | 0   | 0  |
| 0.1062           | 0   | 6.7   | 0.1312           | 0   | 3.3  |
| 0.1087           | 3.1   | 0   | 0.1337           | 12.5  | 0  |
| 0.1112           | <b>3.1<sup>1</sup></b>  | 0   | 0.1362           | 12.5  | 3.3  |
| 0.1137           | 0   | 15.5  | 0.1387           | <b>18.9<sup>3</sup></b>   | 0  |
| 0.1162           | 12.5  | 0   | –                | –   | –  |

Notes:

E=-0.1212 eV is the local extremum for anti inflammatory effect


E= -0.1387 eV is the local extremum for inhibition of development of tumor cells of molecular level

Notes:

\* The result (-E<sub>value</sub>) is the result of hydrogen bonds energy for one parameter of (-E)

\*\* The result (-E<sub>value</sub>) is the total result of hydrogen bonds energy

Figure 1 shows the distribution (% , (-E<sub>value</sub>)/(-E<sub>total value</sub>) of H<sub>2</sub>O molecules in and 1 % of water solution of ZEOLITH detox (product of LavaVitae, Austria) (red line) and control sample deionized water (blue line).



**Fig. 1.** Mathematical model (Ignatov, Mosin, 2013) of 1 % water solution of ZEOLITH detox (product of LavaVitae, Austria)

Notes:

$E = -0.1212$  eV is the local extremum for anti-inflammatory effect

$E = -0.1387$  eV is the local extremum for inhibition of development of tumor cells of molecular level

The experimental data obtained testified the following conclusions from the mathematical model of in 1 % water solution of ZEOLITH detox (product of LavaVitae, Austria) and control deionized water. The distribution ( $\%$ ,  $(-E_{\text{value}})/(-E_{\text{total value}})$ ) of water molecules in mathematical model of in 1 % water solution of ZEOLITH detox (product of LavaVitae, Austria) and control deionized water. The distribution ( $\%$ ,  $(-E_{\text{value}})/(-E_{\text{total value}})$ ) of water molecules in ZEOLITH detox (product of LavaVitae, Austria) according control sample is different. However, for the value  $E = -0.1387$  eV or  $\lambda = 8.95$   $\mu\text{m}$  there is the biggest local extremum (18.9 ( $\%$ ,  $(-E_{\text{value}})/(-E_{\text{total value}})$ )) corresponding to the restructuring of hydrogen bonds among  $\text{H}_2\text{O}$  molecules for inhibition of development of tumor cells of molecular level. This difference may indicate on the different number of hydrogen bonds in water samples, as well as their physical parameters (pH, ORP), resulting in different distribution of  $\text{H}_2\text{O}$  molecules and different values of  $\text{H}_2\text{O}$  molecules with ratios of  $(-E_{\text{value}})/(-E_{\text{total value}})$ . Particularly it was observed the statistical re-structuring of  $\text{H}_2\text{O}$  molecules in water samples according to the energies. The experimental data may prove that stipulates the restructuring of  $\text{H}_2\text{O}$  molecules on molecular level and may be used for the prophylaxis of development of tumor cells. For the value  $E = -0.1212$  eV or  $\lambda = 10.23$   $\mu\text{m}$  there is the bigger local extremum (18.9 ( $\%$ ,  $(-E_{\text{value}})/(-E_{\text{total value}})$ )) corresponding to the restructuring of hydrogen bonds among  $\text{H}_2\text{O}$  molecules for anti inflammatory effect.. The experimental data for ZEOLITH detox may prove that stipulates the restructuring of  $\text{H}_2\text{O}$  molecules on molecular level and the biophysical effects are:

$E = -0.1212$  eV is the local extremum for anti inflammatory effect

$E = -0.1387$  eV is the local extremum for inhibition of development of tumor cells of molecular level

### 3.3. Results with spectral analysis of 1% solution of ZEOLITH Creme

The average energy ( $E_{\text{H...O}}$ ) of hydrogen H...O-bonds among individual  $\text{H}_2\text{O}$  molecules in 1 % solution of ZEOLITH Creme is measured at  $E = -0.1200$  eV. The result for the control sample (deionized water) is  $E = -0.1130$  eV. The results obtained with the NES method are recalculated with the DNES method as a difference of the NES (1 % solution of ZEOLITH Creme) minus the NES (control sample with deionized water) equaled the DNES spectrum of 1 % solution of ZEOLITH



Creme. Thus, the result for 1 % solution of ZEOLITH Creme recalculated with the DNES method is  $\Delta E = -0.007 \pm 0.0011$  eV. The result shows the increasing of the values of the energy of hydrogen bonds in 1 % solution of ZEOLITH detox regarding the deionized water. The results is effect of stimulation on human body. The result is 6.4 times more than statistical reliable result. This shows restructuring of water molecules in configurations of clusters, which influence usefully on human health on molecular and cellular level. The effects are describing with mathematical model of 1 % solution of ZEOLITH Creme.

**3.2. Mathematical model of ZEOLITH detox**

The research with the NES method of water drops is received with 1 % solutions ZEOLITH Creme and deionized water as control samples. The mathematical models of 1 % solution ZEOLITH Creme give the valuable information for the possible number of hydrogen bonds as percent of H<sub>2</sub>O molecules with different values of distribution of energies (Table 1). These distributions are basically connected with the restructuring of H<sub>2</sub>O molecules having the same energies.

**3.3. Mathematical model of ZEOLITH Creme (product of the company LavaVitae)**

The research with the NES method of water drops is received with 1 % solution ZEOLITH Creme, and deionized water as control sample. The mathematical model of 1 % solution ZEOLITH Creme gives the valuable information for the possible number of hydrogen bonds as percent of H<sub>2</sub>O molecules with different values of distribution of energies (Table 2 and Fig. 2). These distributions are basically connected with the restructuring of H<sub>2</sub>O molecules having the same energies.

**Table 2.** The distribution (% , (-E<sub>value</sub>)/(-E<sub>totalvalue</sub>)) of H<sub>2</sub>O molecules in 1 % water solution of ZEOLITH Creme (product of LavaVitae, Austria) and control deionized water

| -E(eV)<br>x-axis | 1 % water<br>solution<br>ZEOLITH<br>Creme<br>(LavaVitae)<br>y-axis<br>(%((-E <sub>value</sub> )* /<br>(-E <sub>total value</sub> ))** | Control<br>Sample<br>Deionized<br>water<br>y-axis<br>(%((-E <sub>value</sub> )* /<br>(-E <sub>total value</sub> ))** | -E(eV)<br>x-axis | 1 % water<br>solution<br>ZEOLITH<br>Creme<br>(LavaVitae)<br>y-axis<br>(%((-E <sub>value</sub> )* /<br>(-E <sub>total<br/>value</sub> ))** | Control<br>Sample<br>Deionized water<br>y-axis<br>(%((-E <sub>value</sub> )* /<br>(-E <sub>total<br/>value</sub> ))** |
|------------------|---|--|------------------|---|---|
| 0.0937           | 0   | 0  | 0.1187           | 0   | 5.7   |
| 0.0962           | 0   | 11.4   | 0.1212           | <b>15.2<sup>2</sup></b>   | 5.7   |
| 0.0987           | 7.8   | 5.7  | 0.1237           | 3.8   | 0   |
| 0.1012           | 7.8   | 5.7  | 0.1262           | 3.8   | 5.7   |
| 0.1037           | 3.8   | 11.4   | 0.1287           | 7.7   | 5.7   |
| 0.1062           | 7.8   | 11.4   | 0.1312           | 7.7   | 0   |
| 0.1087           | 3.8   | 0  | 0.1337           | 7.7   | 0   |
| 0.1112           | <b>3.8<sup>1</sup></b>  | 5.7  | 0.1362           | 3.8   | 5.7   |
| 0.1137           | 0   | 8.8  | 0.1387           | <b>7.7<sup>3</sup></b>  | 5.7   |
| 0.1162           | 7.8   | 5.7  | -                | -   | -   |

Notes:


E=-0.1212 eV is the local extremum for anti-inflammatory effect

Notes:

\* The result (-E<sub>value</sub>) is the result of hydrogen bonds energy for one parameter of (-E)

\*\* The result (-E<sub>value</sub>) is the total result of hydrogen bonds energy

Figure 2 shows the distribution (% , (-E<sub>value</sub>)/(-E<sub>total value</sub>)) of H<sub>2</sub>O molecules in and 1 % of water solution of ZEOLITH Creme (product of LavaVitae, Austria) (red line) and control sample deionized water (blue line).



**Fig. 2.** Mathematical model (Ignatov, Mosin, 2013) of 1% water solution of ZEOLITH Creme (product of LavaVitae, Austria).

Notes:

$E = -0.1212$  eV is the local extremum for anti-inflammatory effect

The experimental data obtained testified the following conclusions from the mathematical model of in 1% water solution of ZEOLITH Creme (product of LavaVitae, Austria) and control deionized water. The distribution ( $\%$ ,  $(-E_{\text{value}})/(-E_{\text{total value}})$ ) of water molecules in mathematical model of in 1% water solution of ZEOLITH Creme (product of LavaVitae, Austria) and control deionized water. The distribution ( $\%$ ,  $(-E_{\text{value}})/(-E_{\text{total value}})$ ) of water molecules in ZEOLITH Creme (product of LavaVitae, Austria) according control sample is different. However, for the value  $E = -0.1212$  eV or  $\lambda = 10.23$   $\mu\text{m}$  there is the biggest local extremum (15.2%,  $(-E_{\text{value}})/(-E_{\text{total value}})$ ) corresponding to the restructuring of hydrogen bonds among  $\text{H}_2\text{O}$  molecules for inhibition of development of tumor cells of molecular level. This difference may indicate on the different number of hydrogen bonds in water samples, resulting in different distribution of  $\text{H}_2\text{O}$  molecules and different values of  $\text{H}_2\text{O}$  molecules with ratios of  $(-E_{\text{value}})/(-E_{\text{total value}})$ . Particularly it was observed the statistical re-structuring of  $\text{H}_2\text{O}$  molecules in water samples according to the energies. The experimental data may prove that stipulates the restructuring of  $\text{H}_2\text{O}$  molecules on molecular level and may be used for cleaning of skin with anti-inflammatory effect. The experimental data for ZEOLITH Crème may prove that stipulates the restructuring of  $\text{H}_2\text{O}$  molecules on molecular level and the biophysical effect is:

$E = -0.1212$  eV is the local extremum for anti-inflammatory effect

#### 4. Results with pH and ORP


There are valid the following results of pH as indicator for acid alkaline medium of the products of Lava Vitae. There are the results also of ORP or Oxidation-reduction potential.

The results are for 1% of solutions of products, which are made from deionized water. This research is performed with Georgi Gluhchev from Bulgarian Academy of Science. The results of pH of deionized water is 6.05 and of ORP is 119.7. Table 3 shows the results of pH and ORP.

**Table 3.** Results of products of company LavaVitae for pH and ORP

| Product         | pH        | ORP (mV) | Coordinates<br>Fig. 2     |
|-----------------|-----------|----------|---------------------------|
| VITA Intense    | 4.07±0.02 | -104.5   | Point 1<br>(4,07; -104.5) |
| BOOST           | 3.60±0.02 | +113.6   | Point 2<br>(3,90;113.6)   |
| ZEOLITH detox   | 8.01±0.02 | +109.5   | Point 3<br>(8,01;103.3)   |
| Deionized water | 6.05±0.02 | +119.7   |                           |

Figure 3 shows the dependence between the acidity and basicity (pH) of electrochemically activated solutions and the oxidation-reduction potential (ORP). The pH value within the interval from 3 to 10 units and the ORP within the interval from -400 mV to +900 mV characterize the area of the biosphere of microorganisms. Outside these ranges of pH and ORP the microorganisms will hardly survive.



**Fig. 3.** The dependence between acidity and basicity (pH) of solutions and the ORP on the biosphere of micro-organisms (point 1; VITA intense), (point 2; BOOST), point 3; ZEOLITH detox).

Owing to the unique porous structure the mineral Zeolites are ideal absorbents and fillers (Gorshteyn et al., 1979), and as sorbents have a number of positive characteristics:

- High adsorption capacity, characterized by low resistance to water pressure;
- Mechanical strength and low abrasion resistance;
- Corrosion-resistance;
- Absorption capacity relatives to many substances, both organic (oil, benzene, phenol, pesticides, etc.) and inorganic (chlorine, ammonia, heavy metals);
- Catalytic activity;
- Relatively low cost;
- Environmental friendliness and ecological safety.

## 5. Discussion and Conclusions

### 5.1. ZEOLITH detox (product of LavaVitae company)

The interaction of ZEOLITH detox with water is quiet complex and results the restructuring



of energy values among H<sub>2</sub>O molecules with a statistically reliable increase of local extremums in DNES-spectra after treatment of ZEOLITH detox with water. These values are measured at -0.1219 eV for ZEOLITH detox. The result for control sample (deionized water) is -0.1185 eV. The results with NES method were recalculated by the DNES method. The result of ZEOLITH detox with DNES method is  $0.0034 \pm 0.0011$  eV.

From the NES and DNES spectrum and mathematical model of 1 % solution of ZEOLITH detox and deionized water as control sample are valid the following conclusions for biophysical effects for ZEOLITH detox (LavaVitae Company)

- Anti-inflammatory effect;
- inhibition of development of tumor cells of molecular level;

Naturally occurring zeolites are rarely pure and are contaminated to varying degrees by other minerals, metals, quarts, or other zeolites. For this reason, naturally occurring zeolites are excluded from many important commercial applications where uniformity and purity are essential. In comparison with zeolite comprises a microporous crystalline aluminosilicate mineral commonly used as adsorbent. Zeolite creates a negative charge of the carcasses compensated by cations (H<sup>+</sup>, Na<sup>+</sup>, K<sup>+</sup>, Ca<sup>2+</sup>, NH<sub>4</sub><sup>+</sup>, etc.), in most cases, capable of cations exchange in solutions. Efficiency of using zeolite is stipulated by the high range of valuable properties (absorption, catalytic, antioxidant, regenerative, antibacterial). There is permanent antioxidant activity of zeolite on enzymes (Dogliotti et al., 2012; Ignatov, Mosin, 2015).

### 5.2. ZEOLITH Creme (product of LavaVitae company)

From the NES and DNES spectrum and mathematical model of 1% solution of ZEOLITH Crème and deionized water as control sample is valid the following conclusion for biophysical effects for ZEOLITH Creme (LavaVitae company)

- anti-inflammatory effect;

Naturally occurring zeolites are rarely pure and are contaminated to varying degrees by other minerals, metals, quarts, or other zeolites. For this reason, naturally occurring zeolites are excluded from many important commercial applications where uniformity and purity are essential. In comparison with zeolite comprises a microporous crystalline aluminosilicate mineral commonly used as adsorbent. Zeolite creates negative charge of the carcasses compensated by cations (H<sup>+</sup>, Na<sup>+</sup>, K<sup>+</sup>, Ca<sup>2+</sup>, NH<sub>4</sub><sup>+</sup>, etc.), in most cases, capable of cations exchange in solutions. Efficiency of using zeolite is stipulated by the high range of valuable properties (absorption, catalytic, antioxidant, regenerative, antibacterial). There is permanent antioxidant activity of zeolite on enzymes (Dogliotti et al., 2012; Ignatov, Mosin, 2015).

## 6. Acknowledgements

The author wish to thank to Georgi Gluhchev for the research of pH and ORP. The author express thankful to Teodora Todorova for the preparation of the figures.

## References

- Antonov, 1995 – Antonov, A. (1995). Research of the Nonequilibrium Processes in the Area in Allocated Systems. Diss. Thesis Doctor of Physical Sciences. Sofia: Blagoevgrad: pp. 1–255.
- Gluhchev et al., 2015a – Gluhchev, G., Ignatov, I., Karadzhov, S., Miloshev, G., Ivanov, I., Mosin, O.V. (2015). Biocidal Effects of Electrochemically Activated Water. *Journal of Health, Medicine and Nursing*, V. 11, pp. 67-83.
- Gluhchev et al., 2015b – Gluhchev, G., Ignatov, I., Karadzhov, S., Miloshev, G., Ivanov, N., Mosin, O.V. (2015). Electrochemically Activated Water: Biophysical and Biological Effects of Anolyte and Catholyte Types of Water. *European Journal of Molecular Biotechnology*, V.1, pp. 12-26.
- Gluhchev et al., 2015c – Gluhchev, G., Ignatov, I., Karadzhov, S., Miloshev, G., Ivanov, N., Mosin, O.V. (2015). Studying the Antimicrobial and Antiviral Effects of Electrochemically Activated NaCl Solutions of Anolyte and Catholyte on a Strain of E. Coli DH5 and Classical Swine Fever (CSF) Virus. *European Journal of Medicine*, 9 (3), pp. 124-138.
- Gluhchev et al., 2015d – Gluhchev, G., Ignatov, I., Karadzhov, S., Miloshev, G., Ivanov, I., Mosin, O.V. (2015). Electrochemically Activated Water. Biophysical and Biological Effects of Anolyte and Catholyte as Types of Water. *Journal of Medicine, Physiology and Biophysics*, Vol. 10, pp. 1-17.

[Gluhchev et al., 2015e](#) – Gluhchev, G., Ignatov, I., Karadzhov, S., Miloshev, G., Ivanov, I., Mosin, O. V. (2015). Studying of Virucidal and Biocidal Effects of Electrochemically Activated Anolyte and Catholyte Types of Water on Classical Swine Fever Virus (CSF) and Bacterium E. coli DH5. *Journal of Medicine, Physiology and Biophysics*, Vol. 13, pp. 1-17.

[Ignatov et al., 2014a](#) – Ignatov, I., Karadzhov, S., Atanasov, A., Ivanova, E., Mosin, O. V. (2014). Electrochemical Aqueous Sodium Chloride Solution (Anolyte and Catholyte) as Types of Water. Mathematical Models. Study of Effects of Anolyte on the Virus of Classical Swine Fever Virus. *Journal of Health, Medicine and Nursing*, Vol. 8, pp. 1-28.

[Ignatov et al., 2014b](#) – Ignatov, I., Mosin, O. V., Bauer, E. (2014). Carbonaceous Fullerene Mineral Shungite and Aluminosilicate Mineral Zeolite. Mathematical Model and Practical Application of Water Solution of Water Shungite and Zeolite. *Journal of Medicine, Physiology and Biophysics*, Vol. 4, pp. 27-44.

[Ignatov et al., 2015](#) – Ignatov, I., Mosin, O.V., Gluhchev, G., Karadzhov, S., Miloshev, G., Ivanov, N. (2015). The Evaluation of Mathematical Model of Interaction of Electrochemically Activated Water Solutions (Anolyte and Catholyte) with Water. *European Reviews of Chemical Research*, Vol. 2 (4), pp. 72-86.

[Ignatov et al., 2015a](#) – Ignatov, I., Gluhchev, G., Karadzhov, S., Miloshev, G., Ivanov, I., Mosin, O. V. (2015). Preparation of Electrochemically Activated Water Solutions (Catholyte/Anolyte) and Studying of their Physical-Chemical Properties. *Journal of Medicine, Physiology and Biophysics*, Vol. 13, pp. 18-38.

[Ignatov et al., 2015b](#) – Ignatov, I., Gluhchev, G., Karadzhov, S., Miloshev, G., Ivanov, I., Mosin, O. V. (2015). Preparation of Electrochemically Activated Water Solutions (Catholyte/Anolyte) and Studying of their Physical-Chemical Properties. *Journal of Health, Medicine and Nursing*, Vol. 13, pp. 64-78.

[Ignatov et al., 2015c](#) – Ignatov, I., Mosin, O. V., Gluhchev, G., Karadzhov, S., Miloshev, G., Ivanov, I., (2015). Studying Electrochemically Activated NaCl Solutions of Anolyte and Catholyte by Methods of Non-Equilibrium Energy Spectrum (NES) and Differential Non-Equilibrium Energy Spectrum (DNES). *Journal of Medicine, Physiology and Biophysics*, Vol. 14, pp. 6-18.

[Ignatov et al., 2015d](#) – Ignatov, I., Gluhchev, G., Karadzhov, S., Ivanov, N., Mosin, O.V. (2015). Preparation of Electrochemically Activated Water Solutions (Catholyte/Anolyte) and Studying Their Physical-Chemical Properties. *Journal of Medicine, Physiology and Biophysics*, Vol. 16, pp. 1-14.

[Ignatov et al., 2016](#) – Ignatov, I., Gluhchev, G., Karadzhov, S., Miloshev, G., Ivanov, I., Mosin, O.V. (2015). Preparation of Electrochemically Activated Water Solutions (Catholyte/Anolyte) and Studying Their Physical-Chemical Properties. *Journal of Medicine, Physiology and Biophysics*, Vol. 11: pp. 1-21.

[Ignatov et al., 2016a](#) – Ignatov, I., Mosin, O.V., Gluhchev, G., Karadzhov, S., Miloshev, G., Ivanov, I. (2016). Studying Electrochemically Activated NaCl Solutions of Anolyte and Catholyte by Methods of Non-Equilibrium Energy Spectrum (NES) and Differential Non-Equilibrium Energy Spectrum (DNES). *Journal of Medicine, Physiology and Biophysics*, Vol. 20: pp. 13-23.

[Ignatov et al., 2016b](#) – Ignatov, I. et al. (2016). Results of Biophysical and Nano Technological Research of ZEOLITH Detox of LavaVitae Company. *Journal of Health, Medicine and Nursing*, Vol. 30, pp. 44-49.

[Ignatov, 2005](#) – Ignatov, I. (2005). Energy Biomedicine. *Gea-Libris, Sofia*, pp. 1–88.

[Ignatov, 2010](#) – Ignatov, I. (2010). Which Water is Optimal for the Origin (Generation) of Life? *Euromedica*, Hanover: 34-35.

[Ignatov, 2011](#) – Ignatov, I. (2011). Entropy and Time in Living Matter. *Euromedica*, 74 p.

[Ignatov, 2012](#) – Ignatov, I. (2012). Origin of Life and Living Matter in Hot Mineral Water, Conference on the Physics, Chemistry and Biology of Water. *Vermont Photonics*, USA.

[Ignatov, 2016](#) – Ignatov, I. (2016). Product of LavaVitae BOOST is Increasing of Energy of Hydrogen Bonds among Water Molecules in Human Body. *Journal of Medicine, Physiology and Biophysics*, Vol. 27, pp. 30-42.

[Ignatov, 2016a](#) – Ignatov, I. (2016). VITA intense – Proofs for Anti-inflammatory, Antioxidant and Inhibition Growth of Tumor Cells Effects. Relaxing Effect of Nervous System. Anti Aging Influence. *Journal of Medicine, Physiology and Biophysics*, Vol. 27, pp. 43-61.

[Ignatov, 2017](#) – Ignatov, I. (2017). Aluminosilicate Mineral Zeolite. Interaction of Water Molecules in Zeolite Table and Mountain Water Sevtopolis from Bulgaria. *Journal of Medicine, Physiology and Biophysics*, Vol. 31, pp. 41-45.

[Ignatov, 2017a](#) – Ignatov, I. (2017). VITA intense and Boost – Products with Natural Vitamins and Minerals for Health. *Journal of Medicine, Physiology and Biophysics*, Vol. 31, pp. 58-78.

[Ignatov, 2017b](#) – Ignatov, I. (2017). ZEOLITH detox for Detoxification and ZELOLITH Creme for Skin Effects as Products of LavaVitae Company. *Journal of Medicine, Physiology and Biophysics*, Vol. 31, pp. 79-86.

[Ignatov, Mosin, 2013a](#) – Ignatov I., Mosin O.V. (2013). Possible Processes for Origin of Life and Living Matter with Modeling of Physiological Processes of Bacterium *Bacillus Subtilis* in Heavy Water as Model System. *Journal of Natural Sciences Research*, Vol. 3 (9): pp. 65-76.

[Ignatov, Mosin, 2013b](#) – Ignatov, I., Mosin, O. V. (2013). Modeling of Possible Processes for Origin of Life and Living Matter in Hot Mineral and Seawater with Deuterium. *Journal of Environment and Earth Science*, Vol. 3(14), pp. 103-118.

[Ignatov, Mosin, 2013c](#) – Ignatov, I., Mosin, O. V. (2013). Structural Mathematical Models Describing Water Clusters. *Journal of Mathematical Theory and Modeling*, Vol.3 (11), pp. 72-87.

[Ignatov, Mosin, 2014a](#) – Ignatov, I., Mosin, O. V. (2014). The Structure and Composition of Carbonaceous Fullerene Containing Mineral Shungite and Microporous Crystalline Aluminosilicate Mineral Zeolite. Mathematical Model of Interaction of Shungite and Zeolite with Water Molecules. *Advances in Physics Theories and Applications*, Vol.28, pp. 10-21.

[Ignatov, Mosin, 2014b](#) – Ignatov, I., Mosin, O.V. (2014). The Structure and Composition of Shungite and Zeolite. Mathematical Model of Distribution of Hydrogen Bonds of Water Molecules in Solution of Shungite and Zeolite. *Journal of Medicine, Physiology and Biophysics*, Vol. 2, pp. 20-36.

[Ignatov, Mosin, 2014c](#) – Ignatov, I., Mosin, O.V. (2014). Mathematical Models of Distribution of Water Molecules Regarding Energies of Hydrogen Bonds. *Journal of Medicine, Physiology and Biophysics*, Vol. 2, pp. 71-94.

[Ignatov, Mosin, 2014d](#) – Ignatov, I., Mosin, O.V. (2014). Mathematical Model of Interaction of Carbonaceous Fullerene Containing Mineral Shungite and Aluminosilicate Mineral Zeolite with Water. *Journal of Medicine, Physiology and Biophysics*, Vol. 3, pp. 15-29.

[Ignatov, Mosin, 2014e](#) – Ignatov, I., Mosin, O.V. (2014). Methods for Measurements of Water Spectrum. Differential Non-equilibrium Energy Spectrum Method (DNES). *Journal of Health, Medicine and Nursing*, Vol. 6, pp. 50-72.

[Ignatov, Mosin, 2014f](#) – Ignatov, I., Mosin, O.V. (2014). Nano Mix of Shungite and Zeolite for Cleaning of Toxins and Increasing of Energy of Hydrogen Bonds among Water Molecules in Human Body. *Journal of Medicine, Physiology and Biophysics*, Vol. 27, pp. 1-10.

[Ignatov, Mosin, 2014g](#) – Ignatov, I., Mosin, O.V. (2014). Mathematical Models of Distribution of Water Molecules Regarding Energies of Hydrogen Bonds. *Journal of Medicine, Physiology and Biophysics*, Vol. 6, pp. 50-72.

[Ignatov, Mosin, 2014h](#) – Ignatov, I., Mosin, O.V. (2014). Structural Models of Water and Ice Regarding the Energy of Hydrogen Bonding. *Nanotechnology Research and Practice*, Vol. 7 (3): pp. 96-117.

[Ignatov, Mosin, 2015](#) – Ignatov, Mosin (2015). Origin of Life and Living Matter in Hot Mineral Water. *Advances in Physics Theories and Applications*, Vol. 39, 1-22.

[Ignatov, Mosin, 2016](#) – Ignatov, I., Mosin, O.V. (2016). Deuterium, Heavy Water and Origin of Life. LAP LAMBERT Academic Publishing, pp. 1-500.

[Ignatov, Mosin, 2016a](#) – Ignatov, I., Mosin, O.V. (2016). Water for Origin of Life. *Altaspera Publishing & Literary Agency Inc.* pp. 1-616. [in Russian]

[Luck et al., 1980](#) – Luck, W., Schiöberg, D., Ulrich, S. (1980). Infrared Investigation of Water Structure in Desalination Membranes. *J. Chem. Soc. Faraday Trans.*, Vol. 2(76), pp. 136-147.

[Mehandjiev et al., 2017](#) – Mehandjiev, D., Ignatov, I., Karadzhov, I., Gluhchev, G., Atanasov, A. (2017). On the Mechanism of Water Electrolysis, *Journal of Medicine, Physiology and Biophysics*, Vol. 31, pp. 23-26.

[Mosin, Ignatov, 2012](#) – Mosin, O.V., Ignatov, I. (2012). Composition and Structural Properties of Fullerene Natural Mineral Shungite. *Nanomaterials and Nanotechnologies*, 2, pp. 25-36.

[Mosin, Ignatov, 2012a](#) – Mosin, O.V., Ignatov, I. (2012). The Composition and Structural Properties of Fullerene Natural Mineral Shungite. *Nanoengineering*, Vol. 18 (12), pp. 17-24 [in Russian].

[Mosin, Ignatov, 2013b](#) – Mosin, O.V., Ignatov, I. (2013). The Structure and Composition of Natural Carbonaceous Fullerene Containing Mineral Shungite. *International Journal of Advanced Scientific and Technical Research*, Vol. 6(11–12), pp. 9–21.

[Mosin, Ignatov, 2015](#) – Mosin, O.V., Ignatov, I. (2015). An Overview of Methods and Approaches for Magnetic Treatment of Water. *Water: Hygiene and Ecology*, Vol. 3-4 (4): pp. 113-130.

[Panayotova, Velikov, 2002](#) – Panayotova, M., Velikov, B. (2002). Kinetics of Heavy Metal Ions Removal by Use of Natural Zeolite. *Journal of Environmental Science and Health*, Vol. 37(2): pp. 139–147.

[Parfen'eva, 1994](#) – Parfen'eva, L.S. (1994). Electrical Conductivity of Shungite Carbon. *Solid State Physics*, Vol. 36(1), pp. 234–236.

[Podchaynov, 2007](#) – Podchaynov, S.F. (2007). Mineral zeolite – a Multiplier of Useful Properties Shungite. Shungites and human safety. *Proceedings of the First All-Russian scientific-practical conference* (3–5 October 2006), ed. J.K Kalinin (Petrozavodsk: Karelian Research Centre of Russian Academy of Sciences), pp. 6–74 [in Russian].