

Copyright © 2017 by Academic Publishing House Researcher s.r.o.



Published in the Slovak Republic
European Journal of Medicine
Has been issued since 2013.
ISSN: 2308-6513
E-ISSN: 2310-3434
2017, 5(2): 43-55

DOI: 10.13187/ejm.2017.2.43
www.ejournal5.com



Research with Model Systems in Biophysics and Biochemistry of Bioinfluence of Dimitar Risimanski

Ignat Ignatov ^{a, *}

^a Scientific Research Center of Medical Biophysics, Sofia, Bulgaria

Abstract

This paper presents the results of evaluation of possible biophysical methods and approaches for registering of various non-ionizing radiation (NIR) wave types of the human body in the electromagnetic and optical range. Many types of NIR (electromagnetic waves, infrared radiation, thermo radiation, bioluminescence) emitted from the human body were reviewed. In particular the results on of spontaneous biophoton emission and delayed luminescence from the human body are submitted along with *infrared thermography (IRT)* results. It was shown that 1 cm² of skin generally emits ~85 photons for 1s. The intensity of biophoton emission ranges from 10⁻¹⁹ to 10⁻¹⁶ W/cm² (approx. ~1–1000 photons·cm⁻²·s⁻¹). The specific photon emission from part of the human thumb was detected as a spectrum of various colors with the method of Color coronal spectral analysis on a device with an electrode made of polyethylene terephthalate (PET hostafan) with applied electric voltage 15 kV, electric impulse duration 10 μs, and electric current frequency 15 kHz. It was established that photons corresponding to a red color emission of visible electromagnetic spectrum have energy at 1.82 eV. The orange color of visible electromagnetic spectrum has energy at 2.05, yellow – 2.14, blue-green (cyan) – 2.43, blue – 2.64, and violet – 3.03 eV. The reliable result measurement norm was at E ≥ 2.53 eV, while the spectral range of the emission was within 380–495 nm and 570–750 nm±5 nm. Also were estimated some important physical characteristics (energy of hydrogen bonds, wetting angle, surface tension) of water by the methods of non-equilibrium energy (NES) and differential non-equilibrium energy (DNES) spectrum of water, that helps understand in general how electromagnetic radiation interacts with water and establish the structural characteristics of water.

Keywords: electromagnetic waves, infrared radiation, thermo radiation, bioluminescence, color coronal spectral analysis, NES, DNES.

1. Introduction

All living organisms have a cellular therefore, a molecular organized structure. The living processes inside of them run on a cellular and a molecular level. Bioelectrical activity is one of the very important physical parameters of living organisms (Ignatov et al., 1998). Bioelectric potentials generated by various cells are widely used in medical diagnostics (Rubik, 2002) and are recorded as electrocardiogram, electromyogram, electroencephalogram, etc. It was proved that the human

* Corresponding author
E-mail addresses: mbioph@dir.bg (I. Ignatov)

body and tissues emanate weak electromagnetic waves, the electric voltage of which is denoted as resting potential, action potential, omega-potential etc. (Dobrin et al., 1979; Adey, 1981). Between the outer surface of the cell membrane and the inner contents of the cell there is always the electric potential difference which is created because of different concentrations of K^+ , Na^+ and Cl^- inside and outside of the cell and their different permeability through the cell membrane (Kiang et al., 2005). Their value in the human body varies $\sim 50\text{--}80$ mV and is defined by the galvanic contact of a voltmeter input with an object that indicates on the galvanic type of their source (Cleary, 1993). When being excited a living cell changes the membrane electric potential due to changes in membrane permeability and active ion movement through the membrane. In cells of excitable tissues (muscle, nervous), these processes can occur within a very short time intervals (milliseconds) and are called “current action” potential. Its magnitude makes up ~ 120 mV. Electromagnetic fields refer to non-ionizing radiation (NIR), i.g. the radiative energy that, instead of producing charged ions when passing through matter, has sufficient energy only for excitation. Nevertheless it is known to cause biological effects (Kwan-Hoong, 2003). The NIR spectrum is divided into two main regions, optical radiations and electromagnetic fields. The optical spectrum can be further sub-divided into ultraviolet, visible, and infra-red. The electromagnetic fields are further divided into radiofrequency (microwave, very high frequency and low frequency radio wave). NIR encompass the long wavelength (> 100 nm) and low photon energy (< 12.4 eV) portion of the electromagnetic spectrum, from 1 Hz to $3 \cdot 10^{15}$ Hz. As a result of research carried out in the 1990-s and subsequent years, it was established the property of animal and plant tissues to generate relatively strong transient NIR electric fields due to mechanical stresses and temperature changes in biological structure (Anderson, 1993). These electric fields are mainly due to the piezoelectric and pyroelectric voltage electric polarization of natural biological structures. Owing to cell metabolism, electric dipoles (polar and ionized molecules) involved in polarization of biostructures are continuously destroyed and restored, i.e. this is a non-equilibrium polarization (Barnes, Greenebaum, 2006). Such type of non-equilibrium electric polarization is known as a main characteristic of electrets (Gubkin, 1978). Electrets include dielectric insulators and semiconductors, which under certain conditions, i.g. under the influence of a strong electrostatic field or ionizing radiation, light and other factors acquire property to generate an external electric field, existing for a long time (days, years) and slowly diminishes because the destruction of their substance by polarization (Sessler, Gerhard-Multhaupt, 1998). Along with the electromagnetic field electrets generate specific electric currents produced by heating – thermally stimulated current (TSC) (Gross, 1964). Electrets belong to the non-galvanic type of electrical sources, which tend to a strong electric field (up to 10^6 V/m) and the infinitesimal electric current ($\sim 10^{-14}$ A/mm²). By analogy with the physical fields the electric field emitted from the human body on its physical characteristics resembles the electric field generated by electrets. The electrets play an important role in functioning of many biological structures as they themselves possess electret properties. The bioelectret field registered on the surface of the human body basically are generated by the basal cells of the epidermis (Marino, 1988). Dermis cells adjacent to the bottom layer of basal cells are surrounded by a conductive interstitial fluid, which electric voltage while grounding of the human body is close to zero (so called ground potential). This interstitial fluid screens off electromagnetic fields of underlying tissues. With the average thickness of the epidermis (~ 0.1 mm) and the maximum value of electric voltage (~ 30.0 V), the electric field strength can reach significant values at ~ 300000 V/m (Seto et al., 1992). The strength of the electric field is quiet sufficient for its influence on the biological processes in cells and surrounding tissues, including the synthesis of proteins and nucleic acids (Liboff et al., 1984; Frey, 1993; Shimizu et al., 1995). This electric field along with the field of transmembrane assymetry of ions concentrated at inside and outside of the membrane ($\sim 10^5$ V/cm²) can participate in the cooperative effects in cell membrane structures (Holzel, Lamprecht, 1994; Miller, 1986). Thus, owing to the bioelectret condition of certain subcellular structures in the cell and its surroundings is generated slowly oscillating electric field that is strong enough to influence the biological processes. This field and the electric field due to the piezoelectric voltage and intramembrane electric field forms the total electromagnetic field of the cell and its supracellular structures. It is known that the human skin emanates electromagnetic waves in close ultraviolet range, optic range and also in close infrared range. Infrared thermal bioradiation is found in the middle infrared range at wavelengths from 8 to 14 μm . At wavelength

of 9.7 μm infrared bioradiation has its maximum value at $t = 36.6$ °C. At this temperature the skin emission is closest to the emission of absolute black body (ABB) being at the same temperature. Infrared emission penetrates the skin surface at a depth of ~ 0.1 mm, and is reflected in accordance with the physical laws of reflection of the visible part of the electromagnetic spectrum. Evidently, radiation energy influences tissues while being absorbed by them. Yu.V. Gulyaev and E.E. Godik (Gulyaev, Godik, 1984) determined that the threshold of skin sensitivity for infrared radiation compiled $\sim 10^{-14}$ W/cm². When thermal influence is applied to the point of threshold skin sensitivity, there is developed a physiological reaction toward the thermal current. The intensity of the radiated thermal current generated by skin makes up $\sim 2.6 \cdot 10^{-2}$ W/cm². The second component of electromagnetic waves is bioluminescence (Young, Roper, 1976; Chang et al., 1998). It is supposed that biophotons, or ultraweak photon emissions of biological objects, are weak electromagnetic waves in the optical range of the spectrum (Cohen, Popp, 1997). The typical observed emission of biological tissues in the visible and ultraviolet frequencies ranges from 10^{-19} to 10^{-16} W/cm² (~ 1 – 1000 photons $\cdot\text{cm}^{-2}\cdot\text{sec}^{-1}$) (Edwards et al., 1989; Choi et al., 2002). This light intensity is much weaker than that one to be seen in the perceptually visible and well-studied spectrum of normal bioluminescence detectable above the background of thermal radiation emitted by tissues at their normal temperature (Niggli, 1993). Bioelectric emission from parts of the human body as thumbs can be easily detected with the method of Color coronal spectral analysis under applying gas electrical discharge of high voltage and frequency developed by I. Ignatov (Ignatov, 2005). This method has big scientific and practical prospects in biophysics and medical diagnostics (Chiang et al., 2005). Its advantages include safety, sterility, clarity and interpretability of the data obtained, ease of storage and subsequent computer data processing, the ability to monitor the development of processes in time, comparing the structural, functional and temporal processes etc. The purpose of this research was studying of possible biophysical methods and approaches for registering various NIR wave's types emitted from the human body (electromagnetic waves, infrared radiation, thermo radiation) and methods of their visualization by different technique including magnetography, *infrared thermography*, *chemiluminescence*, and color coronal gas discharge spectral analysis.

2. Materials and Methods

2.1. Registration of electromagnetic fields

The registration of electromagnetic fields was used with super conductive detectors based on Joseffson junctions – device made by sandwiching a thin layer of insulating nonsuperconducting material between two layers of superconducting cooper pairs (S-I-S). This allows the registering of magnetic fields 10^{10} times weaker than the Earth's magnetic field. The study of electric field nearly the human body was done using a standard Faraday cage formed by conducting material (aluminium foil) blocks external static and non-static electric fields by channeling electricity through the conducting material, providing constant voltage on all sides of the enclosure.

2.2. Color coronal gas discharge spectral analysis

Experiments were carried out by using selective high-frequency electric discharge (SHFED) on a device with the electrode made of polyethylene terephthalate (PET, hostafan) with an electric voltage on the electrode 15 kV, electric impulse duration 10 μs , and electric current frequency 15 kHz. The electrode of the device was made of hostafan, and was filled up with electro-conductive fluid. The spectral range of the emission was in the range 380–495 nm and 570–750 \pm 5 nm. The measurements were measured in electronvolts (eV). Detection of gas discharge glowing was conducted in a dark room equipped with a red filter. On the electrode put a photosensitive paper or color film. The object under study (human thumb) was placed on top of a sheet of photo paper or color film. Between the object and the electrode were generated impulses of the electric voltage 15 kV and electric current frequency – 15–24 kHz; on the reverse side of the electrode was applied the transparent electrically conductive thin copper coating. Under these conditions in the thin contact gas space between the studied object and electrode was generated gas electric discharge in the form of characteristic glow around the object – a corona gas electric discharge in the range of 280–760 nm, illuminates a color photo or a photographic film on which was judged about the bioelectric properties of the studied object. Along with the visible range, for this method were obtained color spectra in UV and IR range. Evaluation of the characteristic parameters of

snapshots was based on the analysis of images treated by standard software package. Statistical processing of the experimental data was performed using the statistical package STATISTISA 6 using Student's t-criterion (at $p < 0.05$).

2.3. NES and DNES experiments on interaction of electromagnetic field with water

The research was made with the method of non-equilibrium spectrum (NES) and differential non-equilibrium spectrum (DNES). The device measures the angle of evaporation of water drops from 72° to 0° . As the main estimation criterion was used the average energy ($\Delta E_{H...O}$) of hydrogen O...H-bonds between H_2O molecules in water's samples. The spectrum of water was measured in the range of energy of hydrogen bonds $0.08-0.1387$ eV or $8.9-13.8$ μm with using a specially designed computer program. The study with Risimanski was performed with deionized water and 1 % of glucose solution.

3. Results and Discussion

3.1. Color coronal gas discharge spectral analysis

Coronal gas discharge effect is indicated by the glow corona electrical discharge (flooding, crown, streamer) on the surface of objects being placed in the alternating electric field of high frequency (10–150 kHz) and electric voltage (5–30 kV) (Kilrian, 1949). In this process in the ionization zone develops the gas corona discharge sliding on dielectric surface, occurring in a nonuniform electric field near the electrode with a small radius of curvature. In the thin air layer with thickness of $\sim 10-100$ μm between the studied object and the electrode are developed the following processes:

1) Excitation, polarization and ionization by electric field of high frequency the main components of air – the molecules of nitrogen (78 % N_2), oxygen (21 % O_2) and carbon dioxide (0.046 % CO_2). In the result of this is formed an ionized gas, i.e. gas with separated electrons having negative charges, creating a conductive medium as plasma;

2) Formation of a weak electric current in the form of free electrons separated from molecules of N_2 , O_2 and CO_2 , which generate gas discharge between the studied object and the electrode. The form of gas discharge glowing, its density and surface brightness distribution is determined mainly by electromagnetic properties of the object;

3) The transition of electrons from lower to higher energy levels and back again, during which there appears a discrete quantum of light radiation in the form of photon radiation. The transition energy of electrons depends on the external electric field and the electronic state of the studied object. Therefore, in different areas surrounding the electric field, the electrons receive different energy impulses, i.e. “skipping” at different energy levels those results in emission of photons with different wavelengths (frequencies) and the energy, coloring the contour of the glow in various spectral colors. Processes outlined above form the total gas electric effect (Ignatov, Mosin, 2012), allows to study the electrical properties of the object at its interaction with an external electromagnetic field (Ignatov, Mosin, 2013a; Ignatov, Mosin, 2013b). It was shown that the electrical conductivity of the object has almost no effect on the formation of the electric images, which mostly depends on the dielectric constant (Pehek et al., 1976).

There is a relationship (1) of the electric discharge per unit area of the recording medium on the following parameters:

$$\sigma = [\alpha - U_p(d_2 + \delta)/d_2] \varepsilon_0(d_2 + \delta) / \delta d_2, \quad (1)$$

$$\text{where: } \delta = d_1/\varepsilon_1 + d_3/\varepsilon_3$$

α – slope rate of electrical pulse;

T – duration of the electrical pulse;

U_p – breakdown voltage of the air layer between the subject and the recording medium;

d_1 – the width of the object;

d_2 – width of the zone of influence of the electromagnetic field;

d_3 – width of the recording medium;

ε_0 – dielectric permittivity of the air ($\varepsilon_0 = 1.00057$ F/m);

ε_1 – dielectric permittivity of the studied object;

ε_3 – dielectric permittivity of the medium.

To calculate the breakdown voltage of the air layer is used this formula:

$$U_p = 312 + 6,2d_2 \quad (2)$$

As a result of mathematical transformations is obtained a quadratic equation describing the width of the air layer:

$$6,2d_2^2 - (\alpha T - 6,2\delta - 312)d_2 + 312\delta = 0 \quad (3)$$

This equation has two solutions:

$$d_2 = [\alpha T - 6,2\delta - 312] \pm [(\alpha T - 6,2\delta - 312)^2 - 7738\delta]^{1/2} \quad (4)$$


The above equations allow calculating maximum and minimum width of the air layer for the occurrence of electric discharge under which is being formed the electrical image of the studied object. Gas discharge characteristics for various biological objects vary in character and light intensity, size of contour glow and color spectrum and depend both on its own electromagnetic radiation and the dielectric constant of the object. The intensity depends on the electric voltage applied on the electrode. Studies have shown that the contours of gas discharge glow at 12 kHz and 15 kHz are homogeneous in their structure. The contour at kHz is 55 % of the contour at 15 kHz and at 24 kHz – only 15 % of the contour at 15 kHz that is important for further analysis and identification of images. The incidence of bioelectrical activity of the body reducing the intensity of gas discharge glow. Pathology in the organism and surrounding tissues also alter the bioelectric activity and the shape and color of gas discharge glow, which is determined mainly by energy of photon emission at the transition of electrons from higher energy levels to the lower ones when being excited by the external electric field. Thus, for red color of the electromagnetic spectrum this energy compiles 1.82 eV, for orange color – 2.05 eV, yellow – 2.14 eV, blue-green (cyan) – 2.43 eV, blue – 2.64 eV, and violet – 3.03 eV. The reliable result norm is at $E \geq 2.53$ eV.

The spectral range of the photon emission for different colors is within 380-495 nm and 570–750 nm±5 nm. The photons, corresponding to the emission with green color in the visible electromagnetic spectrum, are not being detected under those experimental conditions. Thus, the more predominant in the color spectrum yellow, orange, blue, blue-green and purple colors, the more pronounced is gas discharge glow and bioelectric properties of the object. According to the data obtained, the incidence of bioelectrical activity of the body reducing the intensity of gas discharge glow. Studies carried out by A. Antonov and I. Ignatov on 1120 patients shown that the overall drop in the bioelectric activity of the body, as well as pathology in organism alter the bioelectric activity and reduce the apparent size of the gas discharge glow. This dependence is observed for many disorders, although there are not statistical reliable results that this method can be applied in medical diagnostics. The research area was from part of the thumb contacted with transparent electrode. The norm of energy of photon emission compiles 2.54 eV. If the value is over than 2.54 eV this is an indicator of normal bioelectrical status. Some people with high energy status possess the values of photon emission over 2.90 eV. The high values of this parameter are possible with practicing of yoga, sport etc. The emission less than 2.53 eV is characteristic for people with low bioelectrical status. These results are interesting from scientific point of view, because they may provide brilliant prospects for further using of this method for biophysical studies. interesting from scientific point of view, because they may provide prospects for further using of this method for biophysical studies.


Figure 1 shows the results of of Dimitar Risimanski with color coronal discharge. The bioelectrical discharge in norm is on fig. 1a). The code of bioelectrical photography of Risimanski is 1b). The coronal image of the man A.A. before bioinfluence of Risimanski is 3c). The picture is with blue biophoton emission 2.64 eV. The result of the man A.A. after influence on Risimanski is 3d). It is with biophoton emission with blue and predominant violet color and biophoton emission – 3.00 eV.




a). Bioelectrical discharge of a person in norm (1.94 eV)



b). Bioelectrical discharge image of Dimitar Risimanski (3.03 eV)



c). Bioelectrical discharge image of A.A.. (2.64 eV)



d). Bioelectrical discharge image of A.A. (3.00 eV)

Fig. 1. Bioelectrical discharge images of the research with Dimitar Risimanski (I. Ignatov)

3.2. NES and DNES analysis of water

Water seems to be a good model system for studying the interaction with electromagnetic field and structural research. The recent data indicated that water is a complex associated non-equilibrium liquid consisting of associative groups (clusters) containing from 3 to 50 individual H_2O molecules (Keutsch, Saykally, 2011). These associates can be described as unstable groups (dimers, trimers, tetramers, pentamers, hexamers etc.) in which individual H_2O molecules are linked by van der Waals forces, dipole-dipole and other charge-transfer interactions, including hydrogen bonding (Ignatov, Mosin, 2013c). At room temperature, the degree of association of H_2O molecules may vary from 2 to 21. The measurements were performed with using NES and DNES methods. It was established experimentally that the process of evaporation of water drops, the wetting angle θ decreases discretely to 0, and the diameter of water drop basis is only slightly altered, that is a new physical effect (Antonov, Yuskesslieva, 1983). Based on this effect, by means of measurement of the wetting angle within equal intervals of time is determined the function of distribution of H_2O molecules according to the value of $f(\theta)$. The distribution function is denoted as the energy spectrum of the water state. A theoretical research established the dependence between the surface tension of water and the energy of hydrogen bonds among individual H_2O -molecules (Antonov, 1995). The hydrogen bonding results from interaction between electron-deficient H-atom of one H_2O molecule (hydrogen donor) and unshared electron pair of an electronegative O-atom (hydrogen acceptor) on the neighboring H_2O molecule; the structure of hydrogen bonding may be defined as $O \cdots H^{\square+} - O^{\square}$.

For calculation of the function $f(E)$ represented the energy spectrum of water, the experimental dependence between the wetting angle (θ) and the energy of hydrogen bonds (E) is established:

$$f(E) = b \times f(\theta) / 1 - (1 + b \times E)^{1/2}, \quad (4)$$

$$\text{where } b = 14.33 \text{ eV}^{-1} \quad (5)$$

The relation between the wetting angle (θ) and the energy (E) of the hydrogen bonds between H_2O molecules is calculated by the formula:

$$\theta = \arcsin(-1 - 14.33E) \quad (6)$$

The energy spectrum of water is characterized by a non-equilibrium process of water droplets evaporation, therefore, the term non-equilibrium spectrum (NES) of water is used. The energy of hydrogen bonds measured by NES is determined as $\bar{E} = -0,1067 \pm 0,0011 \text{ eV}$.

The difference $\Delta f(E) = f(\text{samples of water}) - f(\text{control sample of water})$ – is called the “differential non-equilibrium energy spectrum of water” (DNES).

Thus, DNES spectrum is an indicator of structural changes of water as a result of various external factors. The cumulative effect of these factors is not the same for the control sample of water and the water sample being under the influence of this factor.

Figure 2 shows NES-spectrum of deionized water that was used as a model system for studying the interaction of electromagnetic field with water. On the X-axis are given three scales. The energies of hydrogen bonds among H_2O molecules are calculated in eV. On the Y-axis is shown the energy distribution function $f(E)$ of H_2O molecules measured in eV^{-1} . It was shown that the window of transparency of the earth atmosphere for the electromagnetic radiation in the middle IR-range almost covers NES-spectrum of water. Arrows A and B designate the energy of hydrogen bonds among H_2O molecules. Arrow C designates the energy at which the human body behaves itself as absolute black body (ABB) at optimum temperature $36.6 \text{ }^\circ\text{C}$ and adsorbs the thermal radiation. A horizontal arrow designates the window of transparency of the earth atmosphere for the electromagnetic radiation in the middle IR-range.




Fig. 2. Non-equilibrium energy spectrum (NES) of water as a result of measurement for 1 year: λ – wavelength, k – wave number

Another important physical parameter was calculated with using NES and DNES methods – the average energy ($\Delta E_{H...O}$) of H...O-bonds between H₂O compiled -0.1067 ± 0.0011 eV. The most remarkable peculiarity of H...O-bond consists in its relatively low strength; it is 5–10 times weaker than chemical covalent bond. In respect of energy hydrogen bond has an intermediate position between covalent bonds and intermolecular van der Waals forces, based on dipole-dipole interactions, holding the neutral molecules together in gasses or liquefied or solidified gasses. Hydrogen bonding produces interatomic distances shorter than the sum of van der Waals radii, and usually involves a limited number of interaction partners. These characteristics become more substantial when acceptors bind H atoms from more electronegative donors. Hydrogen bonds hold H₂O molecules on 15 % closer than if water was a simple liquid with van der Waals interactions. The hydrogen bond energy compiles 5–10 kcal/mole, while the energy of covalent O–H-bonds in H₂O molecule – 109 kcal/mole. With fluctuations of water temperature the average energy of hydrogen H...O-bonds in H₂O molecule associates changes. That is why hydrogen bonds in liquid state are relatively weak and unstable: it is thought that they can easily form and disappear as the result of temperature fluctuations. The next conclusion that can be drawn from our research is that there is the distribution of energies among individual H₂O molecules.

Further we performed two types of temperature-dependent experiments on heat exchange from the surface of the human body by DNES-method. In first experiment we studied heat exchange when the temperature of the human body was higher than the temperature of the surrounding environment (curve 1a and 1b on Fig. 3). In second experiment there was heat exchange when the temperature of the human body was lower than that of the surrounding environment (curve 2a and 2b on Fig. 3). In both experiments it was detected a local maximum at $9.7 \mu\text{m}$ on curve 1 and curve 2 (Fig. 3). This local maximum corresponds to the maximal level of heat emission from the surface of the human body and lays within the “transparency window” of Earth atmosphere to electromagnetic radiation in the mid IR-range of the electromagnetic spectrum. In this range, the electromagnetic radiation emitted by the earth in the surrounding space is being absorbed by the Earth atmosphere. There is a statistical difference between the results of heat emission from the surface of the human body to the surrounding environment and back to the human body according to the *t*-criterion of Student at $p < 0.01$. The local maximum on curve 1a is detected at 7.3 eV^{-1} , while the local maximum on curve 2a – at 2.4 eV^{-1} (Fig. 3). The result of Dimitar Risimanski from him to environment is (-7.2 meV) . The result of Dimitar Risimanski from environment to him is (7.1 meV) . The difference is definite as effective energy is $(-7.5) - (7.5) = (-15.0) \text{ meV}$.




Fig. 3. Differential non-equilibrium energy spectrum (DNES) reflecting the heat exchange of the human body with surrounding environment

4. Conclusion

In frames of this research many types of NIR radiation (electromagnetic waves, infrared radiation, thermo radiation, and bioluminescence) emitted from the human body were studied and carefully scrutinized. The approaches and methods for detecting various types of radiation employed in this research as magnetography, chemiluminescence and color coronal gas discharge spectral analysis can find further application in many branches of applied science and medical diagnostics, while other methods as NES and DNES may be applied for studying the interaction of electromagnetic fields with water and structural studies. The research of Risimanski's bioinfluence was performed with two model systems. First is biophysical analysis on water spectrum. Second is biochemical analysis with 1 % solution of glucose. In the report are presented the phenomenal biophysical results of influence of Dimitar Risimanski (Bulgaria). The results are base for additional biochemical, biological and medical research projects.

References

- [Adey, 1981](#) – Adey, W.R. (1981). Tissue Interaction with Non-ionizing Electromagnetic Fields. *Physiol. Rev.*, 61, pp. 435–514.
- [Anderson, 1993](#) – Anderson, L.E. (1993). Biological Effect of Extremely Low Frequency Electromagnetic Fields: in Vivo Studies. *Am. Ind. Hyg. Assoc. J.*, 54, pp. 186–196.
- [Anosov, Trukhan, 2003](#) – Anosov V.N., Trukhan E.M. (2003). A New Approach to the Problem of Weak Magnetic Fields: An Effect on Living Objects. *Doklady Biochemistry and Biophysics*, 392 (1-6): 274-278.
- [Antonov, 1995](#) – Antonov, A. (1995). Research of the Non-equilibrium Processes in the Area in Allocated Systems. Dissertation thesis “Doctor of physical sciences”, Blagoevgrad, Sofia.
- [Antonov, Yuskesseliyeva, 1985](#) – Antonov, A., Yuskesseliyeva, L. (1985). Selective High Frequency Discharge (Kirlian effect). *Acta Hydrophysica*, Berlin, 5, 29 p.
- [Barnes, Greenebaum, 2006](#) – Barnes, F.S., Greenebaum, B. (eds.) (2006). CRC Handbook on Biological Effects of Electromagnetic Fields. 3d Edition. Boca Raton: CRC Press, Vol. 2, 960 p.
- [Bars, Andre, 1976](#) – Bars, Le., Andre, G. (1976). Biological Effects of Electric Fields on Rats and Rabbits. *Red. Gen. Elect.* (special issue), pp. 91–97.
- [Belousov et al., 1997](#) – Belousov, L.V., Opitz, J.M., Gilbert, S.F. (1997). Life of Alexander G. Gurwitsch and His Relevant Contribution to the Theory of Morphogenetic Fields. *The International journal of developmental biology*, 41(6): pp. 7–771.
- [Belousov et al., 2000](#) – Belousov, L., Popp, F.A., Voeikov, V., van Wijk, R. (eds) (2000). Biophotonics and Coherent Systems. *Moscow University Press*, 133 p.
- [Boveris et al., 1980](#) – Boveris, A., Cadenas, E., Reiter, R., Filipkowski, M., Nakase, Y., Chance, B. (1980). Organ chemiluminescence: Noninvasive assay for oxidative radical reactions. *Proc. Natl. Acad. Sci. USA*, Vol. 77, pp. 347–351.
- [Chang et al., 1998](#) – Chang, J.J., Fisch, J. & Popp, F.A. (eds) (1998). Biophotons. Dordrecht. *Kluwer Academic Publishers*, 417 p.
- [Chiang et al., 2005](#) – Chiang, L.H, Wah Khong, P. & Ghista, D. (2005). Bioenergy Based Medical Diagnostic Application Based on Gas Discharge Visualization. *Conf Proc IEEE Eng. Med. Biol. Soc.*, Vol. 2, 1533 p.
- [Choi et al., 2002](#) – Choi, C., Woo, W.M., Lee, M.B. at al. (2002). Biophoton Emission From the Hands. *J. Korean Physical. Soc.*, Vol. 41, pp. 275–278.
- [Cleary, 1993](#) – Cleary, S.F. (1993). A Review of in Vitro Studies: Low-frequency Electromagnetic Fields. *J. Am. Ind. Hyg. Assoc.*, Vol. 54(4), pp. 178–185.
- [Cohen, 1968](#) – Cohen, D. (1968). Magnetoencephalography: Evidence of Magnetic Fields Produced by Alpha-rhythm Currents. *Science*, Vol. 161(3843), pp. 784–786.
- [Cohen, Popp, 1997](#) – Cohen, S., Popp, F.A. (1997). Biophoton emission of the human body. *Journal of Photochemistry and Photobiology B: Biology*, 40(2): 187–189.
- [Devaraj et al., 1997](#) – Devaraj, B., Usa, M., Inaba, H. (1997). Biophotons: Ultraweak Light Emission from Living Systems. *Curr. Opin. Solid State Mater Sci.*, Vol. 2, 188.
- [Dobrin et al., 1979](#) – Dobrin, R., Kirsch, C., Kirsch, S. et al. (1979). Experimental Measurements of the Human Energy Field. In S. Krippner (ed.). *Psychoenergetic Systems: The Interface of Consciousness, Energy and Matter*. New York, Gordon & Breach, 230 p.

[Esterbauer et al., 1990](#) – Esterbauer, H., Zollner, H., Schaur, R. J. (1990). Aldehydes Formed by Lipid Peroxidation: Mechanisms of Formation, Occurrence, and Determination. In Membrane Lipid Oxidation. Boca Raton, CRC Press, 283 p.

[Frey, 1993](#) – Frey, A.H. (1993). Electromagnetic Field Interactions with Biological Systems. *FASEB Journal*, Vol. 7(2): pp. 272–281.

[Gerardi et al., 2008](#) – Gerardi, G., De Ninno A., Prosdocimi, M. et al. (2008). Effects of Electromagnetic Fields of Low Frequency and Low Intensity on Rat Metabolism. *Biomagnetic Research and Technology*, 6: 3.

[Goodman et al., 1995](#) – Goodman, R., Greenbaum, B. & Marron, M.T. (1995). Effects of Electromagnetic Fields on Molecules and Cells. *Int. Rev. Cytol.*, 158: 279–338.

[Gross, 1964](#) – Gross, B. (1964). Charge Storage in Solid Dielectrics; a Bibliographical Review on the Electret and Related Effects. New York, Elsevier Pub. Co., 230.

[Gubkin, 1978](#) – Gubkin, A.N. (1978). Electrets. Moscow, Nauka. 192 p.

[Gulyaev, Godik, 1984](#) – Gulyaev, Yu.V., Godik, E.E. (1984). On the Possibilities of the Functional Diagnostics of the Biological Subjects Via Their Temporal Dynamics of the Infrared Images. *USSR Academy Nauk Proceedings/Biophysics*, Vol. 277, pp. 1486–1491.

[Gulyaev, Godik, 1990](#) – Gulyaev, Yu.V., Godik, E.E. (1990). Human and Animal Physical Fields. // *Scientific American*, Vol. 5, pp. 74–83.

[Gulyaev, Godik, 1991](#) – Gulyaev, Yu.V., Godik, E.E. (1991). Functional Imaging of the Human Body. *IEEE Engineering in Medicine and Biology*, Vol. 10, pp. 21–29.

[Gurwitsch, 1959](#) – Gurwitsch, A.G. (1959). Die Mitogenetische Strahlung, Ihre Physikalisch-chemischen Grundlagen und Ihre Anwendung in Biologie und Medizin. Jena, Germany, Veb G. Fisher.

[Gurwitsch, 1988](#) – Gurwitsch, A.G. (1988). A historical Review of the Problem of Mitogenetic Radiation. *Experientia*, Vol. 44: 545–550.

[Halliwell, Gutteridge, 1989](#) – Halliwell, B., Gutteridge, J.M.C. (1989). Free Radicals in Biology and Medicine (2nd ed.). Oxford, Clarendon Press.

[Hastings, 1983](#) – Hastings, J.W. (1983). Biological Diversity, Chemical Mechanisms, and the Evolutionary Origins of Bioluminescent Systems. *J. Mol. Evol.*, 19(5): 309–321.

[Holzel, Lamprecht, 1994](#) – Holzel, R., Lamprecht, I. (1994). Wirkungen Elektromagnetischer Felder auf biologische Systeme. *Nachrichtentech Elektron.* Vol. 44(2), pp. 28–32.

[Ignatov et al., 1998](#) – Ignatov, I., Antonov, A., Galabova, T. (1998). Medical Biophysics – Biophysical Fields of Man. *Gea Libris, Sofia*, pp. 1–71.

[Ignatov et al., 2002](#) – Ignatov, I., Antonov, A., Galabova, T. (2002). Scientific Research Studies with Christos Drossinakis (October 2001 – October 2002). *Int. Conference “Man and Nature”*, SRCMB, Sofia.

[Ignatov et al., 2014](#) – Ignatov, I., Mosin, O. V., Stoyanov, Ch. (2014). Fields in Electromagnetic Spectrum Emitted from Human Body. Application in Medicine. *Journal of Health, Medicine and Nursing*, Vol. 7, pp. 1-22.

[Ignatov et al., 2014a](#) – Ignatov, I., Mosin, O. V., Niggli, H., Drossinakis, Ch. (2014). Evaluating of Possible Methods and Approaches for Registering of Electromagnetic Waves Emitted from the Human Body. *Advances in Physics Theories and Applications*, Vol. 30, pp. 15-33

[Ignatov et al., 2014b](#) – Ignatov, I., Mosin, O.V., Drossinakis, Ch. (2014). Infrared Thermal Field Emitted from Human Body. Thermovision. *Journal of Medicine, Physiology, Biophysics*, Vol.1, pp. 1-12.

[Ignatov et al., 2014c](#) – Ignatov, I., Mosin, O. V., Niggli, H., Drossinakis, Ch., Stoyanov, Ch. (2014). Registration of Electromagnetic Waves Emitted the Human Body. *Journal of Medicine, Physiology and Biophysics*, Vol. 5, pp. 1-22.

[Ignatov et al., 2014d](#) – Ignatov, I., Mosin, O. V., Niggli, H., Drossinakis, Ch. (2014). Evaluating of Possible Methods and Approaches for Registering Electromagnetic Waves Emitted from the Human Body. *Nanotechnology Research and Practice*, Vol. 2 (2), pp. 96-116.

[Ignatov et al., 2014e](#) – Ignatov, I., Mosin, O.V., Stoyanov, Ch. (2014). Biophysical Fields. Color Coronal Spectral Analysis. Registration with Water Spectral Analysis. Biophoton Emission. *Journal of Medicine, Physiology and Biophysics*, Vol. 6, pp. 1-22.

[Ignatov et al., 2015](#) – Ignatov, I., Mosin, O.V., Niggli, H., Drossinakis, Ch. (2015). Evaluation of Possible Methods and Approaches for Registering Non-Ionizing Radiation (NIR) Waves Emitting from the Human Body. *Journal of Pharmacy and Alternative Medicine*, Vol. 4, pp. 57-75.

[Ignatov et al., 2016](#) – Ignatov, I. et al. (2016). Results of Biophysical and Nano Technological Research of ZEOLITH Detox of LavaVitae Company. *Journal of Health, Medicine and Nursing*, Vol. 30, pp. 44-49.

[Ignatov, 2005](#) – Ignatov, I. (2005). Energy Biomedicine. *Gea-Libris, Sofia*: pp. 1–88.

[Ignatov, 2016a](#) – Ignatov, I. (2016). Product of LavaVitae BOOST is Increasing of Energy of Hydrogen Bonds among Water Molecules in Human Body. *Journal of Medicine, Physiology and Biophysics*, Vol. 27., pp. 30-42.

[Ignatov, 2016b](#) – Ignatov, I. (2016). VITA intense – Proofs for Anti-inflammatory, Antioxidant and Inhibition Growth of Tumor Cells Effects. Relaxing Effect of Nervous System. Anti Aging Influence. *Journal of Medicine, Physiology and Biophysics*, Vol. 27. pp. 43-61.

[Ignatov, 2017](#) – Ignatov, I. (2017). Aluminosilicate Mineral Zeolite. Interaction of Water Molecules in Zeolite Table and Mountain Water Sevtopolis from Bulgaria. *Journal of Medicine, Physiology and Biophysics*, Vol. 31, pp. 41-45.

[Ignatov, 2017a](#) – Ignatov, I. (2017). VITA intense and Boost – Products with Natural Vitamins and Minerals for Health. *Journal of Medicine, Physiology and Biophysics*, Vol. 31, pp. 58-78.

[Ignatov, 2017b](#) – Ignatov, I. (2017). ZEOLITH detox for Detoxification and ZEOLITH Creme for Skin Effects as Products of LavaVitae Company. *Journal of Medicine, Physiology and Biophysics*, Vol. 31, pp. 79-86.

[Ignatov, Mosin, 2014](#) – Ignatov, I., Mosin, O. V. (2014). Coronal Gas Discharge Effect in Modeling of Non-Equilibrium Conditions with Gas Electric Discharge Simulating Primary Atmosphere and Hydrosphere for Origin of Life and Living Matter, *Journal of Medicine, Physiology and Biophysics*, 5: 47-70.

[Ignatov, Mosin, 2012](#) – Ignatov, I., Mosin, O.V. (2012). Coronal Effect in Biomedical Diagnostics and Study of Bioenergetical Properties of Biological Objects and Water. *Biomedical Radio electronics, Biomedical Technologies and Radio Electronics*, Vol. 12, pp. 13–21 [in Russian].

[Ignatov, Mosin, 2013a](#) – Ignatov, I., Mosin, O.V. (2013). Method for Color coronal (Kirlian) Spectral Analysis. *Biomedical Radio electronics, Biomedical Technologies and Radio Electronics*: Vol. 1, pp. 38–47 [in Russian].

[Ignatov, Mosin, 2013b](#) – Ignatov, I., Mosin O.V. (2013b). Color Crown Spectral Kirlian Analysis in the Modeling of Non-equilibrium Conditions with a Gas electric Discharge that Simulates the Primary Atmosphere. *Nano engineering*, Vol. 12(30), pp. 3–13 [in Russian].

[Ignatov, Mosin, 2013c](#) – Ignatov, I., Mosin, O.V. (2013c). Structural Mathematical Models Describing Water Clusters. *Journal of Mathematical Theory and Modeling*, 3(11), 72–87.

[Ignatov, Mosin, 2014](#) – Ignatov, I., Mosin, O. V. (2014). The Methods for Studying the Structure of Water Clusters (H₂O), where n=3-20, Application in Medicine. *Journal of Health, Medicine and Nursing*, Vol. 7, pp. 23-52.

[Ignatov, Mosin, 2014a](#) – Ignatov, I., Mosin, O.V. (2014). Mathematical Models of Distribution of Water Molecules Regarding Energies of Hydrogen Bonds. *Journal of Medicine, Physiology and Biophysics*, 2: 71-94.

[Ignatov, Mosin, 2014b](#) – Ignatov, I., Mosin, O.V. (2014). Mathematical Models Describing Water Clusters as Interaction among Water Molecules. Distributions of Energies of Hydrogen Bonds. *Journal of Medicine, Physiology and Biophysics*, Vol. 3, pp. 48-70.

[Ignatov, Mosin, 2014c](#) – Ignatov, I., Mosin, O. V. (2014). The Methods for Studying the Structure of Water Clusters (H₂O), where n=3-20. Water Clusters as Nano-structures. *Journal of Health, Medicine and Nursing*, Vol. 8, pp. 29-58.

[Ignatov, Mosin, 2016](#) – Ignatov, I., Mosin, O.V. (2016). Deuterium, Heavy Water and Origin of Life. LAP LAMBERT Academic Publishing, pp. 1-500.

[Ignatov, Mosin, 2016a](#) – Ignatov, I., Mosin, O.V. (2016). Water for Origin of Life Altaspera Publishing & Literary Agency Inc., pp. 1-616.

[Ignatov, Mosin, 2016b](#) – Ignatov, I., Mosin, O.V. (2016). Results of Bioinfluence of Dimitar Risimanski with Biophysical Model Systems, *Journal of Medicine, Physiology and Biophysics*: Vol. 24, pp. 5-17.

Ignatov, Mosin, 2016c – Ignatov, I., Mosin, O.V. (2016). Biophysical Results of Bioinfluence of Dimitar Risimanski as Base of Medical Effects. *Journal of Health, Medicine and Nursing*: Vol. 27, pp. 24-35.

Ignatov, Mosin, 2016d – Ignatov, I., Mosin, O.V. (2016). Structural Alterations among Water Molecules after Bioinfluence of Dimitar Risimanski. *Advances in Physics Theories and Applications*, Vol. 57, pp. 20-44.

Ignatov, Tsvetkova, 2011 – Ignatov, I., Tsvetkova, V. (2011). Water for the Origin of Life and “Informationability” of Water, Kirlian (Electric Images) of Different Types of Water. *Euromedica, Hanover*, pp. 62-65.

Inaba, 1988 – Inaba, H. (1988). Super-high Sensitivity Systems for Detection and Spectral Analysis of Ultraweak Photon Emission from Biological Cells and Tissues. *Experientia*, Vol. 44, pp. 550–559.

Inaba, 2000 – Inaba, H. (2000). Measurement of biophotons from human body. *J. Int. Soc. Life Inf. Sci.*, 18: 448.

Kiang et al., 2005 – Kiang, J.G., Ives J.A., Jonas, W.B. (2005). External Bioenergy-induced Increases in Intracellular Free Calcium Concentrations are Mediated by $\text{Na}^+/\text{Ca}^{2+}$ Exchanger and L-type Calcium Channel. *Mol. Cell Biochem.*, Vol. 271, 51 p.

Kim, 2002 – Kim, T.J. (2002). Biophoton Emission from Fingernails and Fingerprints of Living Human Subjects. *Acupuncture Electrother Res.*, Vol. 27, 85 p.

Kirlian, 1949 – Kirlian, S.D. (1949). Method for Receiving Photographic Pictures of Different Types of Objects. USSR Patent № 106401.

Kwan-Hoong, 2003 – Kwan-Hoong, Ng. (2003). Non-ionizing radiations – Sources, Biological Effects, Emissions and Exposures. *Proceedings of the International Conference on Non-Ionizing Radiation at UNITEN (ICNIR2003). Electromagnetic Fields and Our Health*. 20–22 October 2003.

Liboff et al., 1984 – Liboff, A.R., Williams, T., Strong, D.M., Wistar, R. (1984). Time-varying Magnetic Fields: Effect on DNA Synthesis. *Science*, Vol. 223, pp. 818–820.

Marino, 1988 – Marino, A.A. (Ed.) (1988). *Modern Bioelectricity*. Mariele Dekker, New York, Basel.

Marinov, Ignatov, 2008 – Marinov, M., Ignatov, I. (2008). Color Kirlian Spectral Analysis. Color Observation with Visual Analyzer. *Euromedica, Hanover*, pp. 57-59.

Marinov, Ignatov, 2008 – Marinov, M., Ignatov, I. (2008). Color Kirlian Spectral Analysis. Color Observation with Visual Analyzer. *Euromedica, Hanover*, pp. 57-59.

Miller, 1986 – Miller, M.W. (1986). Extremely Low Frequency (ELF) Electric Fields: Experimental Work on Biological Effects. *CRC Handbook of biological effects of electromagnetic fields*, 138–168.

Mosin, 2011 – Mosin, O.V. (2011). Magnetic Devices for Water Treatment. *C.O.K. Publishing House “Media Technology” (Moscow)*, Vol. 6: 24–27 [in Russian].

Mosin, 2012 – Mosin, O.V. (2012). Advanced Technologies and Equipment for Magnetic Water Treatment (review). *Water Supply and Sanitary Technique*, Vol. 8: 12–32 [in Russian].

Mosin, Ignatov, 2015 – Mosin, O.V., Ignatov, I. (2015). An Overview of Methods and Approaches for Magnetic Treatment of Water. *Water: Hygiene and Ecology*, Vol. 3-4 (4): pp. 113-130.

Motohiro, 2004 – Motohiro, T. (2004). Biophoton Detection as a Novel Technique for Cancer Imaging. *Cancer Science*, Vol. 95(8), pp. 656–661.

Niggli, 1993 – Niggli, H. (1993). Artificial Sunlight Irradiation Induces Ultra Weak Photon Emission in Human Skin Fibroblasts. *Journal of Photochemistry and Photobiology B: Biology*, Vol. 18 (2–3), pp. 281–285.

Nikolaev, 2000 – Nikolaev, Y.A. (2000). Distant Interactions in Bacteria. *Microbiology*, Vol. 69(5), pp. 497–503.

Pehek et al., 1976 – Pehek, J.O., Kyler, H.J., Faust, D.L. (1976). Image Modulating Corona Discharge Photography. *Science*, Vol. 194(4262), pp. 263–270.

Popp et al., 1992 – Popp, F.A., Li, K., Gu, Q. (1992). Recent Advances in Biophoton Research and its Application, *World scientific*, 1–18.

Popp et al., 1994 – Popp, F.A., Quao, G. & Ke-Hsuen, L. (1994). Biophoton Emission: Experimental Background and Theoretical Approaches. *Modern physics Letters B.*, Vol. 8, pp. 21–22.

[Popp et al., 2002](#) – Popp, F.A., Chang, J.J., Herzog, A., Yan, Z. & Yan, Y. (2002). Evidence of Non-classical (squeezed) Light in Biological Systems. *Physics Letters A*, Vol. 293(1–2), pp. 98–102.

[Popp, 2005](#) – Popp, F.A. (2005). Essential Differences Between Coherent and Non-coherent Effects of Photon Emission from Living Organisms. In: X. Shen, R. van Wijk (eds). *Biophotonics*. New York: Springer, 124 p.

[Porter, Wujek, 1988](#) – Porter, N.A., Wujek, D.G. (1988). Reactive Oxygen Species in Chemistry, Biology, and Medicine. In A. Quintanilha, Ed. New York, Plenum Press, pp. 55–79.

[Rattemeyer et al., 1981](#) – Rattemeyer, M., Popp, F.A. & Nagl, W. (1981). Evidence of Photon Emission from DNA in Living Systems. *Nature Wissenshanften*, Vol. 68(11), pp. 572–573.

[Rauhut, 1985](#) – Rauhut, M.M. (1985). Chemiluminescence. In: M. Grayson (Ed). Kirk-Othmer Concise Encyclopedia of Chemical Technology (3rd ed). New York, John Wiley and Sons. 247 p.

[Ring, Hughes, 1986](#) – Ring, E.F.J., Hughes, H. (1986). Real Time Video Thermography in Recent Developments in Medical and Physiological Imaging. *Suppl. Journal of Medical Engineering and Technology*, pp. 86–89.

[Rubik, 2002](#) – Rubik, B. (2002). The biofield hypothesis: its biophysical basis and role in medicine. *J. Altern. Complement Med.*, Vol. 8(6), pp. 703–717.

[Sessler, Gerhard-Multhaupt, 1998](#) – Sessler, G.M., Gerhard-Multhaupt, R. (eds) (1998). Electrets. *Laplacian Press, Morgan Hill, California, USA*.

[Seto et al., 1992](#) – Seto, A., Kusaka, C. Nakazato, S. et al. (1992). Detection of Extraordinary Large Bio-magnetic Field Strength from Human Hand. *Acupuncture Electrother Res. Int. J.*, Vol. 17, 75 p.

[Shimizu et al., 1995](#) – Shimizu H., Suzuki, Y., Okonogi, H. (1995). Biological Effects of Electromagnetic Fields. *Nippon Eiseigaki Zasshi.*, Vol. 50(6), pp. 919–931.

[Sisodia, 2007](#) – Sisodia, M.L. (2007). Microwaves: Introduction to Circuits, Devices and Antennas. *New Delhy, New Age International Ltd.*, 602 p.

[Vladimirov, 1996](#) – Vladimirov, Y.A. (1996). Studies of Antioxidants with Chemiluminescence. In: *Proceedings of the International Symposium on Natural Antioxidants. Molecular Mechanisms and Health Effects*. L. Packer, M.G. Traber & W. Xin (eds.), pp. 125–144.

[Wikswow, Barach, 1980](#) – Wikswow, J., Barach, J. (1980). An Estimation of the Steady Magnetic Field Strength Required to Influence Nerve Condition. *IEEE Trans. Bio-Med. Eng.*, Vol. 27, pp. 722–723.

[Young, Roper, 1976](#) – Young, R.E., Roper, C.F. (1976). Bioluminescent Countershading in Midwater Animals: Evidence from Living Squid. *Science*, Vol. 191(4231), pp. 1046–1048.

[Zhadin, 2001](#) – Zhadin, M.N. (2001). Review of Russian Literature on Biological Action of DC and Low-frequency AC magnetic fields. *Bioelectromagnetics*, Vol. 22, pp. 27–45.

[Zlatkevich, Kamal-Eldin, 2005](#) – Zlatkevich, L., Kamal-Eldin, A. (2005). Analysis of Lipid Oxidation. In: A. Kamal-Eldin & J. Pokorn (Eds.). New York, AOCS Publishing, 281 p.