

Pharmaceutically important uses of *Chameli* (*Jasminum grandiflorum* Linn): A Review

Shailendra Pradhan*

*Department of Dravyaguna, Uttarakhand Ayurved University, Gurukul Campus, Haridwar, Uttarakhand, India

Abstract

Jasminum grandiflorum Linn (Chameli / Yasmin; Oleaceae) is innate to Tropical and warm Temperate regions and cultivated in France, Italy, China, Japan, India, Morocco and Egypt. The plant is recognised to possess positive effects as odontalgic, thermogenic, aphrodisiac, antiseptic, emollient, anthelmintic, deobstruant, suppurative, tonic, in fixing loose teeth, ulcerative stomatitis, leprosy, skin diseases, otorrhoea, otalgia, wounds, corns and aromatherapy. Pharmacological actions of the plant reported so far are spasmolytic, anti-inflammatory, anti-microbial, antioxidant, antiulcer, cytoprotective, chemoprotective, wound healing and anti-acne activity. The present review is an attempt to highlight the various ethnobotanical and traditional uses as well as phytochemical and pharmacological actions reported so far from *J. grandiflorum*.

Keywords

Jasminum grandiflorum , Chameli, Yasmin, Oleaceae



Greentree Group

Received 15/5/15 Accepted 19/6/15 Published 10/7/15

INTRODUCTION

Jasminum grandiflorum is a large scrambling sub erect twining evergreen shrub, which grows up to 10 to 15 m. high^{1, 2}. It is native of Asia, Kashmir, Afghanistan and Persia ascending to an altitude of 700- 2700 m, cultivated in India, wild in sub-tropical North-West Himalayas, Western Ghats, Nilgiris, hill of Tinnavally above 1400 m, France, Italy, China, Japan, India, Morocco and Egypt^{3,4,5,6}. The plant is recognised to possess beneficial effects as odontalgic, thermogenic, aphrodisiac, antiseptic, emollient, anthelmintic, deobstruant, suppurative, tonic, in fixing loose teeth, ulcerative stomatitis, leprosy, skin diseases, otorrhoea, otalgia, wounds, corns and aromatherapy⁷. Our thorough literature search exposed an interesting fact that though the plant is a popular remedy for a variety of ailments, very little effort have been made to verify its efficacy through scientific screenings in animal model and clinical trials. The present review highlights the various folk, Ayurvedic uses, pharmacognostical, phytochemical and pharmacological studies conducted on *J. grandiflorum* and also highlight unexplored

potential of the plant.

Taxonomical/ Scientific Classification⁸

Kingdom: Plantae- Plants

Subkingdom: Tracheobionts- Vascular plants

Division: Magnoliophyta- Flowering plants

Class: Magnoliopsida- Dicotyledons

Order: Scrophulariales

Family: Oleaceae

Genus: *Jasminum*

Species: *grandiflorum*

Classical Names^{9, 10}

Jati, Sauanasyayani, Sumama, Chetika,

Hridyagandha, Malati, Rajaputrika.

Botanical Description^{11, 12}

It is a scrambling deciduous shrub growing to 2-4 m tall. The leaves are opposite, entire ovate to somewhat elliptic in shape with acuminate mucronate apex, petiole almost lacking, imparipinnately compound, with three paired foliates ending with a single leaf at the tip. The leaflets are elongate-lanceolate, acute, 7 to 11 terminal leaflet somewhat large than laterals, narrowing at the base, ovate-lanceolate, acute or acuminate, laterals ovate, terminal one larger than laterals and often partially

united with surfaces with a ciliate margin. Flowers are terminal and axillary cymes, calyx lobes long and linear, more than half as long as the corolla tubes. The fruit is a black berry, elliptic, globose berries when ripe.

Climate, Soil and Propagation

The plant is cultivated in well drained loamy soil and also on a variety of soils such as black, lateritic and clay loam with good drainage system as the plant is highly susceptible to water logging^{13, 14}. It can be propagated by shoot tip culture method. Flowering of jasmine plants starts in the first year itself. The yield being ½, 5 and 10 tonnes/ha flowers within first, second and third year respectively^{15, 16}. The harvesting of the flower is done in the month of May to December (in South India) and July to November (in North India).¹⁷

Pharmacognostical Studies

Leaf Powder Characteristics

The leaf powder is light green in color. The important powder characteristics of the leaf are simple, unbranched trichomes, vessels with simple pits and spiral thickenings, glandular trichome, starch grains and calcium oxalate crystals^{18, 19}.

The physical constants⁶ of the plant are given in the Table 1.

Medicinal Uses

1. Traditional Uses⁷

Plant parts used: Whole plant

The plant is bitter, astringent, acrid, thermogenic, aphrodisiac, antiseptic, anodyne, depurative, emmenagogue, emollient, diuretic, anthelmintic, deobstruant, dentrifice, suppurative and tonic.

Roots: They are useful in cephalalgia, vitiated condition of vata, paralysis, facial paralysis, mental debility, chronic constipation, flatulence, strangury, sterility, dysmenorrhoea, amenorrhoea, ringworm, leprosy, skin diseases and giddiness.

Leaves: They are useful in odontalgia, fixing loose teeth, ulcerative stomatitis, leprosy, skin diseases, otorrhoea, otalgia, strangury, dysmenorrhoea, ulcers, wound and corns.

Flowers: They are useful in stomatopathy, cephalopathy, odontopathy, ophthalmopathy, leprosy, skin diseases, pruritis, strangury, dysmenorrhoea, ulcers, as refrigerant, ophthalmic and vitiated conditions of pitta.

Table 1 Physical Constants of *Jasminum grandiflorum* Linn

S. No.	Physical constant	% w/w
1.	Total ash	10.89
2.	Acid insoluble ash	1.29
3.	Water soluble ash	2.92
4.	Loss on drying	4.25
Petroleum ether extractive		
5.	value	2.61
Chloroform extractive		
6.	value	3.58
7.	Acetone extractive value	8.72
8.	Alcohol extractive value	11.57
9.	Water extractive value	12.14

2. Ethnobotanical Uses

Asian and Indian folk practitioners recommend Jasmine for liver complaints, dysentery, various types of pain including painful menstruation, and skin diseases such as leprosy. In addition, Jasmine oil applied externally is used to soften and smooth the skin, for cancer, heart disease, and a variety of other ills. Aroma therapists believe Jasmine oil can be useful as an antidepressant, as a calming agent to soothe stress, pain, and anxiety, and as an aphrodisiac. Its reputation as an intoxicant is legendary. Apart from that inhaling Jasmine scent increases beta waves in the brain, which are associated with increased states of alertness¹⁰.

3. Uses described in Ayurveda

Leaves are chewed in aphthae, stomatitis, toothache, ulcer in the mouth and leaf-juice or oil obtained from it is dropped in to the ear (Bhavaprakash²⁰). A decoction of the leaf was also used as a gargle (Bhavaprakash and Vrindamadhava²¹). The oil cooked with juice of jati leaves was prescribed in purulent discharge from the ear (Vrindamadhava and Bangasena²¹). Fresh juice of the leaves is a valuable application for soft corns between the toes, for ulceration in the mouth, throat and gums, the leaves fried in ghee are recommended to be applied (Chakradatta²⁰). The use of flowers applied as a plaster to the loins, genitals and pubes as an aphrodisiac. The plant is used in scorpion-string (Mahomedan²²). Charaka used the sprouts or dried flowers, in prescriptions, externally in coryza, nasal hemorrhage and dermatosis. Sushruta used Malati as an ingredient of a medicated clarified butter for external application on infected wounds, for cleansing and sterilizing the interior of ulcer, as an ingredient of hair oil for baldness and alopecia and as an ingredient of an eye-salve for loss of vision. Malati was used externally in leprosy, malignant ulcers and

other virulent skin diseases. The root of Jaati, cooked in goat's milk and mixed with sugar was prescribed for giving relief in pain due to retention of urine and for expelling calculus (Raaja Maarttanda²⁰).

Ayurvedic Properties^{6, 23}:

Rasa: Tikta, Kashaya

Guna: Laghu, Snigdha, Mridu

Veerya: Ushna

Vipaka: Katu

Doshagnata: Tridoshashamaka

Rogagnata: Tridoshajavikara, Dantashoola,

Dantadaurbalya, Mukharoga.

Karma: Mukharoganashaka,

Saumanasyajanana, Medhya, Vajikarana.

Phytochemical studies

Very little phytochemical work has been carried out with the plant *J. grandiflorum*^{24, 25, 26}.

Leaves: 2''-epifraxamoside, demethyl-2''-epifraxamoside, jasminanhydride²⁷, oleacein, 2-(3,4-dihydroxy phenyl)-ethanol, isoquercitrin, ursolic acid²⁸, resin, salicylic acid, jasminine, indole oxygenase²⁹, 3,4-dihydroxy benzoic acid, 2-hydroxy-30, 40-dihydroxyacetophenone and oleanolic acid²⁷.

Flowers: Cis-3-hexenol, 2-vinyl pyridine,

indole, myrcene, linalool, geranyl linalool, α -terpineol, geraniol, linalyl acetate, nerolidol, phytol, isophytol, farnesol, eugenol, benzyl alcohol, p-cresol, methyl benzoate, benzyl cyanide, benzyl acetate, methyl dihydrojasmonate, methyl anthranilate, jasmone, methyl-N-methyl anthranilate, vanillin, cis-3-hexenyl benzoate, benzyl benzoate, methyl palmitate, methyl linoleate²⁵, jasgranoside, jaspolyoside, 8-epi-kingiside, 10-hydroxy-oleuropein, 10-hydroxy ligstroside, oleoside-7,11-dimethyl ester³⁰, 3-O- α -L-rhamnopyranosyl (1 \rightarrow 2)- β -D-xylopyranosyl-hederagenin-28-O- β -D-galactopyranosyl (1 \rightarrow 6)- β -D-galactopyranosyl ester, hederagenin-3-O- β -D-glucopyranosyl (1 \rightarrow 3)- α -L-arabinopyranoside, 2- α ,3 β ,23-trihydroxyolean-12-en-28-oic -O- β -D-glucopyranosyl ester, hederagenin-3-O- β -D-xylopyranosyl (1 \rightarrow 3)- α -L-rhamnopyranosyl (1 \rightarrow 2)- α -L-arabinopyranoside, 2 α ,3 β ,23-trihydroxyolean-12-en-28-oic -O- α -L-rhamnopyranosyl (1 \rightarrow 4)- β -D-glucopyranosyl (1 \rightarrow 6)- β -D-glucopyranosyl ester, hederagenin-3-O- α -L-rhamnopyranosyl (1 \rightarrow 2)- α -L-arabinopyranoside³¹, kaempferol-3-O- α -L-rhamnopyranosyl (1 \rightarrow 3)-[α -L-

rhamnopyranosyl (1→6)-β-D-galactopyranoside, kaempferol-3-O-rutinoside, 7-ketologanin, oleoside-11-methyl ester, 7-glucosyl-11-methyl oleoside, ligstroside and oleuropein³².

Jasmine oil: Methyl jasmonate²⁴, benzyl benzoate, linalool, linalyl acetate, benzyl alcohol, indole, jasmone, methyl anthranilate, P-cresol, geraniol, racemic (5-pent-2-enyl)-5,1-pentanolid, benzyl benzoate, nerol, 1-α-terpineol, d and dl-linalool, γ-jasmolactone, farnesol, nerolidol and eugenol^{6, 26}.

Pharmacological Activities

Spasmolytic activity

Jasmine has spasmolytic activity on guinea pig ileum (post synaptic and not atropine-like) and rat uterus *in vitro*. The spasmolytic effect of Jasmine absolute was most likely to be mediated through cAMP, and not through cGMP. The contradictory effect *in vitro* and *in vivo* has been suggested probably due to the solely physiological effects of jasmine absolute *in vitro* (producing a relaxation) compared with that *in vivo*, where it has a strong psychological input, producing a stimulant effect in man and enhanced movement in animals³³.

Anti-inflammatory activity

Topical anti-inflammatory activity of a polyherbal formulation, Jatyadi ghrita, consists of

Jasminum officinale, *Azadirachta indica*, *Berberis aristata*, *Curcuma longa*, *Picrorrhiza kurroa*, *Rubia cordifolia*, *Trichosanthes dioica*, *Aristolochia indica*, *Hemidesmus indicus*, *Randia spinosa* and *Glycyrrhiza glabra* has been evaluated. The preparation showed nearly 50 percent inhibition of croton oil induced ear edema when compared to Diclofenac sodium, which showed 33 percent inhibition³⁴.

Antimicrobial activity

The antimicrobial activity of ethanol callus extracts of two species of *Jasminum*, (*J. grandiflorum* and *J. sambac*) were evaluated. Preliminary phytochemical analysis of the callus extracts revealed the presence of alkaloids, glycoside, flavanoid, terpenes, tannin, resin, and salicylic acid. The extracts were subjected for screening of *in-vitro* antimicrobial activity against selected disease causing pathogens, viz., *Staphylococcus albus*, *Proteus mirabilis* and *Salmonella typhii*, at the concentrations of 500 mg/ml and 250 mg/ml. The results of antimicrobial activity revealed that all

the extracts showed significant antibacterial activity³⁵.

Antiulcer and antioxidant activities

The antiulcer and antioxidant activities of 70% ethanolic extract of leaves of *J. grandiflorum* (JGLE) were evaluated. Antiulcerogenic activity of JGLE (100 and 200 mg/kg, b.w. orally) was evaluated employing aspirin + pylorus ligation (APL) and alcohol (AL) induced acute gastric ulcer models and ulcer-healing activity using acetic acid induced (AC) chronic ulcer model in rats. The antioxidant activity of JGLE has been assayed by using *in vitro* methods like 2,2-diphenyl-1-picrylhydrazylhydrate (DPPH) assay, reductive ability, superoxide anion scavenging activity, nitric oxide scavenging activity and total phenolic content, in order to explain the role of antioxidant principles in the antiulcerogenic activity of the extract. There was a significant dose-dependent decrease in the ulcerative lesion index produced by all the three models in rats as compared to the standard drug famotidine (20 mg/kg, b. w. orally). The reduction in gastric fluid volume, total acidity and an increase in the pH of the gastric fluid in APL rats proved the antisecretory activity

of JGLE. Additionally, JGLE completely healed the ulcer within 20 days of treatment in AC model as evidenced by histopathological studies. The free radical scavenging activities of JGLE depends on concentration and increased with increasing amount of the extract. These results suggest that leaves of *J. grandiflorum* possess potential antiulcer activity, which may be attributed to its free radical scavenging activity³⁶.

Cytoprotective activity

The cytoprotective effects of *J. grandiflorum* flowers and leaves in 7,12-dimethylbenz (a) anthracene (DMBA) induced chromosomal abnormalities in bone marrow of female wistar rats were evaluated. Oral pretreatment of *J. grandiflorum* flower and leaf extracts to DMBA treated rats significantly reduced the frequency of micronucleated polychromatic erythrocytes in the rat bone marrow. Also, the plant extracts significantly decreased the percentage of aberrant cells; the number of chromatic and chromosomal breaks in DMBA treated rats which proves its cytoprotective effect³⁷.

Chemopreventive and Lipid peroxidative activities

The chemopreventive efficacy and anti-lipid peroxidative potential of *J. grandiflorum* Linn. on 7,12-dimethylbenz (a) anthracene (DMBA)-induced rat mammary carcinogenesis were evaluated. Oral administration of ethanol extract of *J. grandiflorum* flowers (JgEt) at a dose of 300 mg/kg body weight for 14 weeks to DMBA-injected animals completely prevented the formation of tumors in the pre-initiation period. JgEt also exerted significant anti-lipid peroxidative effect and improved the antioxidant defense system in DMBA-treated rats suggesting clearly that JgEt has potent chemopreventive efficacy³⁸.

Breast cancer

Flowers of *J. grandiflorum* are useful to women when brewed as a tonic as it aids in preventing breast cancer and stopping uterine bleeding³⁹.

Wound healing activity

The effect of flower extract of *J. grandiflorum* was studied for its wound healing activity at a dose of 250 mg/kg body weight orally for 10 days using excision and dead space wound models in rats. Extract treated rats exhibited 65%

reduction in the wound area when compared to controls (54%). The wet and dry granulation tissue weight, and hydroxyproline content in a dead space wound model were increased significantly when compared to controls. Histological studies of the tissue obtained on day 10 from the extract-treated group showed increased well organized bands of collagen, more fibroblasts and few inflammatory cells when compared to controls which showed inflammatory cells, scanty collagen fibers and fibroblasts which suggests the use of *J. grandiflorum* flower extract in the management of wound healing⁴⁰.

Anti-acne activity

The anti-acne activity of ten natural products being used as traditional medicine in various skin disorders has been investigated against *Propionibacterium acnes* by broth dilution method.

Minimal inhibition concentrations (MIC) of *J. grandiflorum* extract was found to be below 800 µg/ml⁴¹.

Angiotensin Converting Enzyme (ACE)

Inhibitor Activity

Bioactivity guided fractionation of extract of aerial parts of *J. grandiflorum* led to the

isolation of oleacein. The IC_{50} values of purified ACE inhibitor were between 26-66 mM^{28} .

Clinical Evaluation

A clinical trial was conducted to assess the effect of Jatyadi taila of which *J. grandiflorum* was one of the main ingredients. Group A (8 patients of eczema) were treated with Raktashodhaka vati (2 tab t.d.s), Surakta strong syrup (2 tsf b.d.), Panchatikta ghrita guggulu (15mg b.d.) and external application of Marichyadi taila and group B (n=8) were administered the same schedule as group A except that Jatyadi taila was applied externally instead of Marichyadi taila. The duration of treatment was continued for 3 months and reviewed after 4 weeks. In group A, only 37.5 % patients were cured whereas group B cured 62.5 % patients which suggests that Jatyadi taila was more effective in eczema patients⁴².

CONCLUSION

In recent years, ethnobotanical and traditional uses of natural compounds, especially of plant origin received much attention as they are well tested for their efficacy and generally believed to be safe

for human use. They obviously deserve scrutiny on modern scientific lines such as phytochemical investigation, biological evaluation on experimental animal models, toxicity studies and investigation of molecular mechanism of actions of isolated phytoconstituents. Thorough screening of literature available on *J. grandiflorum* depicted the fact that it is a popular remedy among the various ethnic groups, Vaidyas, Hakims and Ayurvedic practitioners for cure of variety of ailments. Following the traditional and folk claims, very little efforts have been made by the researchers to explore the therapeutic potential of this plant. From the literature, *J. grandiflorum* have been screened for some pharmacological activities and found to possess spasmolytic, anti-inflammatory, antimicrobial, antiulcer, antioxidant, cytoprotective, chemopreventive, breast cancer, wound healing and anti-acne activities but number of other pharmacological activities are yet to be explored. In future study, the isolated principles from Chameli needs to be evaluated in scientific manner using specific experimental animal models and clinical trials to understand the molecular mechanism of action, in search of lead molecule from natural resources.

Acknowledgement

The author wish to thank Dr. Mithilesh Singh, Scientist C, GBPIHED, Sikkim Unit, Gangtok for the assistance provided for this review.

REFERENCES

1. Anonymous. The Wealth of India A Dictionary of Indian Raw Materials and Industrial Products. Publication & Information Directorate, New Delhi: CSIR, 2004: 284-88.
2. Anonymous. Medicinal Plants of India. New Delhi: ICMR, 1987: 96-101.
3. Chopra RN, Chopra IC, Handa KL, Kapur LD. Indigenous Drugs of India. Calcutta: U N Dhur & Sons Pvt Ltd, 1958: 512.
4. Chopra RN, Nayar SL, Chopra IC. Glossary of India Medicinal Plants. New Delhi: CSIR, 2002: 143-145.
5. Kirtikar KR, Basu BD. Indian Medicinal Plants. Allahabad: L M Basu, 1989:1522-1524.
6. Sharma PC, Yelne MB, Dennis TJ. Database on Medicinal Plants used in Ayurveda. Vol. 3. New Delhi: Central Council for Research in Ayurveda and Siddha, 2005: 332-345.
7. Warriar PK, Nambiar VPK, Ramankutty. Indian Medicinal Plants- a Compendium of 500 Species. Vol. 3. Chennai: Orient Longman Pvt Ltd, 2004: 249-253.
8. Edwin JE, Edwin JS. Color Atlas of Medicinal Plants. New Delhi: CBS Publishers and Distributors, 2006:156-157.
9. Frank SD, Amelio SR. A Phyto Cosmetic Desk Reference: Botanicals. London: CRC Press, 1999: 138.
10. Kulkarni PH, Ansari Shahida. The Ayurvedic Plants Indian Medical Science series No. 132. New Delhi: Sri Satguru Publications, Indological and Oriental Publications A division of Book centre, 2004: 191.
11. Cooke T. The Flora of Presidency of Bombay. Vol. II. Calcutta: Botanical Survey of India, 1967:176.
12. Nadkarni AK. K. M. Nadkarni's Indian Materia Medica. Vol. I. Bombay: Popular Prakashan Pvt Ltd, 1976: 701.
13. Davis B. Climbers and Wall Shrubs. Viking: Contains information on 2,000 species and cultivars, giving details of cultivation requirements, 1990:18.
14. Bhattacharjee SK. A preliminary study on the effect of nitrogen, phosphorus and potash fertilizers on *Jasminum grandiflorum* L. Pafai J

- 1983; 5(1): 21-27.
15. Bhattacharjee SK. Effect of different levels of pruning on *Jasminum grandiflorum* L. flower yield. Indian Perfumer 1985; 29(3-4): 181-185.
 16. Bhattacharjee SK, Balakrishna M. Standardization of propagation of jasmine from stem cutting, I. Effect of various growth substances on the regeneration of adventitious roots and the survival of rooted cuttings. Indian Perfumer 1983; 27(1): 32-35.
 17. Bhupal Rao JVR, Divakar NG. Variation for flower yield and essential oil content in *Jasminum grandiflorum* Linn, Proc VIIIth Int Congr of Essent Oil, Cannes-Grasse, France, 12-17 th Oct, Parfum Cosmet Aromes 1980:87.
 18. Mitra R. Bibliography on Pharmacognosy of Medicinal Plants. Lucknow: NBRI, 1985:303-304.
 19. Rashid MA, Ahmed F. Pharmacognostical studies on the leaves of *Jasminum grandiflorum* Linn. Hamdard Medicus 1995; 38(1): 55-62.
 20. Panda H. Herbs cultivation and Medicinal uses. New Delhi: National Institute of Industrial Research, 2000: 324.
 21. Khare CP. Encyclopedia of India Medicinal Plants, Rational Western Therapy, Ayurvedic and other traditional Usage. New York: Springer-Verlag Berlin Heidelberg, 2007: 267-68.
 22. Nadkarni AK. Indian Plants and Drugs with their Medical properties and uses. New Delhi: Asiatic Publishing house, 2001: 205-06.
 23. http://www.indian-herbs-exporters.com/_jasminum_grandiflorum.html.
 24. Rastogi RP, Mehrotra BN. Compendium of Indian Medicinal Plants. Vol. 1. New Delhi: CSIR, 1999:229-230.
 25. Rastogi RP, Mehrotra BN. Compendium of Indian Medicinal Plants. Vol. 2. New Delhi: CSIR, 1999:396.
 26. Rastogi RP, Mehrotra BN. Compendium of Indian Medicinal Plants. Vol. 3. New Delhi: CSIR, 2001:368.
 27. Sadhu SK, Khan MS, Ohtsuki T, Ishibashi M. Secoiridoid components from *Jasminum grandiflorum*. Phytochemistry 2007; 68(13): 1718-21.

28. Brinda S, Ulla WS, George V, Pushpangadan P, Rajasekharan S, Duus JO, et al. Angiotensin Converting Enzyme (ACE) Inhibitors from *Jasminum azoricum* and *Jasminum grandiflorum*. *Planta Med* 1998; 64: 246-250.
29. Divakar NG, Subramanian V, Sugumaran M, Vaidyanathan CS. Indole oxygenase from the leaves of *Jasminum grandiflorum*. *Plant Science Letters* 1979; (15): 177.
30. Zhao GQ, Yin ZF, Dong JX. A new secoiridoid from the flowers of *Jasminum grandiflorum*. *Yao Xue Xue Bao* 2008; 43(3): 513-517.
31. Zhao GQ, Dong JX. Triterpenoid saponin from flower bud of *Jasminum officinale* var. *grandiflorum*. *Zhongguo Zhong Yao Za Zhi* 2008; 33(1): 38-42.
32. Zhao GQ, Xia JJ, Dong JX. Glycoside from the flowers of *Jasminum officinale* var. *grandiflorum*. *Yao Xue Xue Bao* 2007; 42(10): 1066-1069.
33. Lis-Balchin, Hart SM, Lo BWH. Jasmine absolute (*Jasminum grandiflorum* L.) and its mode of action on guinea-pig ileum *in vitro*. *Phytotherapy Research* 2002; 16(5): 437-439.
34. Fulzele SV, Sattkrwar PM, Joshi SB, Dorle AK. Studies on anti-inflammatory Activity of a poly herbal formulation – Jatyadi ghrita. *Indian Drugs* 2002; 39(1): 42-44.
35. Priya Joy, Patric RD. Anti-bacterial activity studies of *Jasminum grandiflorum* and *Jasminum sambac*. *Ethnobotanical Leaflets* 2008; (12): 481-483.
36. Umamaheswari M, Ashokkumar K, Rathidevi R, Sivashanmugam AT, Subbhardadevi V, Ravi TK. Anti ulcer and *in vitro* antioxidant activities of *Jasminum grandiflorum* L. *J Ethnopharmacol* 2007; 110(3): 464-70.
37. Shanmugam M, Kuppusamy P, Krishnamurthy V, Dhanarasu S, Kaliyaperumal K. Protective effect of *Jasminum grandiflorum* Linn. on DMBA-induced chromosomal aberrations in bone marrow of wistar rats. *International Journal of Pharmacology* 2006;4(2): 406-410.
38. Kolanjiappan K, Manoharan S. Chemopreventive efficacy and anti-lipid peroxidative potential of *Jasminum grandiflorum* Linn. on 7,12-dimethylbenz (a) anthracene-

- induced rat mammary carcinogenesis. *Fundamental & Clinical Pharmacology* 2005-Dec; 19 (6): 687-93.
39. Joshi SG. *Oleaceae: Medicinal Plants*. New Delhi: Oxford & IBH Publishing Co. Pvt. Ltd, 2000: 298–300.
40. Nayak BS, Krishna M. Influence of ethanolic extract of *Jasminum grandiflorum* Linn. flower on wound healing activity in rats. *Indian J Physiol Pharmacol* 2007; 51 (2): 189–194.
41. Kumar GS, Khanam S. Anti-acne activity of Natural products. *Indian J Nat Prod* 2004; 30(4): 7-9.
42. Sharma KP, Kushwah HK, Sharma SS. Significance of Jatyadi oil in treatment of Vicharchika. *Sachitra Ayurved* 1993; 46(5): 339-347.