




Contents lists available at [ScienceDirect](http://www.sciencedirect.com)

Asian Pacific Journal of Tropical Biomedicine

journal homepage: www.elsevier.com/locate/apjtb



Original article <https://doi.org/10.1016/j.apjtb.2017.10.003>

Molecular study of astrovirus, adenovirus and norovirus in community acquired diarrhea in children: One Egyptian center study



Maysaa El Sayed Zaki*, Nermen Abo El Kheir

Clinical Pathology Department, Mansoura Faculty of Medicine, Mansoura University, Egypt

ARTICLE INFO

Article history:

Received 24 Aug 2017

Received in revised form 7 Sep 2017

Accepted 30 Sep 2017

Available online 9 Oct 2017

Keywords:

Diarrhea

Astrovirus

Adenovirus

Norovirus

Multiplex-RT PCR

ABSTRACT

Objective: To determine the prevalence of astrovirus, norovirus, adenovirus in children below five years old with diarrhea by multiplex reverse transcriptase polymerase chain reaction (RT-PCR) along with rotavirus antigen detection by enzyme linked immunosorbant assay.

Methods: The study was conducted on children below five years old complaining of acute diarrhea. The study included stool examination by molecular method for detection of norovirus, adenovirus and astrovirus by multiplex RT-PCR. Rotavirus antigen was detected in the stool by enzyme linked immunosorbant assay.

Results: The study included 100 children below 5 years old with acute diarrhea. Multiplex RT-PCR was positive in 34% of the children. The most frequently detected virus was rotavirus (44%), followed by norovirus (30%), adenovirus (20%) and astrovirus (14%). The clinical symptoms were more significantly associated with viral diarrhea such as fever ($P=0.03$), bloody diarrhea ($P=0.025$), vomiting ($P=0.0001$) and watery diarrheas ($P=0.05$). The frequency of diarrhea with viral pathogen was significantly presented in winter season (39.7%). There were significant frequencies of norovirus and adenovirus in age ranging 1–2 years old ($P=0.04$, $P=0.01$ respectively).

Conclusions: The present study spotlights on the prevalence of viral pathogens as an important etiology in diarrhea in children below five years old. Astrovirus, norovirus and adenovirus are common along with rotavirus in this group of patients. Multiplex PCR leads to improve the laboratory diagnosis of these viruses along with antigen detection method. Further longitudinal studies are required to evaluate the epidemiological data associated with these viruses and for proper management of such drastic infection.

1. Introduction

Diarrhea is a common childhood disease worldwide especially under five years old. The morbidity and mortality risks increase in developing countries in Africa and Southeast Asian countries according to the WHO report [1]. The pathogens distributions among diarrheal diseases are not well known in many countries, such as in Egypt. The knowledge of the causative pathogen is crucial for empirical treatment and preventive measures [2].

There are various pathogens that are well known to be the etiology of diarrhea including bacterial, protozoa and viral pathogens. Viruses are recognized as a cause of diarrheal diseases in children below five years old. Among enteric viruses, four categories of viruses are associated with such condition: group A rotavirus (family Reoviridae), norovirus (family Caliciviridae), adenovirus 40/41 (subgenus F), and astrovirus [3–5].

Rotavirus is a major viral cause of diarrhea in Egypt and it was studied in previous studies [6–8]. Norovirus was also reported to be common pathogen among diarrheal diseases in children especially in an outbreak; however, its role in community acquired sporadic diarrheal diseases remains to be elucidated [7,8]. Norovirus a member of family Caliciviridae, is a single stranded RNA virus. The virus is transmitted by feco-oral route by low doses leading to vomiting and diarrhea especially in winter season [9].

*Corresponding author: Maysaa El Sayed Zaki, Clinical Pathology Department, Mansoura Faculty of Medicine, Mansoura University, Egypt.

E-mail: may_s65@hotmail.com (M. El Sayed Zaki).

Peer review under responsibility of Hainan Medical University. The journal implements double-blind peer review practiced by specially invited international editorial board members.

Human adenoviruses, another viral pathogen associated with diarrheal diseases, is a double strand virus belongs to the family Adenoviridae (HAdV). There are 52 serotypes classified into six species from A to G, among which species F serotypes 40 and 41 are associated with diarrheal diseases in children [10]. Human astroviruses, a single-stranded RNA virus belonging to the Astroviridae family are a leading cause of diarrhea in children [11].

The limitation of the studies concerning viral etiology of the diarrheal diseases arises from the limitation of the methodology that can be applied in routine microbiological laboratory to define the viruses associated with such infection. Viral isolation by tissue culture is the sensitive and specific method for laboratory diagnosis [12]; however, it needs specific laboratory manipulation and it is time consuming, especially in diagnosis of community acquired infections. Also, electron microscope is specific but lacks sensitivity with detection limit of 106 viral particles/gram of stool. Antigen detection methods of viral pathogen in stool are rapid, and sensitive but it is not available for all viral pathogens. Molecular detection methods have gained a special appeal in last few years for detection of various pathogens with specific and rapid results [13–17]. The choice of molecular technique as a laboratory method for diagnosis of viruses associated with diarrheal diseases appears to be an accessible method especially if multiplex method is applied for multiple viral detection in the same sample.

Therefore the aim of the present study was to determine the prevalence of astrovirus, norovirus, adenovirus in children below five years old with diarrhea by multiplex reverse transcriptase polymerase chain reaction along with rotavirus antigen detection by enzyme linked immunosorbant assay (ELISA).

2. Materials and methods

2.1. Samples collection

The study is a cross sectional observational study that was carried at Mansoura University Children Hospital from January 2015 to January 2017. The included patients were children with acute diarrhea complaint at or below five years old recruited from out patients clinics. The study included 100 children below 5 years old with acute diarrhea. The mean age \pm SD was (3.4 \pm 1.2) years old with male predominance gender (56%). Diarrhea was defined as presence of frequent loose stool for 3 times or more during 24 h. Children older than 5 years and those without diarrhea were excluded from the study. The study was approved by Mansoura Faculty of Medicine Ethical Committee.

The study was performed according to the declaration of Helsinki and written approval was obtained from the parent of each child.

Each child was subjected to full medical history recording including age, presence of vomiting, fever and abdominal pain. Stool sample was obtained from in a clean container and transported to the laboratory.

Stool sample in the laboratory was subjected to macroscopic examination including the consistency, presence of mucus and blood. Then, each stool was divided into two aliquots: one for rotavirus antigen detection by ELISA (ELISA-Ridascreen, Germany) and the second for nucleic acid extraction for multiplex PCR detection of astrovirus, norovirus and adenovirus. The aliquots were preserved at -20°C till time of analysis.

2.2. Rotavirus antigen detection by ELISA (Ridascreen)

The test uses monoclonal antibodies for detection of rotavirus sixth viral gene protein products in sandwich type ELISA. The method was used as described by the manufacturer.

2.3. Viral DNA and RNA extraction

For each stool sample two extraction methods were used: one for viral RNA extraction for astrovirus and norovirus by QIAamp viral RNA kit and the second for viral DNA extraction for adenovirus by QIAamp DNA stool kit (Qiagen, Germany) according to the manufacturer instructions. The extracted nucleic acid was stored at -20°C till time of amplification.

2.4. Multiplex reverse transcriptase polymerase chain reaction

Both RT step and PCR were performed in the same tube using 5 μL of extracted sample (3 μL of extracted RNA and 2 μL of the extracted DNA). The protocol of the amplification was according to Rohayem *et al.* (2004) [18]. The primers sequences and the amplicon sizes of the used primers for norovirus, astrovirus and adenovirus were summarized in Table 1.

2.5. Statistical analysis

Data was analyzed by SPSS24 (SPSS, Inc., Chicago, IL). The qualitative data were expressed as percentage and *Chi*-square was used to compare between groups. The test was significant if $P < 0.05$.

Table 1

Viruses genes primers and base pair size amplicon of products.

Virus	Primers sequences	Size
Norovirus	<i>Calman-29</i> F:5' TATGGTGATGATGAAATAGTGTC-3	
Norovirus I	<i>Calman-32</i> R:5' ATTCGGGCAGAAGATTG-3	490 bp
Norovirus II	<i>Calman-1</i> F:5' GCACACTGTGTTACACTCC-3	822 bp
	<i>Calman-2</i> R:5' ACATTGGCTCTTGTCCTGG-3	
Astrovirus	<i>Mon340</i> F:5' CGTCATTGTTTGTGTCATACT-3	347 bp
	<i>Mon348</i> R:5' ACATGTGCTGCTGTTACTATG-3	
Adenovirus	<i>Adhex-1</i> F:5' GCCACCGATACGTACTTCAGCCTG-3	261 bp
A-G	<i>Adhex-2</i> R:5' GGCAGTGCCGGAGTAGGGTTAA-3	

Bp stands for base pair.

3. Results

Viral pathogens were detected in 63% of the studied children. The main complaints were fever (46%), vomiting (43%), abdominal pain (36%), watery diarrhea (30%) and bloody diarrhea (29%). The cases were more frequently presented to out patients clinics during winter season (37%). The presence of the studied viruses was detected in 63 patients (63%).

The molecular methods and antigen detection for rotavirus by ELISA revealed the presence of combined viral pathogens aiming 31.7% of the studied children, data not shown.

The most frequently detected virus was rotavirus (44%), followed by norovirus (30%), adenovirus (20%) and astrovirus (14%).

The clinical symptoms were more significantly associated with viral diarrhea such as fever ($P=0.03$), bloody diarrhea ($P=0.025$), vomiting ($P=0.0001$) and watery diarrheas ($P=0.05$). The frequency of diarrhea with viral pathogen was significantly presented in winter season (39.7%).

There was significant frequencies of norovirus and adenovirus in age ranging 1–2 years ($P=0.04$, $P=0.01$ respectively).

4. Discussion

Infectious diarrhea in children is a serious problem in developing countries with high rates of morbidity and mortality. The efforts of improving water sanitary and environmental quality conditions lead to slight improvement; however, it is still a problem [19].

Viral pathogens were detected in 63% of the studied children. The most frequently detected virus was rotavirus (44%). These results are similar to a previous study from Egypt, demonstrating that viral etiology of acute diarrhea in children affected 62% of the patients and rotavirus was the most common viral pathogen (38%) [6]. Various studies reported that rotavirus was the most frequent cause of gastroenteritis in children below 5 years old [20–22].

The adenovirus was detected in 20% of the children. This prevalence is similar to other studies from Nigeria, which presented the presence of adenovirus ranging from 18%–20% [23,24]. However, it is less than which is reported from another study in Egypt (6.8%) [6]. The differences between rates of adenovirus detection may be attributed to the difference of the methods used for laboratory diagnosis and the difference of age group of the studied children.

In the present study, norovirus was detected in 30% of children below 5 years old with gastroenteritis. The prevalence is comparable to prevalence reported to other studies reporting that norovirus frequency was similar to rotavirus but with less severe symptoms, most probably due to increasing pathogen tolerance and limited sterilizing immunity. A transmission model showed that, in highly endemic settings, protection against severe norovirus gastroenteritis could be acquired early in life, resulting in frequent asymptomatic reinfections [2,25]. This is reported in the present study as the frequent positive cases for norovirus are during the first two years.

The most frequently detected virus was rotavirus (44%). The previous studies had shown that rotavirus was the most frequent etiology for diarrhea across all age groups in many geographical regions especially in developing countries. Though, the incidence of rotavirus infections decreased with age, it still remains constant throughout all age groups [26,27].

The distinguished finding of the present study is the prevalence of astrovirus in 14% in children below 5 years old with acute gastroenteritis. The previous study reported the presence of astrovirus in 1.1% generally in children. The result of the present study demonstrates that there may be a difference in the etiology of viral infection in diarrhea within different age groups *i.e.* in children below 5 years old. This may be attributed to more exposure to these viruses in young age with limited immunity to these pathogen [6].

Another distinguished finding in the present study is the prevalence of viral coinfection in combined viral infections were detected by laboratory methods in 31.7% [28,29].

The severity of diarrhea, presence of watery and bloody diarrhea with fever were significantly prominent with viral pathogen than cases without viral etiology. This is similar to the findings of other studies [30,31], indicating that children with severe disease excrete more virus than children with less severe disease. Moreover, viral diarrhea was significantly associated with winter season, which was reported in previous studies [32,33].

The molecular method used in the present study has many advantages as it is rapid and improves the capacity to detect the viruses in accessible samples from out patients clinics. It is easier than traditional tissue culture methods [34,35]. The application of molecular method with antigen detection by ELISA detected viral pathogens in 63% of the studied children. The treatment will be symptomatic with fluid replacement therapy. Moreover, vaccination of children below 5 years old for rotavirus can assist in lowering the infectious diarrhea in those patients [36].

Further studies are needed to establish diagnostic approach for diarrhea in children with community acquired infections in low and middle income countries.

The present study spotlight on the prevalence of viral pathogens as an important etiology in diarrhea in children below 5 years old. Astrovirus, norovirus and adenovirus are common along with rotavirus in this group of patients. Multiplex PCR leads to improve the laboratory diagnosis of these viruses along with antigen detection method. Further longitudinal studies are required to evaluate the epidemiological data associated with these viruses and for proper management of such drastic infection.

Conflict of interest statement

The authors declare that they have no conflict of interest.

References

- [1] Babaniyi OA. Oral rehydration of children with diarrhea in Nigeria, a 12 year renew of impact of morbidity and mortality from diarrhea disease and diarrhea treatment practices. *J Trop Pediatr* 1991; **37**(2): 16–66.
- [2] Krumkamp R, Sarpong N, Schwarz NG, Adlkofer J, Loag W, Eibach D, et al. Correction: gastrointestinal infections and diarrheal disease in Ghanaian infants and children: an outpatient case-control study. *PLoS Negl Trop Dis* 2015; **9**(4). e0003728.
- [3] de Wit MAS, Koopmans MP, Kortbeek LM, Wannet WJB, Vinjé J, van-Leusden F, et al. Sensor, a population-based cohort study on gastroenteritis in the Netherlands: incidence and etiology. *Am J Epidemiol* 2001; **154**(7): 666–74.
- [4] Levidiotou S, Gartzonika C, Papaventsis D, Christaki C, Priavali E, Zotos N, et al. Viral agents of acute gastroenteritis in hospitalized children in Greece. *Clin Microbiol Infect* 2009; **15**(6): 596–8.

- [5] Oh DY, Gaedicke G, Schreier E. Viral agents of acute gastroenteritis in German children: prevalence and molecular diversity. *J Med Virol* 2003; **71**(1): 82-93.
- [6] Zaghoul MZ, El-Sahn SF, Galal ZA. Confection of rotavirus group A, norovirus and adenovirus in Egyptian children with gastroenteritis. *Life Sci J* 2013; **10**(2): 848-52.
- [7] Iritani N, Seto Y, Kubo H, Murakami T, Haruki K, Ayata M, et al. Prevalence of Norwalk like virus infections in cases of viral gastroenteritis among children in Osaka City. *Jpn J Clin Microbiol* 2003; **41**(4): 1756-9.
- [8] Junquera CG, de Baranda CS, Mialdea OG, Serrano EB, Sánchez-Fauquier A. Prevalence and clinical characteristics of norovirus gastroenteritis among hospitalized children in Spain. *Pediatr Infect Dis J* 2009; **28**(7): 604-7.
- [9] Robilotti E, Deresinski S, Pinsky BA. Norovirus. *Clin Microbiol Rev* 2015; **28**(1): 134-64.
- [10] Ghebremedhin B. Human adenovirus: viral pathogen with increasing importance. *Eur J Microbiol Immunol* 2014; **4**(1): 26-33.
- [11] Johnson C, Hargest V, Cortez V, Meliopoulos VA, Schultz-Cherry S. Astrovirus pathogenesis. *Viruses* 2017; **9**(1): 22.
- [12] Al-Yousif Y, Anderson J, Chard-Bergstrom C, Bustamante A, Muenzenberger M, Austin K, et al. Evaluation of a latex agglutination kit (Virogen Rotatest) for detection of bovine rotavirus in fecal samples. *Clin Diagn Lab Immunol* 2001; **8**(3): 496-8.
- [13] Benfield DA, Stotz IJ, Nelson EA, Groon KS. Comparison of a commercial enzyme-linked immunosorbent assay with electron microscopy, fluorescent antibody, and virus isolation for the detection of bovine and porcine rotavirus. *Am J Vet Res* 1984; **45**(10): 1998-2002.
- [14] de Beer M, Peenze I, da Costa Mendes VM, Steele AD. Comparison of electron microscopy, enzyme-linked immunosorbent assay and latex agglutination for the detection of bovine rotavirus in faeces. *J S Afr Vet Assoc* 1997; **68**(3): 93-6.
- [15] Nussbaum DJ, Salord JR, Rimmel DD. Evaluation of quantitative latex agglutination for detection of *Cryptosporidium parvum*, *E. coli* K99, and rotavirus in calf feces. *J Vet Diagn Investig* 1999; **11**(4): 314-8.
- [16] Parwani AV, Rosen BI, Flores J, McCrae MA, Gorziglia M, Saif LJ. Detection and differentiation of bovine group rotavirus serotypes using polymerase chain reaction-generated probes to the VP7 gene. *J Vet Diagn Investig* 1992; **4**(2): 148-58.
- [17] Chinsangaram J, Akita GY, Castro AE, Osburn BI. PCR detection of group A bovine rotaviruses in feces. *J Vet Diagn Investig* 1993; **5**(4): 516-21.
- [18] Rohayem J, Berger S, Juretzek T, Herchenröder O, Mogel M, Poppe M, et al. A simple and rapid single-step multiplex RT-PCR to detect norovirus, astrovirus and adenovirus in clinical stool samples. *J Virol Methods* 2004; **118**(1): 49-59.
- [19] Sánchez-Fauquier A, Montero V, Moreno S, Solé M, Colomina J, Iturriza-Gomara M, et al. Human rotavirus G9 and G3 as major cause of diarrhea in hospitalized children, Spain. *Emerg Infect Dis* 2006; **12**(10): 1536-41.
- [20] Vainio K, Nordbo SA, Njolstad G, Storvold G, Dollner H, Midgaard C, et al. Detection and characterization of group A rotaviruses in children hospitalized with acute gastroenteritis in Norway, 2006-2008. *J Med Virol* 2009; **81**(10): 1839-44.
- [21] Ahmed SF, Mansour AM, Klena JD, Husain TS, Hassan KA, Mohamed F, et al. Rotavirus genotypes associated with acute diarrhea in Egyptian infants. *Pediatr Infect Dis J* 2014; **33**(Suppl 1): S62-8.
- [22] Ibrahim SB, El-Bialy AA, Mohammed MS, El-Sheikh AO, Elhewala A, Bahgat S. Detection of rotavirus in children with acute gastroenteritis in Zagazig University Hospitals in Egypt. *Electron Phys* 2015; **7**(5): 1227-33.
- [23] Babalola MO, Odaibo GN, Olaleye DO, Alonge AO. Enteric adenovirus and norovirus gastroenteritis among under-5 years children in Owo, Ondo State, Nigeria. *Br J Med Med Res* 2015; **9**(8): 23-9.
- [24] Aminu M, Ahmad AA, Umoh JU, de Beer MC, Esona MD, Steele AD. Adenovirus infection in children with diarrhea disease in Northwestern Nigeria. *Ann Afr Med* 2007; **6**(4): 168-73.
- [25] Lopman B, Simmons K, Gambhir M, Vinje J, Parashar U. Epidemiologic implications of asymptomatic reinfection: a mathematical modeling study of norovirus. *Am J Epidemiol* 2014; **179**(4): 507-12.
- [26] Kotloff KL, Nataro JP, Blackwelder WC, Nasrin D, Farag TH, Panchalingam S, et al. Burden and aetiology of diarrhoeal disease in infants and young children in developing countries (the Global Enteric Multicenter Study, GEMS): a prospective, case-control study. *Lancet* 2013; **382**(9888): 209-22.
- [27] Velazquez FR, Matson DO, Calva JJ, Guerrero L, Morrow AL, Carter-Campbell S, et al. Rotavirus infections in infants as protection against subsequent infections. *N Engl J Med* 1996; **335**(14): 1022-8.
- [28] Ferreira CE, Raboni SM, Pereira LA, Nogueira MB, Vidal LR, Almeida SM. Viral acute gastroenteritis: clinical and epidemiological features of co-infected patients. *Braz J Infect Dis* 2012; **16**(3): 267-72.
- [29] Raboni SM, Damasio GAC, Ferreira CE, Pereira LA, Nogueira MB, Vidal LR, et al. Acute gastroenteritis and enteric viruses in hospitalised children in southern Brazil: aetiology, seasonality and clinical outcomes. *Mem Inst Oswaldo Cruz* 2014; **109**(4): 428-35.
- [30] Kang G, Iturriza-Gomara M, Wheeler JG, Crystal P, Monica B, Ramani S, et al. Quantitation of group A rotavirus by real-time reverse-transcription-polymerase chain reaction correlation with clinical severity in children in South India. *J Med Virol* 2004; **73**(1): 118-22.
- [31] Wolffs PF, Bruggeman CA, van Well GT, van Loo IH. Replacing traditional diagnostics of fecal viral pathogens by a comprehensive panel of real-time PCRs. *J Clin Microbiol* 2011; **49**(5): 1926-1231.
- [32] Sherchand JB, Nakagomi O, Dove W, Nakagomi T, Yokoo M, Pandey BD, et al. Molecular epidemiology of rotavirus diarrhea among children aged <5 years in Nepal: predominance of emergent G12 strains during 2 years. *J Infect Dis* 2009; **200**(Suppl 1): S182-7.
- [33] Pun SB, Nakagomi T, Sherchand JB, Pandey BD, Cuevas LE, Cunliffe NA, et al. Detection of G12 human rotaviruses in Nepal. *Emerg Infect Dis* 2007; **13**(3): 482-3.
- [34] Levine MM, Robins-Browne RM. Factors that explain excretion of enteric pathogens by persons without diarrhea. *Clin Infect Dis* 2012; **55**(Suppl 4): S303-11.
- [35] Platts-Mills JA, Gratz J, Mduma E, Svensen E, Amour C, Liu J, et al. Association between stool enteropathogen quantity and disease in Tanzanian children using TaqMan array cards: a nested case-control study. *Am J Trop Med Hyg* 2014; **90**(1): 133-8.
- [36] Mohanty E, Dwibedi B, Kar SK, Acharya AS. Epidemiological features and genetic characterization of virus strains in rotavirus associated gastroenteritis in children of Odisha in Eastern India. *Infect Genet Evol* 2017; **53**: 77-84.