

doi: 10.4103/2221-1691.221129

©2018 by the Asian Pacific Journal of Tropical Biomedicine. All rights reserved.

Pharmacodynamic profiling of optimal sulbactam regimens against carbapenem-resistant *Acinetobacter baumannii* for critically ill patients

Weerayuth Saelim^{1,2}, Wichai Santimaleeworagun^{1,6}✉, Sudaluck Thunyaharn³, Dhitiwat Changpradub⁴, Piraporn Juntanawiwat⁵

¹Department of Pharmacy, Faculty of Pharmacy, Silpakorn University, Nakhon Pathom, 73000, Thailand

²The College of Pharmacotherapy of Thailand, The Pharmacy Council, Nonthaburi, 11000, Thailand

³Faculty of Medical Technology, Nakhonratchasima College, Nakhon Ratchasima, 30000, Thailand

⁴Division of Infectious Diseases, Department of Medicine, Phramongkutklo Hospital, Bangkok, 10400, Thailand

⁵Division of Microbiology, Department of Clinical Pathology, Phramongkutklo Hospital, Bangkok, 10400, Thailand

⁶Antibiotic Optimization and Patient Care Project by Pharmaceutical Initiative for Resistant Bacteria and Infectious Diseases Working Group

ARTICLE INFO

Article history:

Received 2 October 2017

Revision 2 November 2017

Accepted 5 December 2017

Available online 1 January 2018

Keywords:

Acinetobacter baumannii

Critically ill patients

Monte Carlo simulation

MIC

ABSTRACT

Objective: To study the minimum inhibitory concentration (MIC) of sulbactam against carbapenem-resistant *Acinetobacter baumannii* (CR-AB) and to determine the dosage regimens reaching target time of free drug concentration remaining above the MIC ($fT > MIC$). **Methods:** Clinical isolates of CR-AB from patients admitted to Phramongkutklo Hospital, Thailand from January 2014 to December 2015 were obtained. The MIC of sulbactam for each CR-AB isolate was determined using the agar dilution method. Each sulbactam regimen was simulated using the Monte Carlo technique to calculate the probability of target attainment (PTA) and the cumulative fraction of response (CFR) in critically ill patients. PTA was defined by how likely a specific drug dose was to reach 40% and 60% $fT > MIC$. The CFR was the probability of drug dose covering the MIC range of CR-AB. Dosing regimens reaching above 80% of PTA and CFR, were considered as the optimal dosage for documented and empirical therapy, respectively. **Results:** A total of 118 CR-AB isolates were included in the study. The percentile at the fiftieth and ninetieth MIC of sulbactam were 64 and 192 $\mu\text{g/mL}$, respectively. For a MIC of sulbactam of 4 $\mu\text{g/mL}$, all dosage regimens achieved PTA target. However, only a sulbactam dosage of 12 g intravenous daily using 2-4 h infusion or continuous infusion that covered for isolates with a sulbactam MIC of 96 $\mu\text{g/mL}$, met the PTA or CFR targets. **Conclusions:** The MIC of sulbactam against CR-AB is quite high. The sulbactam dose of 12 g/day using prolonged infusion was required to achieve the target $fT > MIC$ for CR-AB treatment.

1. Introduction

Acinetobacter baumannii (*A. baumannii*) is a Gram negative coccobacilli that can cause nosocomial infections, such as respiratory tract infection, bacteremia, urinary tract infections, post-surgical meningitis and intra-abdominal infections[1]. *A. baumannii* is an

emerging carbapenem-resistant pathogen, becoming a global threat[2]. Carbapenem-resistant *A. baumannii* (CR-AB) has several resistance mechanisms, including enzyme production, loss of

This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial-Share Alike 3.0 License, which allows others to remix, tweak and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

©2018 Asian Pacific Journal of Tropical Biomedicine Produced by Wolters Kluwer-Medknow

How to cite this article: Saelim W, Santimaleeworagun W, Thunyaharn S, Changpradub D, Juntanawiwat P. Pharmacodynamic profiling of optimal sulbactam regimens against carbapenem-resistant *Acinetobacter baumannii* for critically ill patients. Asian Pac J Trop Biomed 2018; 8(1): 14-18.

✉Corresponding author: Wichai Santimaleeworagun, Department of Pharmacy, Faculty of Pharmacy, Silpakorn University, Nakhon Pathom, 73000, Thailand.

Tel: +66 34 255800

E-mail: swichai1234@gmail.com

porins, an efflux pump and a change of penicillin binding protein[3]. In Thailand, CR-AB is the most common causative pathogen of nosocomial pneumonia in tertiary care hospitals[4]. CR-AB has been reported to be the most prevalent pathogen in intensive care units in several studies[5,6]. In addition, colistin, sulbactam, and tigecycline are only major treatment options for CR-AB infection[7].

Colistin and tigecycline have good activity against CR-AB. Two studies in Thailand found more than 90% of CR-AB isolates were susceptible to colistin and tigecycline[8,9]. However, the pharmacokinetic properties and toxicities of colistin and tigecycline have limitations. Colistin poorly penetrates some tissues/organs and is nephrotoxic[10,11]. Tigecycline has a large volume of distribution resulting in a low serum concentration[12], so caution should be used in treating *A. baumannii* bacteremia with tigecycline[13]. Since 2013, the US Food and Drug Administration has warned increased risk of death among ventilator-associated pneumonia patients with MDR-AB treated with tigecycline[14,15].

Sulbactam is a β -lactamase inhibitor with activity against CR-AB. Sulbactam is not highly protein bound and penetrates most infected organs with adequate concentrations[16]. Sulbactam can be given in doses as high as 12 g daily without adverse reactions[17]. According to the 2016 guidelines recommended by the Infection Diseases Society of America/American Thoracic Society, sulbactam remains the drug of choice to treat MDR-AB pneumonia[18].

However, sulbactam is one of the β -lactam antibiotics. β -lactam antibiotics have augmented renal clearance and a large volume of distribution may cause inadequate tissue concentration[19]. Sulbactam shows a time-dependent bactericidal action at a percentage of the exposure time. When sulbactam is active, the free drug concentration remains above the minimum inhibitory concentration ($\%fT > MIC$) in pharmacokinetic pharmacodynamic (PKPD) targets[20]. The Monte Carlo Simulation is a technique that randomly selects a pharmacokinetics parameter value from its distribution. That process is repeated many times to generate the pharmacokinetic parameter value incorporated with the structural pharmacokinetics model to predict the appropriate dosing regimen achieving the PKPD targets[21].

Thus, the aim of this study is to determine the pharmacodynamics of sulbactam by determining its MIC. It also aims to develop a potential dosage regimen to achieve PKPD targets using the probability target of attainment (PTA) and the cumulative fraction of response (CFR) for CR-AB treatment of critically ill patients.

2. Materials and methods

2.1. Bacterial isolates

The study was conducted at Phramongkutklao Hospital in Bangkok, Thailand, a 1 200-bed tertiary care center, from January 2014 to December 2015. All clinical isolates of CR-AB obtained from patients were included in the study. Each isolate was grown in tryptic

soy broth containing 20% glycerol and kept at -70°C until used.

2.2. Determination of multidrug-resistant isolates

CR-AB was identified using the disk diffusion test and defined as resistance to carbapenems [imipenem (10 μg) or meropenem (10 μg)] [22]; the other antibiotics used during this test were: aminoglycosides [gentamicin (30 μg) or amikacin (30 μg)], antipseudomonal penicillins [piperacillin/tazobactam (100 $\mu\text{g}/10$ μg)], cephalosporins [ceftazidime (30 μg) or cefepime (30 μg)], sulfa drugs [trimethoprim-sulfamethoxazole (1.25 $\mu\text{g}/23.75$ μg)] and fluoroquinolones [ciprofloxacin (5 μg)]. The methods used followed the Clinical and Laboratory Standards Institute, guidelines, version 2017[23]. Isolates with a clear zone ≥ 11 mm to colistin (5 μg) were interpreted as susceptible.

2.3. MIC determination of sulbactam

The MIC of sulbactam was determined using the agar dilution method with Müller-Hinton agar (Oxoid) plates. The serial sulbactam (Wago, Japan) concentrations were freshly prepared between 1 and 1 024 $\mu\text{g}/\text{mL}$. A quality control strain, *Escherichia coli* ATCC 25922 (Department of Medical Sciences Culture Collection, Bangkok, Thailand) was used[23]. This study investigated MIC range, MIC50, and MIC90 of sulbactam against CR-AB. MIC range was defined as a list; the MIC value was just the difference between the largest and smallest values. MIC50 and MIC90 values were defined as the lowest concentration of sulbactam at which 50% and 90% of the isolates were inhibited, respectively.

2.4. Pharmacokinetic pharmacodynamic model study

All pharmacokinetic parameters obtained from published studies of critically ill patients were collected[24,25]. The concentration versus time was studied using a two-compartment model for critically ill patients and a one-compartment model for critically ill patients who received continuous renal replacement therapy. The pharmacokinetic and pharmacodynamic properties of sulbactam were represented by the percentage of free drug time above the MIC during the interval time ($\%fT > MIC$). The PKPD goal was defined as 40% to 60% $fT > MIC$ which was the good outcome related to the efficacy[20]. Dosage simulations were conducted using various dosages per day and dosage intervals at durations of infusion.

2.5. Monte Carlo Simulation

The PKPD investigation was conducted using a 10 000-subject Monte Carlo Simulation (Oracle Crystal Ball Classroom Faculty Edition-Oracle 1-Click Crystal Ball 201, Thailand). The Monte Carlo Program used to calculate $\%fT > MIC$ for intravenous dosage regimens of sulbactam depended on the linear pharmacokinetic behavior of the agent.

The PTA was defined by how likely a specific drug dose reached a target PKPD index ($fT > MIC$) [26]. In the present study, a target PKPD index was 40% and 60% $fT > MIC$. The CFR was the probability of drug dose covering a specified bacterial population [26]. Our bacterial population was the MIC of sulbactam among CR-AB isolates obtained from patients.

CFR was calculated by the cumulative fraction of proportional bacteria of each sulbactam MIC multiplied by PTA of each sulbactam MIC. Dosing regimen that reached above 80% of PTA and CFR was considered the optimal dosage for documented therapy and empirical therapy, respectively.

This study was approved by the institutional review board of the Royal Thai Army Medical Department and Phramongkutklao Hospital, Bangkok, Thailand (approval No. Q014h/59 issued on 24 November 2016).

3. Results

3.1. Characteristics and antimicrobial susceptibilities of CR-AB

One hundred eighteen isolates of CR-AB were collected during the study period. Seventy-one percent of the isolates were from blood, 17% from the skin and soft tissue, 6% from intra-abdominal specimens, and 6% from other sources. Seventy-seven percent of the isolates were obtained from sterile sites. Using the disk diffusion method, most CR-AB isolates (90%) in our study were found to be resistant to gentamicin, amikacin, piperacillin/tazobactam, ceftazidime, cefepime, and ciprofloxacin, making them extensively drug-resistant *A. baumannii*. Of all the study isolates, 100% were susceptible to colistin and 91.7% were susceptible to tigecycline.

3.2. Minimum inhibitory concentrations of study isolates

The MIC range, MIC50, and MIC90 for sulbactam against studied isolates were 8 to >1 024 $\mu\text{g/mL}$, 64 $\mu\text{g/mL}$, and 192 $\mu\text{g/mL}$, respectively. Each MIC value of sulbactam included 8 $\mu\text{g/mL}$ (0.8%), 16 $\mu\text{g/mL}$ (5.1%), 32 $\mu\text{g/mL}$ (11.9%), 64 $\mu\text{g/mL}$ (42.4%), 80 $\mu\text{g/mL}$ (15.3%), 96 $\mu\text{g/mL}$ (5.1%), 128 $\mu\text{g/mL}$ (7.6%), 192 $\mu\text{g/mL}$ (5.1%), 256 $\mu\text{g/mL}$ (0.8%), 512 $\mu\text{g/mL}$ (3.4%), and >1 024 $\mu\text{g/mL}$ (2.5%).

3.3. PTA

The PTA for the different sulbactam regimens at specific MICs, with targets of 40% $fT > MIC$ and 60% $fT > MIC$ is shown in Figures 1A and 1B for critically ill patients. Figures 1C and 1D indicate PTA among critically ill patients with CRRT. Among critically ill patients, for pathogens with a MIC of 4 $\mu\text{g/mL}$, all dosage regimens achieved the PTA target. However, only a sulbactam dosage of 12 g intravenous daily using 2-4 h infusion or continuous infusion that

covered for isolates with a sulbactam MIC of 96 $\mu\text{g/mL}$, met the PTA at 40% and 60% $fT > MIC$. None of all sulbactam dosage regimens reached the PTA target for critically ill patients with CRRT.




Figure 1. PTA for different sulbactam regimens at specific MICs, with targets of 40% $fT > MIC$ and 60% $fT > MIC$.

A: $fT > MIC$ 40 % among critically ill patients; B: $fT > MIC$ 60 % among critically ill patients; C: $fT > MIC$ 40 % among critically ill patients with CRRT; D: $fT > MIC$ 60 % among critically ill patients with CRRT.

3.4. CFR

Using a CFR >80%, only 4 drug regimens were determined to be appropriate for sulbactam: 3 g infused over 2 h given every 6 h, 3 g infused over 4 h given every 6 h, 12 g infused over 24 h given every 24 h and 4 g infused over 4 h given every 8 h (Table 1). However, none of the studied regimens gave a CFR >80% among patients with

CRRT.

Table 1

Cumulative fraction of response of sulbactam with various drug regimens (%).

D o s a g e regimen	Infusion time (h)	Critically ill		Critically ill with CRRT	
		40%	60%	40%	60%
		<i>fT</i> >MIC	<i>fT</i> >MIC	<i>fT</i> >MIC	<i>fT</i> >MIC
1 g/6 h	2	32.79	28.81	7.69	6.29
	4	32.57	30.46	7.38	6.71
2 g/6 h	2	71.62	65.77	24.12	20.12
	4	71.36	68.60	23.21	21.23
3 g/6 h	2	84.84	80.96	40.58	33.84
	4	84.80	83.06	39.64	36.06
4 g/8 h	2	84.51	79.64	40.27	31.34
	4	84.88	81.65	40.60	34.22
9 g/6 h	24	74.29	74.29	26.83	26.82
12 g/6 h	24	83.87	83.86	37.87	37.87

4. Discussion

CR-AB is the leading causative pathogen presenting the high mortality rate (73.3%) among critically ill patients[27]. Colistin is the agent most commonly used to treat MDR-AB and extensively drug-resistant *A. baumannii*[28]. In our study, all isolates were susceptible to colistin. However, colistin has nephrotoxicity and poor tissue penetration that limits its usefulness[10,11]. Sulbactam has been purported to be a good option to treat CR-AB[29].

With our study, MIC₅₀ and MIC₉₀ values of sulbactam against CR-AB were 64 µg/mL and 192 µg/mL, respectively. In Thailand, two studies performed at Siriraj Hospital[30] and at Queen Sirikit National Institute[31] showed values of MIC₅₀/MIC₉₀ at 32/32 and 16/89.6 µg/mL, respectively. However, unlike other related studies conducted in Thailand, the MIC₅₀ and MIC₉₀ in the present study presented higher than ever before. These distinguished MIC results might be explained because almost CR-AB isolates in our study comprised extensively drug-resistant *A. baumannii* and more than one half of isolates (60%) was obtained from critically ill patients at the ICU ward of a university-affiliated hospital.

Generally, the pharmacokinetics of sulbactam among critically ill patients differed from the general population in the aspects of volume of distribution (V_d). The reported V_d values in Thai healthy volunteers were 3.69 liters[32] while among critically ill patients, V_d of sulbactam were 14.56 liters[24]. The larger V_d values among critically ill patients effect lower serum sulbactam levels. The inadequate sulbactam concentration might be resolved by using a higher sulbactam dose and prolonged or continuous infusion as in our recommended dose of sulbactam at 12 g daily regimens. Our suggestion was similar to the results from reporting that 12 g of sulbactam daily could be achieved at the desired PTA[24].

Sulbactam is unavailable as a single agent in Thailand. Only sulbactam in combination with cefoperazone or ampicillin is available. A sulbactam dose of 12 g daily in a combination form with cefoperazone or ampicillin might result in adverse drug reactions. Thus, patients complying with a high dose of sulbactam should be closely monitored. However, several related studies have indicated

that sulbactam in combination with colistin, fosfomycin or imipenem could reduce the MIC of sulbactam against CR-AB[33–35]. Thus, the beneficial synergism of a sulbactam combination might be necessary toward the increasing PTA and CFR targets.

Our study has some limitations. First, the isolates of the CR-AB were from MIC distributions at a university-affiliated hospital which might be dissimilar when taken from other types of hospital. Second, our simulation used plasma pharmacokinetics and not tissue pharmacokinetics. Lastly, this study only suggested the probable dose of sulbactam to achieve the PKPD index. Further clinical studies are needed to determine the most beneficial dosage regimens.

In conclusion, the present study shows the MIC of sulbactam against CR-AB is quite high. However, sulbactam could be maximized in a dosage as high as 12 g daily with prolonged or continuous infusion, especially in treatment of critically ill patients.

Conflict of interest statement

We declare that there is no conflict of interest.

Acknowledgements

We would like to thank the Microbiology Laboratory Unit of Phramongkutklao Hospital for keeping the MRD-AB isolates. We also thank Dr. Tossawan Jitwasinkul, Faculty of Pharmacy, Silpakorn University for her supportive knowledge and experience.

References

- [1]Bergogne-Berezin E, Towner J. *Acinetobacter* spp. as nosocomial pathogens: microbiological, clinical, and epidemiological features. *Clin Microbiol Rev* 1996; **9**(2): 148-165.
- [2]Pogue JM, Mann T, Barber KE, Kaye KS. Carbapenem-resistant *Acinetobacter baumannii*: epidemiology, surveillance and management. *Expert Rev Anti Infect Ther* 2013; **11**(4): 383-393.
- [3]Potron A, Poirel L, Nordmann P. Emerging broad-spectrum resistance in *Pseudomonas aeruginosa* and *Acinetobacter baumannii*: mechanisms and epidemiology. *Int J Antimicrob Agents* 2015; **45**(6): 568-585.
- [4]Werarak P, Waiwarawut J, Tharavichitkul P, Pothirat C, Rungruanghiranya S, Geater SL, et al. *Acinetobacter baumannii* nosocomial pneumonia in tertiary care hospitals in Thailand. *J Med Assoc Thai* 2012; **95**: S23-S33.
- [5]Santimaleeworagun W, Wongpoowarak P, Chayakul P, Pattharachayakul S, Tansakul P, Garey KW. Clinical outcomes of patients infected with carbapenem-resistant *Acinetobacter baumannii* treated with single or combination antibiotic therapy. *J Med Assoc Thai* 2011; **94**(7): 863-870.
- [6]Wisplinghoff H, Bischoff T, Tallent SM, Seifert H, Wenzel RP, Edmond MB. Nosocomial bloodstream infections in US hospitals: analysis of 24 179 cases from a prospective nationwide surveillance study. *Clin Infect Dis* 2004; **39**(3): 309-317.
- [7]Viehman JA, Nguyen MH, Doi Y. Treatment options for carbapenem-

- resistant and extensively drug-resistant *Acinetobacter baumannii* infections. *Drugs* 2014; **74**(12): 1315-1333.
- [8]Tiengrim S, Tribuddharat C, Thamlikitkul V. In vitro activity of tigecycline against clinical isolates of multidrug-resistant *Acinetobacter baumannii* in Siriraj Hospital, Thailand. *J Med Assoc Thai* 2006; **89**: S102-105.
- [9]Piewngam P, Kiratisin P. Comparative assessment of antimicrobial susceptibility testing for tigecycline and colistin against *Acinetobacter baumannii* clinical isolates, including multidrug-resistant isolates. *Int J Antimicrob Agents* 2014; **44**(5): 396-401.
- [10]Imberti R, Cusato M, Villani P, Carnevale L, Iotti GA, Langer M, et al. Steady-state pharmacokinetics and BAL concentration of colistin in critically ill patients after IV colistin methanesulfonate administration. *Chest* 2010; **138**(6): 1333-1339.
- [11]Ordooei JA, Shokouhi S, Sahraei Z. A review on colistin nephrotoxicity. *Eur J Clin Pharmacol* 2015; **71**(7): 801-810.
- [12]Rodvold KA, Gotfried MH, Cwik M, Korth-Bradley JM, Dukart G, Ellis-Grosse EJ. Serum, tissue and body fluid concentrations of tigecycline after a single 100 mg dose. *J Antimicrob Chemother* 2006; **58**(6): 1221-1229.
- [13]Peleg AY, Potoski BA, Rea R, Adams J, Sethi J, Capitano B, et al. *Acinetobacter baumannii* bloodstream infection while receiving tigecycline: a cautionary report. *J Antimicrob Chemother* 2007; **59**(1): 128-131.
- [14]Freire AT, Melnyk V, Kim MJ, Datsenko O, Dzyublik O, Glumcher F, et al. Comparison of tigecycline with imipenem/cilastatin for the treatment of hospital-acquired pneumonia. *Diagn Microbiol Infect Dis* 2010; **68**(2): 140-151.
- [15]US Food and Drug administration. FDA Drug Safety Communication: FDA warns of increased risk of death with IV antibacterial Tygacil (tigecycline) and approves new Boxed Warning. USFDA, 2013. [Online] Available from: <https://www.fda.gov/Drugs/DrugSafety/ucm369580.htm>. [Accessed on: 2017 April 15].
- [16]Adnan S, Paterson DL, Lipman J, Roberts JA. Ampicillin/sulbactam: its potential use in treating infections in critically ill patients. *Int J Antimicrob Agents* 2013; **42**(5): 384-389.
- [17]Betrosian AP, Frantzeskaki F, Xanthaki A, Douzinas EE. Efficacy and safety of high-dose ampicillin/sulbactam vs. colistin as monotherapy for the treatment of multidrug resistant *Acinetobacter baumannii* ventilator-associated pneumonia. *J Infect* 2008; **56**(6): 432-436
- [18]Kalil AC, Metersky ML, Klompas M, Muscedere J, Sweeney DA, Palmer LB, et al. Management of adults with hospital-acquired and ventilator-associated pneumonia: 2016 clinical practice guidelines by the Infectious Diseases Society of America and the American Thoracic Society. *Clin Infect Dis* 2016; **63**(5): e61-e111.
- [19]Sime FB, Udy AA, Roberts JA. Augmented renal clearance in critically ill patients: etiology, definition and implications for beta-lactam dose optimization. *Curr Opin Pharmacol* 2015; **24**: 1-6.
- [20]Yokoyama Y, Matsumoto K, Ikawa K, Watanabe E, Shigemi A, Umezaki Y, et al. Pharmacokinetic/pharmacodynamic evaluation of sulbactam against *Acinetobacter baumannii* in in vitro and murine thigh and lung infection models. *Int J Antimicrob Agents* 2014; **43**(6): 547-552.
- [21]Roberts JA, Kirkpatrick CM, Lipman J. Monte Carlo simulations: maximizing antibiotic pharmacokinetic data to optimize clinical practice for critically ill patients. *J Antimicrob Chemother* 2011; **66**(2): 227-231.
- [22]Falagas ME, Karageorgopoulos DE. Pandrug resistance (PDR), extensive drug resistance (XDR), and multidrug resistance (MDR) among Gram-negative bacilli: need for international harmonization in terminology. *Clin Infect Dis* 2008; **46**(7): 1121-1122.
- [23]Clinical and Laboratory Standard Institute. *Performance standards for antimicrobial susceptibility testing; twenty-seventh informational supplement*. CLSI document M100S; Wayne: PA; 2017.
- [24]Jaruratanasirikul S, Wongpoowarak W, Wattanavijitkul T, Sukarnjanaset W, Samaeng M, Nawakitranngsan M, et al. Population pharmacokinetics and pharmacodynamics modeling to optimize dosage regimens of sulbactam in critically ill patients with severe sepsis caused by *Acinetobacter baumannii*. *Antimicrob Agents Chemother* 2016; **60**(12): 7236-7244.
- [25]Gao C, Tong J, Yu K, Sun Z, An R, Du Z. Pharmacokinetics of cefoperazone/sulbactam in critically ill patients receiving continuous venovenous hemofiltration. *Eur J Clin Pharmacol* 2016; **72**(7): 823-830.
- [26]Asín-Prieto E, Rodríguez-Gascón A, Isla A. Applications of the pharmacokinetic/pharmacodynamic (PK/PD) analysis of antimicrobial agents. *J Infect Chemother* 2015; **21**(5): 319-329.
- [27]Nazer LH, Kharabsheh A, Rimawi D, Mubarak S, Hawari F. Characteristics and outcomes of *Acinetobacter baumannii* infections in critically ill patients with cancer: a matched case-control study. *Microb Drug Resist* 2015; **21**(5): 556-561.
- [28]Khawcharoenporn T, Pruetpongpun N, Tiamsak P, Rutchana-wech S, Mundy LM, Apisarnthanarak A. Colistin-based treatment for extensively drug-resistant *Acinetobacter baumannii* pneumonia. *Int J Antimicrob Agents* 2014; **43**(4): 378-382.
- [29]Chen H, Liu Q, Chen Z, Li C. Efficacy of sulbactam for the treatment of *Acinetobacter baumannii* complex infection: A systematic review and meta-analysis. *J Infect Chemother* 2017; **23**(5): 278-285.
- [30]Pongpech P, Amornopparattanakul S, Panapakdee S, Fungwithaya S, Nannha P, Dhiraputra C, et al. Antibacterial activity of carbapenem-based combinations against multidrug-resistant *Acinetobacter baumannii*. *J Med Assoc Thai* 2010; **93**(2): 161-171.
- [31]Punpanich W, Munsrichoom A, Srisarang S, Treeratweeraphong V. In vitro activities of colistin and ampicillin/sulbactam against *Acinetobacter baumannii*. *J Med Assoc Thai* 2011; **94**: S95-100.
- [32]Jaruratanasirikul S, Wongpoowarak W, Aeinlang N, Jullangkoon M. Pharmacodynamics modeling to optimize dosage regimens of sulbactam. *Antimicrob Agents Chemother* 2013; **57**(7): 3441-3444.
- [33]Santimaleeworagun W, Wongpoowarak P, Chayakul P, Pattharachayakul S, Tansakul P, Garey KW. In vitro activity of colistin or sulbactam in combination with fosfomycin or imipenem against clinical isolates of carbapenem-resistant *Acinetobacter baumannii* producing OXA-23 carbapenemases. *Southeast Asian J Trop Med Public Health* 2011; **42**(4): 890-900.
- [34]Thamlikitkul V, Tiengrim S. In vitro activity of colistin plus sulbactam against extensive-drug-resistant *Acinetobacter baumannii* by checkerboard method. *J Med Assoc Thai* 2014; **97**: S1-6.
- [35]Laishram S, Anandan S, Devi BY, Elakkiya M, Priyanka B, Bhuvaneshwari T, et al. Determination of synergy between sulbactam, meropenem and colistin in carbapenem-resistant *Klebsiella pneumoniae* and *Acinetobacter baumannii* isolates and correlation with the molecular mechanism of resistance. *J Chemother* 2016; **28**(4): 297-303.