



Effective Biosurfactants Production by *Pseudomonas aeruginosa* and its Efficacy on Different Oils

K.V. Sekar¹, Sarita Kumari^{2*}, A. Nagasathya³, S. Palanivel⁴ and Subramanyam Nambaru⁵

^{1,2}Department of Microbiology, The Oxford College of Science, Bangalore-102, India.

^{3,4}P.G. and Research Dept. of Microbiology, J.J. College of Arts and Science, Pudukkottai- 622404, India.

⁵Research Scholar, CAS in Marine Biology, Annamalai University, Parangipettai, Tamil Nadu, India.

Abstract: A rhamnolipid producing bacterium, *Pseudomonas aeruginosa* was isolated from contaminated soil with oily wastes. The *Pseudomonas aeruginosa* grown with glucose and corn oil as a carbon source produced bio-surfactant. This bio-surfactant was purified by procedures that included chloroform-ethanol extraction and 0.05M bicarbonate treatments. The active compound was identified as rhamnolipid by using thin layer chromatography. The emulsification activity of bio-surfactant, the coconut oil responded better than the olive oil, groundnut oil and sunflower oil and gave a maximum level of 1 cm.

Keywords: Biosurfactant production, *Pseudomonas aeruginosa*, rhamnolipid, test efficacy.

1. Introduction

Biosurfactants are amphiphilic compounds produced on living surfaces, mostly microbial cell surfaces or excreted extra cellularly. It contains hydrophobic and hydrophilic moieties that reduce surface tension and interfacial tensions between individual molecules at the surface and interface respectively. Well known synthetic surfactants are used for a wide variety of purposes, such as emulsification, foaming, detergency, solubilization, wetting and spreading. Almost all surfactants currently in use are chemically derived from petroleum. However, the interest in microbial surfactants has been steadily increasing in recent years due to their diversity, environment friendly characteristics, the possibility of their production through fermentation and their potential applications in such areas such as the environmental protection, surface crude oil recovery, health care and food processing industries (Zajic, 1976). Biosurfactants have the following

advantages compared with chemical surfactants; lower toxicity, higher biodegradability (Swisher, 1970), better environmental compatibility (Parkinson, 1985), higher foaming (Haferburg and Hommel, 1986), higher selectivity and specific activity at extreme temperature, pH levels and salinity. Most bio-surfactant studies have used bacteria grown on hydrocarbon substrates. Few workers have utilized carbohydrates (Suzuki *et al.*, 1974; Cooper *et al.*, 1981). *Pseudomonas aeruginosa* has been one of the most widely studied bio-surfactants. Several rhamnolipid bio-surfactants produced by *Pseudomonas aeruginosa* strains have been characterized (Edwards and Hayashi, 1965; Hirayama and Kato, 1982; Hisatsuka *et al.*, 1985). Reiling *et al.*, (1986) developed a continuous production of *Pseudomonas aeruginosa* to achieve 2.25g rhamnolipids as a L-rhamnose for scientific and industrial purposes. In this paper, we consider the production and test for the efficacy of extracting bio-surfactants on different oil samples.

*Corresponding author:
E-mail: saritamicro@gmail.com.

2. Materials and Methods

Pseudomonas aeruginosa was isolated from soil contaminated with oily wastes. The organism was maintained in nutrient agar slants in an atmosphere of 37°C. The organisms were grown in 50ml sterile nutrient broth in a 250ml flask at 37°C. Erlenmeyer flasks of 1000ml capacity containing 250ml of the mineral salt medium with glucose and corn oil as the carbon source were individually inoculated with 5ml of particular inoculums. The mineral salt medium contained (g l^{-1}) $(\text{NH}_4)_2 \text{SO}_4$, 0.5945; $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 0.0001; $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$, 0.002; $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.061; NaCl , 0.124; Glucose, 0.112; Corn oil, 1.25ml and phosphate buffer, 250ml. The phosphate buffer contained NaH_2PO_4 , 0.39; Na_2HPO_4 , 0.35. The flasks were incubated in a rotatory shaker incubator at 30°C for 72 hours. The culture so obtained was used for the extraction of bio-surfactants.

Rhamnolipid was recovered from the culture supernatant after the removal of cells by centrifugation at 6,800rpm for 20 minutes. The precipitate was dissolved in 25ml of 0.05M bicarbonate (pH 8.6) re-acidified and re-centrifuged at 12,000rpm for 20 minutes. The precipitate was then extracted with chloroform-ethanol (2:1) three times and it is then concentrated by evaporation at room temperature to obtain a bio-surfactant concentrate. Following the evaporation, the bio-surfactant concentrate is dissolved in 25ml of 0.05M bicarbonate (pH 8.6).

The homogeneity was checked and the identification was made by thin layer chromatography (TLC). Precoated TLC plates spotted with samples were developed with following solvent systems chloroform: methanol: acetic acid (65:15:2 by volume). The compounds on the plates were visualized by heating after spraying with 2% anthrone solution in 0.2% H_2SO_4 . Identification was performed by running rhamnolipid standards simultaneously.

Equal volumes of extracted bio-surfactant in 0.05M *Pseudomonas aeruginosa* bicarbonate and the

oil samples i.e., olive oil, coconut oil, groundnut oil and sunflower oil being tested (5ml+5ml) were taken in screw cap tubes. These tubes were then vortexed well and incubated at room temperature. The tubes were examined periodically for a decrease in the thickness of the oil layers and hence an increase in the bio-surfactant layer. Observations were made every 1 hour to 8 hours.

3. Results

Using an enrichment culture technique a bio-surfactant producing soil bacteria was isolated and identified as *Pseudomonas aeruginosa* on the basis of various physiological and biochemical tests as described in Bergey's manual of determinative bacteriology (1974). The bacterium is gram negative, motile, rod-like and grows aerobically. The enrichment of organism in the mineral salt medium incorporated with trace amounts of glucose and 0.05% corn oil act as a carbon source as well as to induce the bio-surfactant production. Rhamnolipid was recovered from culture supernatant by centrifugation, precipitation with hydrochloric acid, purified by re-centrifugation and bicarbonate treatment. The purified samples were detected as one spot of the thin layer chromatography with above mentioned solvent systems (for detailed see the materials and methods section). The concentrate of bio-surfactant obtained following extraction from the production medium was tested with 4 different samples; coconut oil, olive oil, groundnut oil and sunflower oil. When tested with coconut oil, the level of bio-surfactant was found to increase by 1cm in 8 hours which was found to be greatest among four samples tested. Following incubation with olive oil, groundnut oil and sunflower oil, the thickness of the bio-surfactant layer were found to rise by 0.5cm, 0.3cm and 0.2cm in 8 hrs respectively (Fig. 1; Table 1).




Fig. 1. Test efficacy of four different oils.

Table 1. Test efficacy of four different oils

S. No.	Name of sample	Level of aqueous phase (Bio-surfactants) at									
		0 hours	1 hours	2 hours	3 hours	4 hours	5 hours	6 hours	7 hours	8 hours	
1	Coconut Oil	4	4	4.1	4.1	4.3	4.5	4.6	4.9	5.0	
2	Olive Oil	4	4	4	4.1	4.1	4.1	4.2	4.4	4.5	
3	Groundnut Oil	4	4	4	4	4.1	4.1	4.2	4.3	4.3	
4	Sunflower Oil	4	4	4	4	4.1	4.1	4.1	4.2	4.2	

4. Discussion

The result obtained from the present investigation revealed the ability of *Pseudomonas aeruginosa* in producing bio-surfactant in minimal media incorporating with glucose and corn oil. During growth on *Pseudomonas aeruginosa* undergoes two distinct types of metabolism; exponential growth linked with amino acid catabolism and stationary growth linked with glucose metabolism. This behavior is known as reverse diauxic (Hamilton and Dewes, 1960). At this transition point, bio-surfactant production was initiated. Rhamnolipids are also produced by *Pseudomonas aeruginosa* in media containing glucose or glycerol as a carbon source, particularly when the cells become limited for nitrogen (Guerra-Santos et al., 1986; Mulligan and Gibbs, 1989; Wagner et al., 1983), indicating that they may serve other roles besides being involved in solubilizing hydrophobic substrates. Like that of our study, Primary isolation was made in Nutrient broth followed by enrichment in minimal medium containing traces of glucose and 0.05% corn oil. Corn oil serving as the sole carbon source produces significantly increased levels of both cellular mass and rhamnolipids.

The increase in rhamnose cannot be attributed solely to the presence of more cells since cell mass was 2-3 times greater when corn oil was used. In addition to corn oil, other vegetable oils including soya bean oil, coconut oil and cottonseed oil result in similar cell growth and rhamnolipid production (Linhardt et al., 1989). Several types of organisms are involved in the production of bio-surfactant, but *Pseudomonas aeruginosa* have been reported by other workers as excellent bio-surfactant producer and have been widely studied. Corn oil is readily available at low cost and its use as a carbon source in the absence of glucose avoids potential sugar contamination of rhamnose product and produces significantly increased levels of both cellular mass and rhamnolipid. This enrichment induced the production of biosurfactants. For the detection of rhamnolipids, four independent tests were carried out. These included the lowering of the interfacial tension (IFT) by rhamnolipids (Guerra-Santos et al., 1984), the detection of thin layer chromatography (Koch et al., 1988), the haemolysis of erythrocytes by rhamnolipids (Johnson and Marrazzo et al., 1980) and the growth inhibition of *Bacillus subtilis* exerted by rhamnolipids (Itoh et al., 1971).

In our present investigation, the complete separation of rhamnolipids was performed by TLC.

The efficacy of bio-surfactant concentrates extracted was tested with four different oil samples viz; coconut oil, groundnut oil, sunflower oil and olive oil. The greatest activity was found in coconut oil followed by olive oil, groundnut oil and sunflower oil in the above sequence. In coconut oil, the production got increased 1cm, emulsify. The thickness also increased in olive oil, this may be due to saturated fatty acids in olive oil and is hence relatively easy for the organism to act upon. The complete emulsification of sunflower oil would hence take a long time compared to the above mentioned saturated oils due to the longer time is taken for the organism to break down the unsaturated fatty acids. The addition of hydrocarbon forced the existing cells to produce high concentrations of bio-surfactants.

5. Conclusion

The present study demonstrated the bio-surfactant producing potential of *Pseudomonas aeruginosa* isolated from oil contaminated soil using corn oil as sole carbon source. Extracted bio-surfactant showed emulsifying property. The ability of the produced bio-surfactant to emulsify different oils is an important feature which can be used for bioremediation of oil spills. Due to efficient emulsification properties, it can be used in industrial and environmental applications.

Acknowledgment

We thank Head Department of Microbiology, The Oxford College of Science, Bangalore.

References

- [1]. Cooper, D.G. and Zajic, J.E. (1980). Surface active compounds from microorganisms. *Adv. Appl. Microbiol.* 26: 229-253.
- [2]. Cooper, D.G., MacDonald, C.R., Duff, S.J.B. and Kosaric, N. (1981). Enhanced production of surfactin from *Bacillus subtilis* by continuous product removal and metal cation additions. *Appl. Environ. Microbiol.* 42: 408-412.
- [3]. Edwards, J. and Hayashi, J. (1965). Structure of a rhamnolipid from *Pseudomonas aeruginosa*. *Arch. Biochem. Biophys.* 111: 415-421.
- [4]. Guerra-Santos, L., Kappeli, O. and Fiechter, A. (1984). *P. aeruginosa* biosurfactant production

- in continuous culture with glucose as carbon source. *Appl. Environ. Microbiol.* 48: 301-305.
- [5]. Guerra-Santos, L., Kappeli, O. and Fiechter, A. 1986. Dependence of *Pseudomonas aeruginosa* continuous culture biosurfactant production on nutritional and environmental factors. *Appl. Environ. Microbiol.* 24: 443-448.
- [6]. Haferburg, D. and Hommel, R. (1986). *Adv. Biochem. Eng./ Biotechnol.* 33: 53-93.
- [7]. Hamilton, W.A. and Dawes, E.A. (1960). The nature of the diauxic effect with glucose and organic acids in *Pseudomonas aeruginosa*. *Proc. Biochem. Soc. Biochem. J.* 76: 70.
- [8]. Harvey, S., Elashvili, I., Valdes, J.J., Kamely, D. and Chakrabarty, A.M. (1990). Enhanced removal of the Exxon Valdez spilled oil from the Alaskan gravel by a microbial surfactant. *Bio/Technology.* 8: 228-230.
- [9]. Hirayama, T. and Kato, I. (1982). Novel rhamnolipids from *P. aeruginosa*. *FEBS Lett.* 139: 81-85.
- [10]. Hisatsuka, K., Nakahara, T., Sano, N. and Yamada, K. (1971). Formation of rhamnolipid by *Pseudomonas aeruginosa*: its function in hydrocarbon fermentations. *Agric. Biol. Chem.* 35: 686-692.
- [11]. Itoh, S., Honda, H., Tomita, F. and Suzuki, T. (1971). Rhamnolipid produced by *Pseudomonas aeruginosa* grown on n-paraffin. *J. Antibiot.* 24: 855-859.
- [12]. Johnson, M.K. and Bose-Marrazzo, D. (1980). Production and properties of heat-stable extracellular haemolysin from *Pseudomonas aeruginosa*. *Infect. Immun.* 29: 1028-1033.
- Linhardt, R.J., Bakhit, R. and Daniels, L. 1989. Microbially produced rhamnolipid as a source of rhamnose. *Biotechnol. Bioeng.* 33: 365-368.
- [13]. Kappeli, O. and Finnerty, W.R. (1980). Characteristics of hexadecane partition by the growth medium of *Acinetobacter sp.* *Biotechnol. Bioeng.* 22: 495-503.
- [14]. Koch, A.K., Reiser, J., Kappeli, O. and Fiechter, A. (1988). Genetic construction of lactose-utilizing strains of *Pseudomonas aeruginosa* and their application in biosurfactant production. *Bio/Technology.* 6: 1335-1339.
- [15]. Mulligan, C.N. and Gibbs, B.F. (1989). Correlation of nitrogen metabolism with biosurfactant production by *Pseudomonas aeruginosa*. *Appl. Environ. Microbiol.* 55: 3016-3019.
- [16]. Parkinson, M. 1985. Biosurfactants. *Biotech. Adv.* 3: 65-83.
- [17]. Rapp, P., Bock, H., Wray, V. and Wagner, F. (1979). Formation, isolation and characterization of trehalose dimycolates from *Rhodococcus erythropolis* grown on n-alkanes. *J. Gen. Microbiol.* 115: 491-503.
- [18]. Reiling, H.E., Thanei-Wyes, U., Guerra-Santos, L., Hirt and Kappeli, O. (1985). Pilot production of rhamnolipid biosurfactant by *Pseudomonas aeruginosa*. *Appl. Environ. Microbiol.* 51: 985-989.
- [19]. Robert, M., Mercade, M.E., Bosch, M.P., Parra, J.L., Espuny, M.J., Manresa, M.A. and Guinea, J. (1989). Effect of the carbon source on biosurfactant production by *Pseudomonas aeruginosa* 44T. *Biotechnol. Lett.* 11: 871-874.
- [20]. Suzuki, T., Tanaka, H. and Itoh, H. (1974). Sucrose lipids of *Arthrobacter*, *Corneybacterium* and *Nocardia* grown on sucrose. *Agric. Biol. Chem.* 38: 557-563.
- [21]. Swisher, R.D. (1970). Surfactant Biodegradation. Marcel Dekker, New York.
- [22]. Van Dyke, Gulley, M.I.S.L., Lee, H. and Trevors, J.T. (1993). Evaluation of microbial surfactants for recovery of hydrophobic pollutants from soil. *J. Ind. Microbiol.* 11: 163-170.
- [23]. Wagner, F., Bock, H., Kretschmer, A., Lang, S. and Syldatk, C. (1983). Production and chemical characterization of surfactants from *Rhodococcus erythropolis* and *Pseudomonas sp.* MUB grown on hydrocarbons. p.55-60. In J. E. Zajiic, D.G. Cooper, T. R. Jack and N. Kosaric (ed.), Microbial enhanced oil recovery. PennWell Publishing. Tucson, Ariz.
- [24]. Zajiic, J.E. and Steffens, W. 1974. Biosurfactants. *CRC Crit. Rev. Biotechnol.* 1: 87-101.