

Record of the largest longtail tuna, *Thunnus tonggol* (Bleeker 1851) from off shore Salalah in the Sultanate of Oman

Dawood Al-Mamari¹ • Lubna Al-Kharusi¹ • Fatma Al-Kiyumi¹ • Saeed Al-Shogebai² • Ibrahim Al-Anboori¹ • Rashed Al-Seneadi¹ • Sergiy Khrorv¹

¹Marine Science and Fisheries Centre, Ministry of Agriculture and Fisheries Wealth, Muscat, Sultanate of Oman

²Fisheries Research Center, Ministry of Agriculture and Fisheries Wealth, Salalah, Sultanate of Oman

Correspondence: Dawood Al-Mamari, Marine Science and Fisheries Centre, Ministry of Agriculture and Fisheries Wealth; Email: almaamari007@hotmail.com

Received: 12 Oct 2014, Received in revised form: 23 Dec 2014, Accepted: 29 Dec 2014, Published online: 31 Dec 2014

Citation: Al-Mamari D, Al-Kharusi L, Al-Kiyumi F, Al-Shogebai S, Al-Anboori I, Al-Seneadi R and Khrorv S (2014) Record of the largest size of longtail tuna, *Thunnus tonggol* (Bleeker 1851) from off shore Salalah in the Sultanate of Oman. Journal of Fisheries 2(3): 215-216. DOI: [dx.doi.org/10.17017/jfish.v2i3.2014.57](https://doi.org/10.17017/jfish.v2i3.2014.57)

Abstract

Capture of a great Longtail tuna, *Thunnus tonggol* from off Shore of the city of Salalah, Arabian Sea, Sultanate of Oman in January 2013 is described in this paper. Its morphometric and meristic characters are discussed.

Keywords: Longtail tuna, *Thunnus tonggol*, Sultanate of Oman, Dohfar, Arabian Sea

INTRODUCTION

Longtail tuna, *Thunnus tonggol* (Bleeker, 1851) is an epipelagic migratory tuna species and cosmopolitan. It is widely distributed in tropical to temperate provinces of the Indo-Pacific between 47°N and 33°S (Froese and Pauly 2009). Its diffusion is exclusive compared with those of other species in the genus *Thunnus* that commonly variety through open oceans, in that longtail tuna almost completely inhabit neritic areas close to landmasses, and are hardly found offshore (Yesaki 1994). Longtail tuna is the second smallest of eight species of the genus *Thunnus* and grows to a maximum size of 142 cm in total length and 35.9 kg in weight (IGFA 2008). Due to their coastal distribution, longtail tuna are greatly exploited by small-scale artisanal and commercial fisheries in at least 17 countries throughout Indo-Pacific (Yesaki 1994). They are commonly targeted by purse-seine, gillnet, and trolling (Yesaki 1994). Thailand, Indonesia, Malaysia, and Iran contribute greatest to stated annual landings, which touched 248 000 ton in 2007 (FAO, 2009).

METHODOLOGY

On the 6th of January 2013 a great specimen of *Thunnus tonggol* was captured along the coast of the Arabian Sea, city of Salalah by a local fisherman using longline. Specimen was kept on ice and taken to the laboratory. Fish was weighed nearest kg, measured for fork and total length in cm, and sexed by visual examination of the gonads. Morphometrics and meristics were characters measured by following Marr and Schaefer (1949).

RESULTS

The fish was 145.2 cm in total length (TL) and 27.418 kg in body weight (Figure 1). Morphometrics and meristics were presented in Table 1.

Unusual caught of large longtail tuna in Arabian Sea, city of Salalah could indicate the suitable environment in this area. However the Arabian Sea is rich in various nutrients that make the most of the areas suitable for the abundance and diversity of fish breeding. More research is needed and important for the study of fish stocks and diversity in the Arabian Sea.



Figure 1: *Thunnus tonggol*, 145.2 cm total length, captured by long line from the coast of Salalah City in the Arabian Sea

In addition, Arabian Sea is open to Indian Ocean, where a continuous migration of fish in the region and it is expected the presence of new species of fishes has not yet been recorded.

Table 1: Morphometric measure and meristic characters of *Thunnus tonggol* along the coast of the Arabian Sea, city of Salalah, Sultanate of Oman

Morphometric characters	Length (cm)
Total Length (TL)	145.2
Fork Length (FL)	135
% in TL	93
Head Length (HL)	31
Pre orbital Length	10.5
% in HL	33.9
Postorbital Length	17.3
% in HL	55.8
Pre dorsal fin length 1	36.6
% in TL	25.2
Post dorsal fin length 1	63.1
% in TL	43.5
Pre dorsal fin length 2	66.1
% in TL	45.5
Post dorsal fin length 2	77
% in TL	53
Pre pectoral fin length	32.5
% in TL	22.4
Pectoral fin length	20.2
% in TL	13.9
Pre pelvic Length	34.4
% in TL	23.7
Pelvic Length	14
% in TL	9.6
Pre anus length	71
% in TL	48.9
Pre anal fin length	73.3
% in TL	50.5
Post anal fin length	85.1
% in TL	58.6
Maximum body depth	27.4
% in TL	18.9
Caudal peduncle length	3.6
% in TL	2.5
Meristic characters	
Number of Dorsal fin	13
Number of upper- adipose	9
Number of lower- adipose	8

ACKNOWLEDGEMENT

We would like to acknowledge the ministry of Agriculture and Fisheries Wealth, the Marine Science and Fisheries Center and the Agriculture and Fisheries Development Fund for giving us the occasion to work on the fish samples within the Tuna project in Sultanate of Oman.

REFERENCES

- Bleeker P (1851). Nieuwebijdrage tot de kennis der Percoïdei, Scleroparei, Sciaenoïdei, Maenoïdei, Chaetodontoïdei en Scomberoïdei van den Soenda-Molukschen Archipel. *Natuurkundig Tijdschrift voor Nederlandsch Indië* 2: 163-179.
- FAO (2009) FISHSTAT Plus: universal software for fishery statistical time series, version 2.3. FAO Fisheries Department, Fishery Information, Data and Statistics Unit.
- Froese R and Pauly DE (2009). FishBase. World Wide Web electronic publication. www.fishbase.org, accessed on 7 February 2009.
- IGFA (International Game Fishing Association)(2008). World Record Game Fishes. International Game Fish Association, FL, USA.
- Marr JC and Schaefer MB (1948) Definitions of body dimensions used in describing tunas-U.S. Fish. Midlife Service Fisher Bull 47: 241-244
- Murray T, Swanson P and Carey C (1984) Tuna resources and tuna research in New Zealand. Country Statement - New Zealand. South Pacific Commission, 16th Regional Technical Meeting on Fisheries. Noumea, New Caledonia 13-17 Aug. 1984.
- Yesaki M (1994). A review of the biology and fisheries for longtail tuna (*Thunnus tonggol*) in the Indo-Pacific Region. FAO Fisheries Technical Paper 336: 370-387