

Mouse models of epithelial ovarian cancer for preclinical studies

Sergey Karakashev¹, Ru-Gang Zhang^{1*}

¹ Immunology, Microenvironment & Metastasis Program, The Wistar Institute, Philadelphia, PA 19104, USA

ABSTRACT

Epithelial ovarian cancer (EOC) is the leading cause of gynecological cancer-related mortality in the developed world. EOC is a heterogeneous disease represented by several histological and molecular subtypes. Therefore, exploration of relevant preclinical animal models that consider the heterogenic nature of EOC is of great importance for the development of novel therapeutic strategies that can be translated clinically to combat this devastating disease. In this review, we discuss recent progress in the development of preclinical mouse models for EOC study as well as their advantages and limitations.

Keywords: Epithelial ovarian cancer; Patient-derived xenografts; Orthotopic mouse model; Subcutaneous mouse model; Intraperitoneal mouse model; Syngeneic mouse model; Genetic engineered mouse model

INTRODUCTION

More than 85% of ovarian cancers are of epithelial origin. Epithelial ovarian cancer (EOC) is broadly divided into two types. Type I EOC includes clear cell, endometrioid, mucinous, and low-grade serous carcinomas, while type II EOC consists primarily of high-grade serous carcinomas (Kurman & Shih *et al.*, 2011). Despite recent advances in our

understanding of EOC pathology, it remains a devastating disease with poor outcome compared to many other types of cancer (Siegel *et al.*, 2020). Indeed, EOC survival rates in patients have only slightly improved in the last several decades, and EOC remains a major cause of cancer-related deaths in women worldwide (Torre *et al.*, 2018). The high mortality rate is, in part, a result of limited therapeutic options and therefore novel therapeutic strategies are greatly needed. Despite the significant progress that has been made in recent decades in our understanding of the molecular mechanisms underlying ovarian cancer initiation and progression, novel therapies to improve outcome in EOC patients remain elusive. EOC is a genetically heterogeneous disease represented by several histological and molecular subtypes that are distinct in their response to different anticancer agents (The Cancer Genome Atlas Research Network, 2011). Given its heterogenic nature, EOC requires appropriate preclinical animal models to recapitulate the pathobiology of its various subtypes. Mouse models are crucial not only for improving our understanding of the pathobiology underlying EOC initiation and progression, but more importantly for developing clinically relevant therapeutic strategies (House *et al.*, 2014). Recent advances in preclinical EOC *in vivo* models have led to the development of several FDA-approved therapies. Preclinical EOC *in vivo* models have been widely used to demonstrate

Received: 28 December 2020; Accepted: 26 January 2021; Online: 28 January 2021

Foundation items: This work was supported by the US National Institutes of Health (R01CA160331, R01CA163377, R01CA202919, R01CA239128, P01AG031862, P50CA228991 to R.G.Z. and K99CA241395 to S.K.), US Department of Defense (OC180109 and OC190181 to R.G.Z.). The Honorable Tina Brozman Foundation for Ovarian Cancer Research (to R.G.Z.), and Ovarian Cancer Research Alliance Collaborative Research Development Grant (to R.G.Z.). Core facilities support was provided by a Cancer Centre Support Grant (CA010815) to the Wistar Institute

*Corresponding author, E-mail: rzhang@wistar.org

DOI: [10.24272/j.issn.2095-8137.2020.382](https://doi.org/10.24272/j.issn.2095-8137.2020.382)

Open Access

This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Copyright ©2021 Editorial Office of Zoological Research, Kunming Institute of Zoology, Chinese Academy of Sciences

the efficacy of FDA-approved poly(ADP-ribose) polymerase (PARP) inhibitors, such as olaparib, niraparib, and rucaparib, for EOC maintenance therapy (Alhilli et al., 2016; Kortmann et al., 2011). Bevacizumab, a vascular endothelial growth factor (VEGF) inhibitor, is another effective anticancer therapy in EOC *in vivo* models and also recently approved by the FDA (Byrne et al., 2003; Hu et al., 2005). However, despite this success, novel preclinical EOC models are required due to the heterogeneity of EOC and the development of resistance to approved anticancer therapies.

XENOGRAFT EOC MODELS

Ovarian cancer cell lines derived from ascites or primary EOCs have been extensively used to study the molecular mechanisms involved in the regulation of tumor progression and chemoresistance and for the development of novel therapeutics. Historically, ovarian cancer cell lines engrafted in immunocompromised mice were the first EOC mouse models developed. Mice subjected to ionizing radiation were the first animal hosts used for ovarian cancer xenografts. Subsequently, several immunodeficient mouse strains were developed and gained wide popularity, including nude mice with low levels of T-lymphocytes due to spontaneous deletion of *Foxn1* (Cordier & Haumont, 1980), SCID mice with severe B-cell and T-cell immunodeficiency due to mutation of *Prkdc* (Pla & Mahouy, 1991), and NOD/SCID mice with B-cell and T-cell immunodeficiency and compromised cytokine signaling (Shultz et al., 2005). These mouse strains have allowed the use of the large panel of EOC human cell lines generated in the last few decades and represent various EOC subtypes. In

general, EOC xenografts are fast growing tumors that permit *in vivo* testing of previously observed *in vitro* experiments. The low cost, high engraftment probability, and low tumor size variability mouse to mouse make these models popular in the EOC field. These xenografts are currently used to study various aspects of EOC progression such as tumor initiation, growth, invasion, metastasis, and sensitivity to anticancer agents. One of the greatest advantages of EOC mouse xenografts is the ability to use genetically modified cell lines. Xenograft EOC mouse models can be classified based on engraftment strategy or source of engrafted material, as follows:

Classification based on engraftment strategy

EOC cells can be engrafted subcutaneously, intraperitoneally, or orthotopically. Each method has its own advantages and limitations for meeting specific needs (Table 1). Some models are suitable for studies on the early stages of cancer, whereas others are useful for studies on invasion and metastasis. Subcutaneous xenograft models were the first animal models applied for the study of EOC and are still widely used today. Subcutaneous xenograft models gained popularity due to their simplicity, reliability, and low cost. Subcutaneous EOC cell injections do not require specialized expertise or equipment. Furthermore, tumors are formed within several weeks and their growth is limited to the injection site. Tumor size can be easily measured from the early stages of tumor formation, thus making subcutaneous xenograft models a good choice for drug response evaluation. However, the absence of a physiologically relevant tumor microenvironment, poor vascularization, and improper anatomical location are major

Table 1 Classification of EOC mouse models based on injection site

Advantages	Disadvantages
Subcutaneous model	
√ Easy to perform	√ Model is not physiologically relevant
√ Suitable for monitoring tumor growth	√ Absence of tumor microenvironment
√ Low variability in tumor size	√ Not suitable for studying angiogenesis
√ Good for routine evaluation of drug efficiency	√ Not suitable for studying tumor dissemination
√ Low cost	
Intraperitoneal model	
√ Easy to perform	√ Tumor growth measurements require advanced methods
√ Good model for studying late stages of EOC	√ Not suitable for studying early stages of EOC
√ Suitable for studying EOC dissemination	√ No primary tumor formation
√ Ascites formation	√ Not suitable for studying angiogenesis
√ Suitable for immunological studies	
√ Low cost	
Orthotopic model	
√ Tumor microenvironment recapitulates physiological conditions	√ Difficult to perform
√ Suitable for studying all stages of disease	√ Tumor growth measurement requires advanced methods
√ Suitable for studying angiogenesis and tumor microenvironment	√ High cost
√ Suitable for immunological studies	
√ Good model to study disease progression	
√ Ascites formation	

limitations, which present a barrier to the use of such models in the study of tumor development, invasion, metastasis, angiogenesis, and other processes. Although, subcutaneous xenografts are particularly useful at the early stages of *in vivo* drug-response evaluation, the results obtained need to be confirmed by more advanced physiologically and clinically relevant *in vivo* EOC mouse models.

Orthotopic xenograft models are free from the multiple constraints of subcutaneous xenograft models. Mouse ovaries are encapsulated in a membranous structure called ovarian bursa. Cancer cells can be injected intrabursally, thus allowing tumors to develop in a physiologically relevant microenvironment (Fu & Hoffman, 1993; Kiguchi et al., 1998). Orthotopic xenograft models can be employed to study cancer progression at various stages. Nutrients and signaling provided by the mouse ovary microenvironment allow for tumor development in conditions similar to those in EOC patients. Thus, orthotopic xenograft mouse models are an important tool for understanding the molecular mechanism underlying EOC progression. After the initial phases of tumor growth in the mouse ovarian bursa, cancer cells can invade and disseminate from the primary site to form ascites or metastasis, making it an appropriate model for studies on late-stage EOC progression. Orthotopic xenografts are also widely used in assessing the efficacy of novel therapeutic approaches. The tumor microenvironment plays a key role in determining responses to anticancer agents and contributes to drug resistance. As such, orthotopic xenograft models are an important step in preclinical studies (Hansen et al., 2016). However, despite their notable advantages, these models are technically challenging, which limits their application in EOC research. Intrabursal cell injections require specialized training and are also time consuming and labor intensive compared to other xenograft models. Furthermore, technical challenges result in low implantation success and tumor size variability among mice. These obstacles constrain the use of orthotopic xenograft EOC models, especially for studies on the efficacy of anticancer agents with marginal effects as they require large-sized cohorts to detect significant differences among treatment groups. The inability to monitor tumor growth by conventional methods also makes orthotopic xenograft models less desirable for routine *in vivo* experiments, although this disadvantage can be overcome by using non-invasive, advanced imaging techniques such as *in vivo* bioluminescence and computed tomography (Cordero et al., 2010). Another limitation of intrabursal xenograft models is the anatomical difference between mouse and human ovaries. Unlike the mouse ovary, human ovaries are not covered by bursa, which prevents certain cell lines from leaving the primary site of injection. Metastasis and formation of ascites are primary reasons of lethality among EOC patients, and therefore orthotopic models are not ideal for studies on the late stages of tumor progression.

Intraperitoneal xenograft EOC mouse models are widely used to study cancer cell dissemination. Cancer cells injected into the peritoneal cavity form tumor nodules on the surface of

the liver and spleen, similar to that observed in patients with advanced stages of EOC (Ward et al., 1987). In addition, EOC cells can invade the peritoneum and diaphragm, which makes these models suitable for studying metastasis and dissemination. Ovarian cancer cells growing intraperitoneally can also lead to the development of ascites, one of the key clinical characteristics of EOC in patients (Ahmed & Stenvers, 2013). Ascites development is tightly linked to the immune response against cancer cells and therefore intraperitoneal EOC models are widely used to evaluate the efficacy of anticancer agents modulating ascites production (Shaw et al., 2004). Technical challenges associated with the assessment of tumor growth are a major disadvantage of intraperitoneal EOC xenograft models. Ascites development is highly variable animal to animal and therefore this model requires large groups to assess ascites production statistically.

Classification based on source of cancer cells

EOC xenograft mouse models can be classified by the EOC cell source (Table 2). Established EOC cell lines are the most common source of EOC cells used in *in vivo* mouse models. EOC is a heterogeneous disease represented by several subtypes, including high-grade serous ovarian carcinoma (HGSOC), ovarian clear cell carcinoma (OCCC), ovarian endometrioid carcinoma (OEC), and mucinous carcinoma. HGSOC is the most common type, representing ~70% of EOC cases (Bajrami et al., 2014). For many years, ovarian cancer surface epithelium was considered the primary origin site of EOC (Kurman & Shih Ie, 2010). In the last decade, however, extensive studies have demonstrated that HGSOC most likely originates from fallopian tube fimbria (Kurman & Shih Ie, 2011). This finding highlights the importance of choosing the right EOC cell line for relevant xenograft preclinical models. For example, the two most common cell lines used in earlier EOC xenograft mouse model studies were SKOV3 and A2870, which were later shown to have originated from ovarian surface epithelium and endometrium tissues, and therefore were not representative of HGSOC (Beaufort et al., 2014; Domcke et al., 2013). Inappropriate use of cell lines may contribute, at least partially, to the clinical failure of many previously developed therapeutic strategies. However, in the last decade, around 40 EOC cell lines representing different histo-subtypes of EOC have been characterized. Several studies have classified EOC cell lines based on their origin, histo-subtype, genetic alteration, and ability to grow *in vivo* (Beaufort et al., 2014; Domcke et al., 2013; Hernandez et al., 2016; Ince et al., 2015). These studies represent a valuable source of information and facilitate the choice of cell lines in experimental design. Appropriate choice of EOC cell lines and routine analysis of cell line authenticity have become requirements in the EOC field. Usage of established EOC cell lines *in vivo* has many advantages, including high engraftment rate, fast tumor growth, low tumor size variability, good reproducibility, and ability to use the same models in different research groups. However, despite these advantages, it is important to keep in mind that the majority of EOC cell lines

Table 2 Classification of EOC mouse models based on source of injected cells

Advantages	Disadvantages
Established human EOC cell lines	
√ Availability of cell lines	√ Cell alteration due to high number of passages
√ Large number of cell lines with various genetic backgrounds	√ Not suitable for immunological studies
√ Easy maintenance	
√ Cells can be genetically manipulated	
√ Low cost	
Established mouse EOC cell lines	
√ Availability of cell lines	√ Cell alteration due to high number of passages
√ Suitable for immunological studies	
√ Easy maintenance	
√ Cells can be genetically manipulated	
√ Low cost	
Patient-derived xenografts	
√ PDX derived from EOC patients, not altered by <i>in vitro</i> culture	√ Slow tumor growth
√ Recapitulates EOC tumor microenvironment	√ Difficult maintenance
	√ Specialized training required
	√ Limited access to tumor samples
	√ PDXs cannot be genetically manipulated

have originated from highly advanced tumors. These cells have been exposed to different levels of selection during chemotherapy and recurrence and have accumulated multiple genetic alterations required for survival *in vitro*.

Patient-derived xenografts (PDXs) are the most clinically relevant models currently used in the EOC field (Ricci et al., 2014). PDXs are developed by direct engraftment of EOC patient tissue into immunodeficient mice. Similar to EOC cell lines, EOC specimens can be engrafted subcutaneously, orthotopically, or intraperitoneally. After initial inoculation, PDXs are usually grown for several months before they are harvested for further analysis and transplantation to a new group of animals for *in vivo* experiments. PDXs clinically recapitulate EOC better than any other xenograft model and have gained popularity for preclinical development of novel therapeutic strategies (Heo et al., 2017). Correlations between drug response in patients and PDXs derived from the same patients before treatment have been observed in several studies (Izumchenko et al., 2017; Topp et al., 2014). PDXs are also suitable for studies on aspects of cancer biology that are difficult to study *in vitro* or in other animal models, e.g., vascularization, cancer cell stemness, and tumor microenvironment. However, one of the main disadvantages that PDXs share with other xenograft models is the lack of an immune response in immunocompromised mice. Xenograft mice cannot be used to study the effects of immunotherapies, one of the greatest achievements in cancer therapy in the past several years. This obstacle can be overcome by using humanized immune system mouse models, which utilize immunodeficient mice engrafted with human immune cells or immune cell producing tissues (Choi et al., 2018). Humanized immune system mice are effective at recapitulating an immune response and have gained traction in the development of immune therapies, such as immune checkpoint blockades

(Gitto et al., 2020). Other disadvantages of PDX models are their high cost associated with long-term experiments and technical challenges associated with surgical procedures. In addition, the slow growth rate and lack of stability due to clonal selection during the first several passages limit the use of PDX models within small research groups.

SYNGENEIC EOC MODELS

Syngeneic EOC animal models have recently gained popularity in EOC research. They are particularly useful in studying antitumor immunity and are widely used in developing novel therapeutic strategies based on anticancer agents modulating antitumor immune response (Zhu et al., 2016). Syngeneic models are also suited for studying the tumor microenvironment as both the cancer cells and surrounding tissues belong to the same species (Said et al., 2007). The ID8 cell line is the most widely used in syngeneic EOC mouse models and is derived from C57BL/6 mouse ovarian surface epithelial cells transformed by subsequent serial passage *in vitro* (Roby et al., 2000). Similar to xenograft models, syngeneic models can be classified by the site of injection. Cancer cells, such as the ID8 mouse ovarian cancer cell line, can be injected subcutaneously, orthotopically, or intraperitoneally into C57BL/6 mice depending on experimental needs. The ID8 syngeneic model has been further improved by the introduction of genetic alterations common in EOC such as inactivation of *Trp53*, *Brca1*, and *Brca2* (Walton et al., 2016). Recently, to address EOC heterogeneity, a new generation of syngeneic models have been developed with various genetic backgrounds. These models were developed by the introduction of serial genetic alterations such as loss of *Trp53*, *Brca1*, *Pten*, or *Nf1*, or overexpression of *Ccne1*, *Akt2*, *Kras*, *Brd4*, or *Smarca4* in fallopian tube cell lines, and are

able to better phenocopy EOC characteristics (Iyer et al., 2020; Zhang et al., 2021). These novel syngeneic models should improve our understanding of EOC progression and assist in the development of novel therapeutic strategies such as immunotherapies and PARP inhibition. Other advantages of syngeneic animal models are fast tumor growth, wide availability, and relatively low cost as there is no need to use immunodeficient mice. However, despite these advantages, the immune systems of different species may have substantial variances and therefore any finding obtained using such models will require further validation by other EOC mouse models.

GENETICALLY ENGINEERED MOUSE EOC MODELS

Genetically engineered mouse models (GEMMs) are another important tool for investigating different aspects of EOC pathobiology (Table 3). GEMMs have greatly improved our understanding of EOC initiation. For example, methods such as the *Cre/loxP* system have allowed tissue-specific gene knockin and knockout *in vivo* and have facilitated identification of several EOC driver genes (Hoess & Abremski, 1985). One of the early challenges in the development of EOC GEMMs was lack of knowledge about specific gene promoters associated with tissues involved in EOC development. This obstacle was initially overcome by intrabursal injection of the adenovirus encoding Cre-recombinase (Ad-Cre) under the control of the cytomegalovirus (CMV) promoter (Hardy et al., 1997). Ad-Cre injection into the ovarian bursa results in excision of conditional *LoxP* site-flanked alleles in surrounding tissues (Wu et al., 2007). Alterations in several oncogenes and tumor suppressor genes such as *Trp53*, *Rb1*, *Myc*, *Akt*, *Pik3ca*, *Pten*, and *Arid1a* are believed to be the major driver genes promoting cancer progression in different EOC subtypes (Marcotte et al., 2012). Indeed, early study demonstrated that simultaneous inactivation of *Trp53* and *Rb1*, but not inactivation of either of these genes alone, can lead to the development of EOC resembling HGSOE, with serous histology and metastases to multiple organs (Flesken-Nikitin et al., 2003). Several other studies have also demonstrated that inactivation of other combinations of genes such as *Myc;Trp53;Brca1* (Xing & Orsulic, 2006) or

Pten;Pik3ca (Kinross et al., 2012) can promote HGSOE development. This approach has also been employed for the development of GEMMs recapitulating other EOC subtypes. For example, the discovery of the *Arid1a* mutation as a major driver in the development of OCCC and OEC has been confirmed by several GEMMs (Wiegand et al., 2010). Genetically, loss-of-function *ARID1A* mutation typically co-exists with activation of the PI3K pathway through gain-of-function mutation in *PIK3CA* in OCCC or inactivation of *PTEN* in OEC (Wiegand et al., 2014). Indeed, simultaneous inactivation of *Arid1a* and induction of *Pik3ca* (*Pik3ca^{H1047R}*) mutation through intra-bursal Ad-Cre injection can drive the formation of mouse EOC that fully recapitulates OCCC pathobiology, including development of ascites and peritoneal dissemination (Chandler et al., 2015). In addition, simultaneous inactivation of *Pten* and *Arid1a* can lead to the development of mouse EOC resembling OEC (Guan et al., 2014). Interestingly, *Arid1a* inactivation promotes epithelial differentiation of cancer cells in OEC mouse models driven by concurrent inactivation of *Pten* and *Apc* (Zhai et al., 2016), highlighting the importance of genetic context in determining GEMM phenotypes.

Another approach for EOC GEMM development is selective expression of Cre-recombinase in tissues associated with EOC using tissue-specific gene promoters (Metzger & Chambon, 2001). Müllerian-derived epithelia, including fallopian tube epithelium, one of the tissues believed to be the origin of EOC, can be genetically modified using the *Amhr2* gene promoter (Connolly et al., 2003). For example, the use of the *Amhr2* promoter for Cre-mediated excision of *Dicer1* or *Pten* in combination with *Trp53* in Müllerian-derived epithelia can result in the development of tumors that morphologically resemble HGSOE (Kim et al., 2012). Recently, several other gene promoters specific to Müllerian-derived epithelia, such as *Pax8* and *Ovgp1*, have been used to introduce genetic alterations associated with EOC. Cre-recombinase expression controlled by the *Pax8* promoter has been applied to selectively inactivate *Trp53*, *Brca1*, and *Brca2* in fallopian tubes and endometrium, resulting in the development of EOC histologically similar to HGSOE (Perets et al., 2013). Similarly, tamoxifen-induced expression of Cre-recombinase under the *Ovgp1* gene promoter has been used for inactivation of *Apc*

Table 3 List of selected EOC GEMMs

Gene alteration strategy	Target genes	Phenotype	References
AdCre-intrabursal injection	<i>Trp53;Rb1</i>	HGSOE	Flesken-Nikitin et al., 2003
AdCre-intrabursal injection	<i>Myc;Trp53;Brca1</i>	HGSOE	Xing & Orsulic, 2006
AdCre-intrabursal injection	<i>Pten;Pik3ca</i>	HGSOE	Kinross et al., 2012
AdCre-intrabursal injection	<i>Arid1a;Pik3ca</i>	OCCC	Chandler et al., 2015
AdCre-intrabursal injection	<i>Pten;Arid1a</i>	OEC	Guan et al., 2014
AdCre-intrabursal injection	<i>Arid1a;Pten;Apc</i>	OEC	Zhai et al., 2016
<i>Amhr2</i> -mediated Cre expression	<i>Dicer1;Pten;Trp53</i>	HGSOE	Kim et al., 2012
<i>Pax8</i> -mediated Cre expression	<i>Trp53;Brca1;Brca2</i>	HGSOE	Perets et al., 2013
<i>Ovgp1</i> -mediated Cre expression	<i>Apc;Pten</i>	OEC	Wu et al., 2016
<i>Ovgp1</i> -mediated Cre expression	<i>Trp53;Pten;Brca1</i>	HGSOE	Zhai et al., 2017

and *Pten* in other animal models, resulting in tumors resembling OEC (Wu et al., 2016). Interestingly, tamoxifen-induced expression of Cre-recombinase under the *Ovgp1* gene promoter can inactivate *Trp53*, *Pten*, and *Brca1*, leading to the development of HGSOE (Zhai et al., 2017). Recently, CRISPR/Cas9 technology has allowed much more rapid development of GEMMs at a significantly lower cost compared with traditional breeding protocols. CRISPR/Cas9-mediated knockout or knockin can be performed with mouse embryos of different genetic background, eliminating the need for crossing animals to achieve the desired combination of genetic alterations (Mou et al., 2015). CRISPR/Cas9-mediated gene editing can also be achieved *in vivo* by intrabursal injection of lentiviral particles encoding CRISPR/Cas9 machinery together with gRNA sequences targeting genes involved in EOC development. GEMMs have been widely used for studies on the basic mechanisms of EOC initiation and progression as well as the development of novel therapeutics. Notably, GEMMs allow the exploration of novel therapeutic strategies in an immunocompetent and genetically defined manner. GEMMs are also particularly useful in developing therapeutic approaches that aim to target EOC at the early stages. The major limitations of GEMMs are their inability to monitor tumor growth, high cost associated with breeding colony maintenance, technical difficulties associated with intrabursal injection of the adenovirus encoding Cre-recombinase, and the lack of tissue-specific promoters uniquely suited to drive Cre-expression for the various EOC histo-subtypes.

CONCLUSIONS AND FUTURE PERSPECTIVES

The past decade has seen unprecedented progress in our understanding of the genetic components of EOCs. However, multiple challenges for improving the survival of EOC patients remain. First, due to the lack of effective screening strategies and absence of specific symptoms, most EOC patients are diagnosed with advanced disease. Second, with the advance in knowledge on EOC biology, mechanistic understandings need to be urgently translated into new therapeutic approaches. To meet these challenges, animal models that faithfully recapitulate the initiation, progression, and response to therapeutics are of utmost importance. Although rat and hen spontaneous ovarian cancer models have been reported (Karnezis & Cho, 2017; Lengyel et al., 2014), mouse models are the main experimental platform in EOC research. It is also important to note that there is no superior EOC mouse model, and each model has its own advantages and disadvantages. Therefore, mouse model choice should be determined by the nature of the scientific questions or preclinical experiments on a case-by-case basis. The usage of several mouse models to answer the same scientific question may also increase confidence in *in vivo* observations and increase the potential of translating preclinical findings.

COMPETING INTERESTS

The authors declare that they have no competing interests.

AUTHORS' CONTRIBUTIONS

S.K. and R.G.Z. conceived the review and wrote the manuscript. All authors read and approved the final version of the manuscript.

REFERENCES

- Ahmed N, Stenvers KL. 2013. Getting to know ovarian cancer ascites: opportunities for targeted therapy-based translational research. *Frontiers in Oncology*, **3**: 256.
- Alhilli MM, Becker MA, Weroha SJ, Flatten KS, Hurley RM, Harrell MI, et al. 2016. *In vivo* anti-tumor activity of the PARP inhibitor niraparib in homologous recombination deficient and proficient ovarian carcinoma. *Gynecologic Oncology*, **143**(2): 379–388.
- Bajrami I, Frankum JR, Konde A, Miller RE, Rehman FL, Brough R, et al. 2014. Genome-wide profiling of genetic synthetic lethality identifies CDK12 as a novel determinant of PARP1/2 inhibitor sensitivity. *Cancer Research*, **74**(1): 287–297.
- Beaufort CM, Helmijr JCA, Piskorz AM, Hoogstraat M, Ruigrok-Ritstier K, Besselink N, et al. 2014. Ovarian cancer cell line panel (OCCP): clinical importance of *in vitro* morphological subtypes. *PLoS One*, **9**(9): e103988.
- Byrne AT, Ross L, Holash J, Nakanishi M, Hu LM, Hofmann JI, et al. 2003. Vascular endothelial growth factor-trap decreases tumor burden, inhibits ascites, and causes dramatic vascular remodeling in an ovarian cancer model. *Clinical Cancer Research*, **9**(15): 5721–5728.
- Chandler RL, Damrauer JS, Raab JR, Schisler JC, Wilkerson MD, Didion JP, et al. 2015. Coexistent ARID1A-PIK3CA mutations promote ovarian clear-cell tumorigenesis through pro-tumorigenic inflammatory cytokine signalling. *Nature Communications*, **6**(1): 6118.
- Choi Y, Lee S, Kim K, Kim SH, Chung YJ, Lee C. 2018. Studying cancer immunotherapy using patient-derived xenografts (PDXs) in humanized mice. *Experimental & Molecular Medicine*, **50**(8): 99.
- Connolly DC, Bao RD, Nikitin AY, Stephens KC, Poole TW, Hua X, et al. 2003. Female mice chimeric for expression of the simian virus 40 TAg under control of the *MISIR* promoter develop epithelial ovarian cancer. *Cancer Research*, **63**(6): 1389–1397.
- Cordero AB, Kwon Y, Hua X, Godwin AK. 2010. *In vivo* imaging and therapeutic treatments in an orthotopic mouse model of ovarian cancer. *Journal of Visualized Experiments*, (42): 2125.
- Cordier AC, Haumont SM. 1980. Development of thymus, parathyroids, and ultimo-branchial bodies in NMRI and nude mice. *American Journal of Anatomy*, **157**(3): 227–263.
- Domcke S, Sinha R, Levine DA, Sander C, Schultz N. 2013. Evaluating cell lines as tumour models by comparison of genomic profiles. *Nature Communications*, **4**(1): 2126.
- Flesken-Nikitin A, Choi KC, Eng JP, Schmidt EN, Nikitin AY. 2003. Induction of carcinogenesis by concurrent inactivation of *p53* and *Rb1* in the mouse ovarian surface epithelium. *Cancer Research*, **63**(13): 3459–3463.
- Fu X, Hoffman RM. 1993. Human ovarian carcinoma metastatic models constructed in nude mice by orthotopic transplantation of histologically-intact patient specimens. *Anticancer Research*, **13**(2): 283–286.
- Gitto SB, Kim H, Rafail S, Omran DK, Medvedev S, Kinose Y, et al. 2020. An autologous humanized patient-derived-xenograft platform to evaluate immunotherapy in ovarian cancer. *Gynecologic Oncology*, **156**(1): 222–232.

- Guan B, Rahmanto YS, Wu RC, Wang YH, Wang Z, Wang TL, et al. 2014. Roles of deletion of Arid1a, a tumor suppressor, in mouse ovarian tumorigenesis. *Journal of the National Cancer Institute*, **106**(7): dju146.
- Hansen JM, Coleman RL, Sood AK. 2016. Targeting the tumour microenvironment in ovarian cancer. *European Journal of Cancer*, **56**: 131–143.
- Hardy S, Kitamura M, Harris-Stansil T, Dai Y, Phipps ML. 1997. Construction of adenovirus vectors through Cre-lox recombination. *Journal of Virology*, **71**(3): 1842–1849.
- Heo EJ, Cho YJ, Cho WC, Hong JE, Jeon HK, Oh DY, et al. 2017. Patient-derived xenograft models of epithelial ovarian cancer for preclinical studies. *Cancer Research and Treatment*, **49**(4): 915–926.
- Hernandez L, Kim MK, Lyle LT, Bunch KP, House CD, Ning F, et al. 2016. Characterization of ovarian cancer cell lines as in vivo models for preclinical studies. *Gynecologic Oncology*, **142**(2): 332–340.
- Hoess RH, Abremski K. 1985. Mechanism of strand cleavage and exchange in the Cre-lox site-specific recombination system. *Journal of Molecular Biology*, **181**(3): 351–362.
- House CD, Hernandez L, Annunziata CM. 2014. Recent technological advances in using mouse models to study ovarian cancer. *Frontiers in Oncology*, **4**: 26.
- Hu LM, Hofmann J, Holash J, Yancopoulos GD, Sood AK, Jaffe RB. 2005. Vascular endothelial growth factor trap combined with paclitaxel strikingly inhibits tumor and ascites, prolonging survival in a human ovarian cancer model. *Clinical Cancer Research*, **11**(19 Pt 1): 6966–6971.
- Ince TA, Sousa AD, Jones MA, Harrell JC, Agoston ES, Krohn M, et al. 2015. Characterization of twenty-five ovarian tumour cell lines that phenocopy primary tumours. *Nature Communications*, **6**(1): 7419.
- Iyer S, Zhang S, Yucel S, Horn H, Smith SG, Reinhardt F, et al. 2020. Genetically defined syngeneic mouse models of ovarian cancer as tools for the discovery of combination immunotherapy. *Cancer Discovery*, doi: [10.1158/2159-8290.CD-20-0818](https://doi.org/10.1158/2159-8290.CD-20-0818).
- Izumchenko E, Paz K, Ciznadija D, Sloma I, Katz A, Vasquez-Dunddel D, et al. 2017. Patient-derived xenografts effectively capture responses to oncology therapy in a heterogeneous cohort of patients with solid tumors. *Annual Oncology*, **28**(10): 2595–2605.
- Kamezis AN, Cho KR. 2017. Preclinical models of ovarian cancer: pathogenesis, problems, and implications for prevention. *Clinical Obstetrics and Gynecology*, **60**(4): 789–800.
- Kiguchi K, Kubota T, Aoki D, Udagawa Y, Yamanouchi S, Saga M, et al. 1998. A patient-like orthotopic implantation nude mouse model of highly metastatic human ovarian cancer. *Clinical & Experimental Metastasis*, **16**(8): 751–756.
- Kim J, Coffey DM, Creighton CJ, Yu ZF, Hawkins SM, Matzuk MM. 2012. High-grade serous ovarian cancer arises from fallopian tube in a mouse model. *Proceedings of the National Academy of Sciences of the United States of America*, **109**(10): 3921–3926.
- Kinross KM, Montgomery KG, Kleinschmidt M, Waring P, Ivetac I, Tikoo A, et al. 2012. An activating *Pik3ca* mutation coupled with *Pten* loss is sufficient to initiate ovarian tumorigenesis in mice. *The Journal of Clinical Investigation*, **122**(2): 553–557.
- Kortmann U, Mcalpine JN, Xue H, Guan J, Ha G, Tully S, et al. 2011. Tumor growth inhibition by olaparib in *BRCA2* germline-mutated patient-derived ovarian cancer tissue xenografts. *Clinical Cancer Research*, **17**(4): 783–791.
- Kurman RJ, Shih I M. 2010. The origin and pathogenesis of epithelial ovarian cancer: a proposed unifying theory. *The American Journal of Surgical Pathology*, **34**(3): 433–443.
- Kurman RJ, Shih I M. 2011. Molecular pathogenesis and extraovarian origin of epithelial ovarian cancer-shifting the paradigm. *Human Pathology*, **42**(7): 918–931.
- Lengyel E, Burdette JE, Kenny HA, Matei D, Pilrose J, Haluska P, et al. 2014. Epithelial ovarian cancer experimental models. *Oncogene*, **33**(28): 3619–3633.
- Marcotte R, Brown KR, Suarez F, Sayad A, Karamboulas K, Krzyzanowski PM, et al. 2012. Essential gene profiles in breast, pancreatic, and ovarian cancer cells. *Cancer Discovery*, **2**(2): 172–189.
- Metzger D, Chambon P. 2001. Site- and time-specific gene targeting in the mouse. *Methods*, **24**(1): 71–80.
- Mou HW, Kennedy Z, Anderson DG, Yin H, Xue W. 2015. Precision cancer mouse models through genome editing with CRISPR-Cas9. *Genome Medicine*, **7**(1): 53.
- Perets R, Wyant GA, Muto KW, Bijron JG, Poole BB, Chin KT, et al. 2013. Transformation of the fallopian tube secretory epithelium leads to high-grade serous ovarian cancer in *Brca;Tp53;Pten* models. *Cancer Cell*, **24**(6): 751–765.
- Pla M, Mahouy G. 1991. The SCID mouse. *Nouvelle Revue Francaise D'hematologie*, **33**(6): 489–491.
- Ricci F, Bizzaro F, Cesca M, Guffanti F, Ganzinelli M, Decio A, et al. 2014. Patient-derived ovarian tumor xenografts recapitulate human clinicopathology and genetic alterations. *Cancer Research*, **74**(23): 6980–6990.
- Roby KF, Taylor CC, Sweetwood JP, Cheng Y, Pace JL, Tawfik O, et al. 2000. Development of a syngeneic mouse model for events related to ovarian cancer. *Carcinogenesis*, **21**(4): 585–591.
- Said N, Socha MJ, Olearczyk JJ, Elmarakby AA, Imig JD, Motamed K. 2007. Normalization of the ovarian cancer microenvironment by SPARC. *Molecular Cancer Research*, **5**(10): 1015–1030.
- Shaw TJ, Senterman MK, Dawson K, Crane CA, Vanderhyden BC. 2004. Characterization of intraperitoneal, orthotopic, and metastatic xenograft models of human ovarian cancer. *Molecular Therapy*, **10**(6): 1032–1042.
- Shultz LD, Lyons BL, Burzenski LM, Gott B, Chen XH, Chaleff S, et al. 2005. Human lymphoid and myeloid cell development in NOD/LtSz-*scid* *IL2Rγ^{null}* mice engrafted with mobilized human hemopoietic stem cells. *The Journal of Immunology*, **174**(10): 6477–6489.
- Siegel RL, Miller KD, Jemal A. 2020. Cancer statistics, 2020. *CA: A Cancer Journal for Clinicians*, **70**(1): 7–30.
- The Cancer Genome Atlas Research Network. 2011. Integrated genomic analyses of ovarian carcinoma. *Nature*, **474**(7353): 609–615.
- Topp MD, Hartley L, Cook M, Heong V, Boehm E, McShane L, et al. 2014. Molecular correlates of platinum response in human high-grade serous ovarian cancer patient-derived xenografts. *Molecular Oncology*, **8**(3): 656–668.
- Torre LA, Trabert B, Desantis CE, Miller KD, Samimi G, Runowicz CD, et al. 2018. Ovarian cancer statistics, 2018. *CA: A Cancer Journal for Clinicians*, **68**(4): 284–296.
- Walton J, Blagih J, Ennis D, Leung E, Dowson S, Farquharson M, et al. 2016. CRISPR/Cas9-mediated *Trp53* and *Brca2* knockout to generate

- improved murine models of ovarian high-grade serous carcinoma. *Cancer Research*, **76**(20): 6118–6129.
- Ward BG, Wallace K, Shepherd JH, Balkwill FR. 1987. Intraperitoneal xenografts of human epithelial ovarian cancer in nude mice. *Cancer Research*, **47**(10): 2662–2667.
- Wiegand KC, Hennessy BT, Leung S, Wang YM, Ju ZL, McGahren M, et al. 2014. A functional proteogenomic analysis of endometrioid and clear cell carcinomas using reverse phase protein array and mutation analysis: protein expression is histotype-specific and loss of *ARID1A*/BAF250a is associated with AKT phosphorylation. *BMC Cancer*, **14**(1): 120.
- Wiegand KC, Shah SP, Al-Agha OM, Zhao YJ, Tse K, Zeng T, et al. 2010. *ARID1A* mutations in endometriosis-associated ovarian carcinomas. *New England Journal of Medicine*, **363**(16): 1532–1543.
- Wu R, Hendrix-Lucas N, Kuick R, Zhai YL, Schwartz DR, Akyol A, et al. 2007. Mouse model of human ovarian endometrioid adenocarcinoma based on somatic defects in the Wnt/ β -catenin and PI3K/Pten signaling pathways. *Cancer Cell*, **11**(4): 321–333.
- Wu R, Zhai YL, Kuick R, Karnezis AN, Garcia P, Naseem A, et al. 2016. Impact of oviductal versus ovarian epithelial cell of origin on ovarian endometrioid carcinoma phenotype in the mouse. *The Journal of Pathology*, **240**(3): 341–351.
- Xing DY, Orsulic S. 2006. A mouse model for the molecular characterization of *brca1*-associated ovarian carcinoma. *Cancer Research*, **66**(18): 8949–8953.
- Zhai YL, Kuick R, Tipton C, Wu R, Sessine M, Wang Z, et al. 2016. *Arid1a* inactivation in an *Apc*- and *Pten*-defective mouse ovarian cancer model enhances epithelial differentiation and prolongs survival. *The Journal of Pathology*, **238**(1): 21–30.
- Zhai YL, Wu R, Kuick R, Sessine MS, Schulman S, Green M, et al. 2017. High-grade serous carcinomas arise in the mouse oviduct via defects linked to the human disease. *The Journal of Pathology*, **243**(1): 16–25.
- Zhang S, Iyer S, Ran H, Dolgalev I, Gu SQ, Wei W, et al. 2021. Genetically defined, syngeneic organoid platform for developing combination therapies for ovarian cancer. *Cancer Discovery*, doi: [10.1158/2159-8290.CD-20-0455](https://doi.org/10.1158/2159-8290.CD-20-0455).
- Zhu HR, Bengsch F, Svoronos N, Rutkowski MR, Bitler BG, Allegrezza MJ, et al. 2016. BET bromodomain inhibition promotes anti-tumor immunity by suppressing PD-L1 expression. *Cell Reports*, **16**(11): 2829–2837.