

A Study of Haemodynamic and Aerobic Fitness Profile of Football Players

Kumar¹, Ashok & Singh², Mandeep.

¹Assistant Professor, Department of Sports Science, Punjabi University Patiala-147001, (Punjab), India.
Email: akashokin@gmail.com.

²PG Student, Department of Sports Science, Punjabi University Patiala-147001, (Punjab), India.

Abstract

The purpose of this study was to observe haemodynamic and aerobic fitness profile of football players. Thirty (N=30) male college football players between the age of 17 and 28 years volunteered for this study. The mean age, height and weight of football players was 21±3years, 172.63±8.64cm and 71.56±10.75Kg respectively. The mean resting pulse rate, systolic blood pressure, diastolic blood pressure, pulse pressure, mean arterial pressure, rate pressure product, stroke volume, cardiac output and VO₂ max was 69±4beats/minute, 120±3mmHg, 78±3mmHg, 42.07±4.71mmHg, 91.67±2.94 mmHg, 83.06±5.05beats.min⁻¹.mmHg, 57.38±5.65ml/beat, 3.97±0.28 L/minute and 42.60±2.26 ml.kg⁻¹.min⁻¹ respectively. It was concluded from the results of this study that college level football players have less values of haemodynamic and aerobic fitness variables than elite level football players and athletes of other countries. Thus, they may need to focus on improving haemodynamic and aerobic fitness variables and this could be a focus of their training programme.

Key Words: Mean Arterial Pressure, Stroke Volume, Cardiac Output, VO₂max

Introduction

The performance of the football players depend on different kinds of physiological characteristics. Exceptional endurance performance capacity has long been known as important prerequisite for on-field performance of football players (Ekblom, 1986; Reilly, 1997; Bangsbo et al., 2006). For instance, a player's aerobic endurance capacity facilitates performance continuity, which is limited by endurance, throughout a 90-120 minute game. In addition, it influences the recuperation capabilities following high-intensity games and training units and the recovery following concise high-intensity exercise spurts throughout the games or training units (Ekblom, 1986; Reilly, 1997; Reilly et al., 2008). There is significant agreement that the maximum value for uptake, transport, and use of

oxygen is a good sign of the working of the respiratory, cardiovascular and musculoskeletal systems. This is one of the reasons why research has shown great interest in the determination of haemodynamic and aerobic fitness variables, in a direct or indirect way, facilitating the understanding of physiological aspects related to performance of football players during matches and training programme. Therefore, the purpose of this study was to observe selected haemodynamic and aerobic fitness variables of college football players so that players and coaches could know the status of the physiological adaptations of football training programme.

Materials & Methods

Thirty male college football players of state and inter-university level of participation from Mastuana College of Physical Education, Sangrur (Punjab) India volunteered to participate in this study and their age ranged between 17 to 28 years. Selected anthropometric measurements on the subjects were recorded by using standardized procedure as described by *Weiner & Lourie (1969)*. The aerobic fitness (i.e. $VO_2\text{max}$) was predicted by using resting pulse rate for 20 seconds and enter the number of beats that counted, along with subject's age, into the following equation (*Uth et al., 2004 and Tanaka et al., 2001*).

$$VO_2\text{max (ml/kg/min)} = 15.3 \times (\text{MHR/RHR})$$

$$\text{where, MHR} = \text{Maximum heart rate (beats/minute)} = 208 - (0.7 \times \text{Age})$$

$$\text{RHR} = \text{Resting heart rate (beats/minute)} = 20 \text{ second heart rate} \times 3$$

The haemodynamic variables like Pulse Pressure (PP = SBP- DBP), Mean Arterial Pressure (MAP = Diastolic pressure + 1/3 (Systolic - Diastolic pressure)) (*Andy et al., 1985*), Rate pressure product (RPP = HR x SBP /100) (*Nagpal et al., 2007 & Balogun et al., 1990*) was estimated by using various equations.

Statistical Analysis

Statistical analysis was performed with SPSS version 16.0 (free trial, SPSS Inc, Chicago). Results are shown as Mean and Standard Deviation. The alpha level for the data analysis was determined at $p < 0.05$.

Results & Discussion

Table 1 shows the number and percent distribution of football players according to their playing positions in the present study. As per their playing positions of the football players 43.3% (N=13) were

back, 33.3% (N=10) forward, 16.7% (N=5) center and 6.7% (N=2) goalkeeper.

Table 1: Number of Football Players according to their playing positions

Playing position↓	Number	Percent
1-forward	10	33.3
2-center	5	16.7
3-back	13	43.3
4-goalkeeper	2	6.7
Total	30	100.0

Table 2: Mean ±SD of Haemodynamic & Aerobic Fitness variables of football players

Variables	Mean± Std. Deviation
Age, year	21±3
Height, cms	172.63±8.64
Weight, Kg	71.56±10.75
$VO_2 \text{ max, mL.kg}^{-1}.\text{min}^{-1}$	42.60±2.26
Pulse Rate, beats/minute	69±4
Systolic Blood Pressure, mmHg	120±3
Diastolic Blood Pressure, mmHg	78±3
Pulse Pressure, mmHg	42.07±4.71
Mean Arterial Pressure, mmHg	91.67±2.94
Rate Pressure Product, $\text{beats.min}^{-1}.\text{mmHg}$	83.06±5.05
Stroke Volume, ml/beat	57.38±5.65
Cardiac Output, L/minute	3.97±0.28

The mean age, height and weight of male football players were 21±3 years, 172.63±8.64 cms and 71.56±10.75Kg respectively. The mean Systolic Blood Pressure, Diastolic Blood Pressure, Pulse Pressure, Mean Arterial Pressure, Rate Pressure Product, Stroke Volume and Cardiac Output of football players at rest was 120±3 mmHg, 78±3 mmHg, 42.07±4.71 mmHg, 91.67±2.94 mmHg, 83.06±5.05 $\text{beats.min}^{-1}.\text{mmHg}$, 57.38±5.65 ml/beat and 3.97±0.28 L/minute respectively. The mean VO_2 max and Pulse Rate of football players at rest was 42.60±2.26 $\text{ml.kg}^{-1}.\text{min}^{-1}$ and 69±4 beats/minute respectively (Table 2).

Testing of the haemodynamic and aerobic fitness variables can provide an insight to the football player's current physical capability. Also, assessment of the football player's current level of aerobic fitness reveals strengths and relative weaknesses that can become the basis for the development of an optimal training program (Mirzaei *et al.*, 2009). The results of the measurements in this study were relatively comparable to the values reported in the literature for example McMillian (2005) reported that the mean VO₂max of elite soccer players was between 55 and 68 ml/kg/min but in the present it was 42.60±2.26 ml/kg/min. This value was similar to that reported for in other team sport but was much lower than that of elite endurance athletes (90ml/kg/min). However, players from different positions might have different level of cardiorespiratory level. Reilly *et al.* (2000) reported that among the elite Danish players, full-backs and midfield players possessed the highest values for VO₂max, whereas goalkeepers and central defenders had the lowest but in the present study no such trend related to playing position was observed and this may be due to less number of players, their training or level of participation. It is important to note that the results of aerobic fitness components vary depending upon the relationship of the athletes training schedule and competitive schedule (Roemmich *et al.* 1997). Maximum oxygen uptake is considered to be a valid indicator of the function of respiratory, cardiovascular and muscular systems working together (Impellizzeri & Marcora, 2007). Studies performed on untrained (UT) males have shown that a

good VO₂max is above 40 ml/kg/min; a measure reported above 50 ml/kg/min is considered excellent. In relationship to sport, endurance specific athletes such as cyclists have been shown to have a VO₂max values of the order of 75 ml/kg/min (Saltin & Astrand, 1967). Martial art athletes generally exhibit greater cardiorespiratory endurance than untrained individuals, but not as great as athletes who focus on cardiorespiratory endurance as their primary fitness component for success in their sport, such as cyclists. In regard to martial art disciplines, a study of highly trained competitive black belt karate practitioners were found to have a VO₂max of 57.5 ml/kg/min while the lesser skilled competitive white belt karate practitioners were found to have a VO₂max of 57.2 ml/kg/min (Imamura *et al.*, 1998). By contrast, VO₂max values of 63.8 ml/kg/min were found in 60 England International boxers (Smith, 2006). Wrestlers have been reported to have values for VO₂max of 60.2 ml/kg/min (Yoon, 2002). Crisafulli *et al.* (2009) reported that maximum oxygen uptake for martial art athletes may be as low as 48.5 ml/kg/min and as high as 63.2 ml/kg/min depending upon the specific type of training discipline. Heart rate increases sharply during the first 1–2 minute of exercise, with the magnitude of the increase depending on the intensity of exercise. The increase in heart rate is brought about by parasympathetic withdrawal and activation of the sympathetic nervous system. After approximately 30 min of heavy exercise heart rate begins to drift upward. The increase in heart rate is proportional to the

decrease in stroke volume, so cardiac output is maintained during exercise. Stroke volume exhibits a pattern of initial increase, plateaus, and then displays a negative (downward) drift. Stroke volume increases rapidly during the first minutes of exercise and plateaus at a maximal level after a workload of approximately 40–50% of $\text{VO}_{2\text{max}}$ has been achieved (Åstrand, et al., 1964). As for light to moderate exercise, the increase in stroke volume results from an increased venous return, leading to the Frank-Starling mechanism, and increased contractility owing to sympathetic nerve stimulation. Thus, changes in stroke volume occur because left ventricular end-diastolic volume increases and left ventricular end-systolic volume decreases (Poliner, et al., 1980). Left ventricular end-diastolic volume increases because of the return of blood to the heart by the active muscle pump, increased venoconstriction (which decreases venous pooling, thereby increasing venous return), and increased cardiac output. Left ventricular end-systolic volume decreases owing to augmented contractility of the heart, which effectively ejects more blood from the ventricle, leaving a smaller residual volume. The downward shift in stroke volume after exercise is most likely due to thermoregulatory stress; plasma loss and a redirection of blood to the cutaneous vessels in an attempt to dissipate heat (Rowell, 1986). This effectively reduces venous return and thus causes the reduction in stroke volume. Diastolic blood pressure remains relatively constant because of peripheral vasodilatation, which facilitates blood flow to the working muscles. The rise in systolic blood pressure and the lack of a significant change in diastolic blood

pressure cause the mean arterial pressure (MAP) to rise only slightly, following the pattern of systolic blood pressure. The increase in mean arterial pressure is determined by the relative changes in cardiac output and total peripheral resistance. Since cardiac output increases more than resistance decreases, mean arterial pressure increases slightly during dynamic exercise. However, the increase in mean arterial pressure would be much greater if resistance did not decrease.

Conclusion: It was concluded from the results of this study that college level football players of this study have lower values of haemodynamic and aerobic fitness variables than elite level football players and athletes of other countries. Thus, they may need to focus on improving haemodynamic and aerobic fitness variables and this could be a focus of their training programme.

Acknowledgment: The authors thank all the subjects who voluntarily participated in this study.

References

- Andy J. J., G. O. A. Ladipo and S. P. Owolabi. 1985. Blood pressure distribution of Nigerians in the first two decades of life. *Annals of Tropical Paediatrics*, **5(3)**: 113–118.
- Åstrand, P.; Cuddy, T. E.; Saltin, B.; and Stenberg, J. 1964. Cardiac output during submaximal and maximal work. *Journal of Applied Physiology*, **19(2)**: 268–274.
- Balogun J. A.; Obajuluwa, V. A.; and Abereje. O. K. 1990. Anthropometric determinants of resting blood pressure and heart rate of Nigerian school children," *Annals of Tropical Paediatrics*, **10(4)**: 425–431.
- Bangsbo J.; Mohr, M.; and Krstrup, P. 2006. Physical and metabolic demands of training and match-play in the elite football player. *J Sports Sci*. **24(7)**: 665-674.
- Crisafulli, A.; Vitelli, S.; Cappai, I.; Milia, R.; Tocco, F.; Melis, F.; and Conco. 2009. Physiological responses and energy cost during a simulation of muay thai boxing match.

- Applied Physiology, Nutrition, and Metabolism*. **34**: 143-150.
- Eklom, B. 1986. Applied physiology of soccer. *Sports Med.*, **3**: 50–60.
- Imamura H.; Yoshimura, Y.; Kazuhiro, K.; Nishimura, S.; and Nakazawa, A.T. 1998. Maximal oxygen uptake, body composition and strength of highly competitive and novice karate practitioners. *Appl. Human Sci.*; **17**: 215-218.
- Impellizzeri, F., and Marcora, S. 2007. The physiology of mountain biking. *Sports Medicine*, **37(1)**: 59-71.
- McMillan, K.; Helgerud, J.; Grant, S. J.; Newell, J.; Wilson, J.; Macdonald, R.; Hoff, J. 2005. Lactate threshold responses to a season of professional British youth soccer. *Br. J. Sports Med.*, **39(7)**: 432-436.
- Mirzaei, B., Curby, G., Rahmani-Nia, F. and Moghadasi, M. 2009. Physiological profile of elite Iranian junior freestyle wrestlers. *Journal of Strength and Conditioning Research* **25(8)**: 2339-2344.
- Nagpal, S., Walia, L., Hemlata; Sood, N.; and Ahuja, G.K. 2007. Effect of exercise on Rate Pressure Product in Premenopausal and Postmenopausal women with coronary artery disease. *Indian J. Physiol. Pharmacol.*, **51(3)**: 279-283.
- Poliner, L. R.; Dehmer, G. J.; Lewis, S. E.; Parkey, R. W.; Blomqvist, C. G.; and Willerson, J. T. 1980. Left ventricular performance in normal subjects: A comparison of the responses to exercise in the upright supine positions. *Circulation*. **62**: 528–534.
- Reilly, T., Bangsbo, J., and Franks, A. 2000. Anthropometric and physiological predisposition for elite soccer. *Journal of Sports Science*, **18**: 669-683.
- Reilly, T.; Drust, B.; and Clarke, N. (2008). Muscle fatigue during football match-play. *Sports Med.*, **38(5)**: 357-367.
- Reilly T. 1997. Energetics of high-intensity exercise (soccer) with particular reference to fatigue. *J. Sports Sci.*, **15(3)**: 257-263.
- Roemmich, J.; and Sinning, W. 1997. Weight loss and wrestling training: effects on nutrition, growth, maturation, body composition, and strength. *Journal of Applied Physiology*, **82**: 1751-1759.
- Rowell, L. B. 1986. *Human Circulation: Regulation during Physical Stress*. New York: Oxford University Press.
- Smith M.S. (2006). Physiological profile of senior and junior England international amateur boxers. *J Sports Sci Med.*, CSSI:74-89.
- Tanaka, H.; Monhan, K.D.; and Seals, D.G. 2001. Age-predicted maximal heart rate revisited. *Am. Coll. Cardiol.*, **37**: 153-156.
- Uth, N., Sørensen, H., Overgaard, K.; and Pedersen, P.K. 2004. Estimation of VO₂max from the ratio between HRmax and HRrest-the Heart Rate Ratio Method. *Eur. J. Appl. Physiol.*, **91(1)**: 111-5.
- Weiner J.S. and Lourie J.A.(1969). *Human Biology: A Guide to Field Method*. IBP no.9, Blackwell, London.
- Yoon, J. 2002. Physiological profiles of elite senior wrestlers. *Sports Medicine*, **32(4)**: 225-233.

Conflict of Interest: None declared.



Instructions to Contributors

Journal of Exercise Science and Physiotherapy (JESP) is a scientific peer reviewed journal, publishing research studies and review articles in the field of sports injuries, physiotherapy, exercise physiology, sports rehabilitation, diseases and exercise, sports psychology, nutrition, physical education, sports biomechanics, kinesiology and sports education.

JESP is published annually in June but is planned to be published biannually in the near future. Articles written in English should be sent to the editor of JESP. The research paper for publication in the journal could be sent not only in typed manuscript form but also through the Internet using email attachment. Keeping in view the postal delays, authors, therefore, are encouraged to submit their research material/articles as email directly to satishsk1@rediffmail.com

Mission

JESP aims to present easy access to the scientific knowledge for health & sport conscious individuals. The purpose is to minimise the problems like the delays in publishing process of the articles by drawing advantage from electronic medium.

Instructions to Authors

The articles may be submitted to the Editor, JESP (email: satishsk1@rediffmail.com OR editor@efha-in.org). Manuscripts are considered if they are submitted only to JESP and therefore should not be under consideration for publication elsewhere, either in part or in whole. Authors are responsible for the scientific context and legal aspects of the articles. Editors may make necessary changes to articles in accordance with "Instructions to Authors". Normally the complete research paper should not exceed 15 typed pages including tables and figures; however under exceptional circumstances this limit can be relaxed. There is no page and reference limitation for the review articles. All files related to manuscripts should be submitted electronically. Internet transmission and e-mail attachments are preferred; CD-ROM is acceptable for large files. Text files should be in Microsoft Word 6.0 and later versions. Each figure, table, photograph or other image should be submitted electronically. The manuscripts should be submitted in Times New Roman font, 12-point type, double-spaced with 3 cm margins on all sides. The reference style used by the Journal is the Harvard System of referencing.

Format for Research Article

Include the following sections in research articles without page space:

Title: Capital letters.

Authors: Surname followed by initials and institution(s) where the study was conducted.

Abstract Include an abstract of not more than 150 words that includes objectives, methods, results and conclusions.

KEY WORDS: To assist in indexing the journal, list up to 6 key words (not from title), which your article could be indexed.

INTRODUCTION

METHODS

RESULTS

DISCUSSION

CONCLUSION

ACKNOWLEDGMENTS: Provide information sufficient to identify sources of support, technical assistance, and intellectual contributions not associated with authorship.

REFERENCES: Each citation in the text must be noted by surname and year in parentheses and must appear in the reference section as an alphabetic order. Example for citation in the text; a) for single author (Gür, 1999), b) for two authors (Gür and Akova, 2001), c) more than two authors (Gür et al., 2000).

Journal Article:

Akova, B., Sürmen-Gür, E., Gür, H., Dirican, M., Sarandöl, E. and Küçükoglu, S. 2001. Exercise-induced oxidative stress and muscular performance in healthy women: role of vitamin E supplementation and endogenous estradiol. *European Journal of Applied Physiology*, **84**: 141-147.

Journal Article, Article not in English:

Seker-Aygül, Z., Akova, B. and Gür, H. 2001. The relationship of stress and stress management factors with injury in soccer players. *Turkish Journal of Sports Medicine*, **36**: 71-80. (In Turkish: English abstract).

Journal Article in press:

Gür, H., Cakin, N., Akova, B., Okay, E. and Küçükoglu, S. 2002. Concentric versus combined concentric- eccentric isokinetic training: Effects on functional capacity and symptoms in patients with osteoarthritis of the knee. *Archives of Physical Medicine and Rehabilitation*, in press.

Journal Article in electronic format:

Weigand, D.A., Carr, S., Petherick, C. and Taylor, A. 2001. Motivational climate in Sport and Physical Education: The role of significant others. *European Journal of Sports Science (serial online)* **1(4)**, (13 screens/inclusive page), October. Available from URL: <http://www.humankinetics.com/ejss>

Book:

Guyton, A.C. and Hall, J.E. 1996. *Textbook of medical physiology*. 9th edition. W. B. Saunders Company, London.

Chapter in edited book:

Wilson, C.H. 1984. Exercise for arthritis. In: *Therapeutic exercise*. Ed: Basmajian, J.V. 4th edition. Baltimore: Williams and Wilkins. 529-545.

Thesis:

Özyener, F. 2000. *Effects of work intensity on dynamics of pulmonary gas exchange during exercise in humans*. Doctoral thesis, University of London, London. 79.

Thesis not in English:

Özer, Ö. 2001. *The effect of muscle exercise to oxygen kinetics in chronic smokers*. Doctoral thesis, University of Uludag, Bursa. 1-54. (In Turkish: English abstract).

LEGENDS

AUTHOR BIOGRAPHY: Please include a brief biography, approximately 50 words per author, for each author including (1) academic title/degrees, (2) institution affiliation, (3) research focus, (4) post and e-mail addresses.

TABLES, FIGURES, etc.: The location within the article where e.g. figures and tables appear should be indicated in the text. These materials (e.g. table, figure) should be submitted separately from the text document. Each item should be produced in Microsoft Office (Word, Excel, or PowerPoint), GIF or JPEG format. Submit each item separately and number sequentially as (main author) Figure 1.jpg, (main author) Figure2.jpg, and so on, for example, GurFigure1.jpg and GurTable1.doc.

FORMAT FOR REVIEW ARTICLE

Include the following sections in research articles without page space:

TITLE: Capital letters.

AUTHORS: Surname followed by initials and institution(s) where the study was conducted.

ABSTRACT Include an abstract of not more than 300 words.

KEY WORDS: To assist in indexing the journal, list up to 6 key words (not from title), which your article could be indexed

MAIN TEXT

CONCLUSION

ACKNOWLEDGMENT

REFERENCES: Each citation in the text must be noted by surname and year in parentheses and must appear in the reference section as an alphabetic order. Example for

citation in the text; a) for single author (Gür, 1999), b) for two authors (Gür and Akova, 2001), c) more than two authors (Gür et al., 2000). See the reference section of research article for detail.

LEGENDS

AUTHOR BIOGRAPHY: For detail see the author biography section of research article.

TABLES, FIGURES, etc.: For details see the tables, figures, section of research article.

REVIEW AND PUBLISHING POLICY

Upon receipt of a manuscript the editor will notify the main author that the submission has been received. The manuscript will undergo a blinded review process. Under this process the editor will forward (electronically) the manuscript to two independent reviewers. Normally the review process will take approximately 4 weeks. After the review process has been completed the main author will be notified via E-mail either of acceptance, revision or rejection of the manuscript. If the main author has not been informed of the results of the review process after 8 weeks from the date of confirmation of receipt of the manuscript (by the editor), the author should notify the editor-in-chief directly. If revision was recommended, this should be completed and returned to the editor within 4 weeks; failure to return the revised manuscript within the allotted time span will lead to rejection of the manuscript. When a manuscript has been accepted for publication it will be forwarded to the editor-in-chief for final approval. The editor-in-chief will work with the main author to make necessary revisions in order for the manuscript to appear in the Journal of Exercise Science and Physiotherapy. The final proof of the manuscript will be sent to the main author as an E-mail attachment. If the main author requires any minor changes to the finished work, these should be submitted electronically to the editor-in-chief within one week. Manuscript accepted in the first half of the year will be published not later than the second half of the year.

Checklist for Authors

- Double-space manuscript text and use 3cm margins on all sides, and Times New Roman, 12-point type, English and SI units.
- Include with text a title, abstract, key words, figure and table lists, figure and table legends.
- Submit all materials electronically.
- Include complete author names, addresses, e-mail, phone and fax numbers.
- Include a short (~50 words) author biography for each author.

**Editor-in-Chief,
JESP**