



A Comprehensive Review of Different Liver Toxicants Used in Experimental Pharmacology

Bigoniya P. *, Singh C. S., Shukla A.

Department of Pharmacology, Radharaman College of Pharmacy, Fatehpur Dobra, Ratibad, Bhopal- 462 002, M.P., India

ABSTRACT

Liver is a primary organ involved in metabolism of food and drugs. Liver disorders are mainly caused by toxic chemicals, such as antibiotics, chemotherapeutic agents, peroxidised oil, aflatoxin, carbon tetrachloride, chlorinated hydrocarbons etc. Toxic liver injury produced by drugs and chemicals may virtually mimic any form of naturally occurring liver disease. In view of multiplicity and complexity of the liver functions, it is obvious that no single test can establish the disturbances in liver function. Thus, a battery of liver function tests is employed for accurate diagnosis, to assess the severity of damage, to judge prognosis and to evaluate therapy. To study hepatoprotective potential of any herbal product, isolated phytochemical or synthetically developed moiety induction of experimental liver damage is prerequisite. The selection of toxicant depends on type and nature of liver damage required as every toxicant has its unique mode of action producing specific type of destruction. The hepatoprotective mechanism of any drug under study can be explained based on its protective behavior against different toxicants. This review is precise compilation of experimental liver pharmacology taking concern dose, route, and schedule of different liver toxicants along with evaluation biochemical, histological and functional ability parameters. The literature compilation will definitely help researcher in design and assessment of studies involving liver function.

Keywords: Hepatotxicity, liver toxicants, biochemical parameters, liver function tests, liver histopathology, bromosulphthalein.

INTRODUCTION

The liver is the largest organ in the body and serves many vital functions such as remove damaged red blood cells from the blood in co-ordination with spleen, produces bile, clotting factors, stores vitamins, minerals, protein, fats and glucose from diet. [1-2] The most important task of the liver is to filter toxic substances from the body, like alcohol, chemotherapeutic drugs, antibiotics and toxicants. If accumulation of toxins is faster than the liver metabolizing ability, hepatic damage may occur.

Besides the chemicals and toxicant, there are several factors that increase the risk of hepatic injury that includes different races, as blacks and Hispanics may be more susceptible to isoniazid toxicity, alcohol ingestion, elderly persons have increased risk of hepatic injury because of decreased clearance, drug to drug interactions and reduced hepatic blood flow. Females are more prone to hepatic injury.

***Corresponding author: Dr. Papiya Bigoniya,**

Department of Pharmacology, Radharaman College of Pharmacy, Fatehpur Dobra, Ratibad, Bhopal- 462 002, M.P., India **Email:** p_bigoniya2@hotmail.com

Unique gene encodes in each P₄₅₀ protein, long acting drugs, host factors, persons suffering from AIDS, malnourished and fasting person may be susceptible to hepatic injury because of low glutathione stores. [3]

The liver metabolizes virtually every drug or toxin introduced in the body. Metabolism of drugs or toxins occurs in 2 phases. [4] In the phase 1 reaction, the drug is made polar by oxidation or hydroxylation. All drugs may not undergo this step, and some may directly undergo the phase 2 reaction. The cytochrome P₄₅₀ enzymes catalyze phase 1 reactions. [5-6] Most of these intermediate products are transient and highly reactive. These reactions may result in the formation of metabolites that are far more toxic than the parent substrate and may result in liver injury. Phase 2 reactions may involve conjugation with a moiety (i.e., acetate, amino acid, sulfate, glutathione, glucuronic acid). [7] Subsequently, drugs with high molecular weight may be excreted in bile, while the kidneys excrete the smaller water soluble molecules. Several chemical agents like phenobarbital, phenytoin, carbamazepine, primidone, ethanol, glucocorticoids, rifampin, griseofulvin, quinine and omeprazole induces P₄₅₀ enzyme activity, whereas in

contrast, amiodarone, cimetidine, erythromycin, isoniazid, ketoconazole, metronidazole, sulfonamides and quinidine inhibit the P₄₅₀ enzyme.

Some hepatotoxins found in nature are the products of plants (e.g. albitocin, cycasin, pyrrolizidines, saffrole, tannic acid, indospicine), fungal (e.g. aflatoxin, phalloidin, luteoskyrin)^[8] or bacterial metabolism (e.g. *Corynebacterium diphtheriae*, *Clostridium botulinus*, *Streptococcus hemolyticus* and some strains of *E. coli*) or minerals. Many hepatotoxic agents are products of chemicals or pharmaceutical industry. Others are industrial byproducts or waste materials that by polluting the environment may cause hepatotoxicity in humans as well as animals.^[9] Hepatic injury can have several forms, classified as:

- Acute hepatitis or chronic hepatitis (inflammatory liver disease)
- Hepatositis (non-inflammatory disease)
- Cirrhosis (degenerative disorder resulting in fibrosis of the liver)

Some agents interfere with bile secretion with little or no overt injury to hepatic parenchyma. Other leads to necrosis, cirrhosis or carcinoma. Some toxicants produce curious degeneration or vascular lesion. Drugs or chemicals can produce the entire range of known hepatic lesion.

MECHANISMS OF HEPATIC INJURY

The mechanism of hepatic injury can be categorized as pathophysiological mechanism and chemical induced mechanism.

Pathophysiological hepatic damage

Disruption of the hepatocyte

Covalent binding of the chemical agent and toxicant to intracellular proteins can cause a decrease in ATP levels, leading to actin disruption. Disassembly of actin fibrils at the surface of the hepatocyte causes blebs and rupture of the membrane.^[10]

Disruption of the transport proteins

Chemical agents and toxicants that affect transport proteins at the canalicular membrane can interrupt bile flow.^[11] Loss of villous processes and interruption of transport pumps such as multidrug resistance-associated protein-3 prevent excretion of bilirubin, causing cholestasis.

Cytolytic T-cell activation

Covalent binding of chemical agents and toxicants to the P₄₅₀ enzyme acts as an immunogen, activating T cells, cytokines and stimulating a multifaceted immune response.^[12]

Apoptosis of hepatocytes

Activation of the apoptotic pathway by the tumor necrosis factor-alpha receptors (TNF- α) of Fas may trigger the cascade of intercellular caspases, which results in cell death.^[13]

Mitochondrial disruption

Certain chemical agents inhibit mitochondrial function by a dual effect on both beta-oxidation energy productions by inhibiting the synthesis of nicotinamide adenine dinucleotide and flavin adenine dinucleotide, resulting in decreased ATP production.^[14]

Bile duct injury

Toxic metabolites excreted in bile may cause injury to the bile duct epithelium.

Chemical induced hepatic damage

Liver disorders are mainly caused by toxic chemicals, such as antibiotics chemotherapeutic agents, peroxidised oil, aflatoxin, CCl₄, chlorinated hydrocarbons etc. Most of the

hepatotoxic chemicals damage liver cells mainly by inducing lipid peroxidation and by generation of reactive oxidative intermediates in liver.^[15]

Direct or intrinsic or predictable drug reactions

Drug or one of its metabolites that fall into this category is either cause reproducible direct toxicity to the liver or lowers the host defense mechanism. The adverse effects occur in most individuals who consume them in dose-dependent manner e.g. carbon tetrachloride.

Indirect or unpredictable idiosyncratic drug reactions

Drugs that fall into this group cause immune mediated toxicity, which is independent of concentrations. The drug or one of its metabolites acts as a hapten and induces hypersensitivity in the host. The hepatotoxicity by this group of agents does not occur regularly in all individuals and the effects are usually not dose related e.g. acetaminophen. The drugs causing particular type of liver diseases are tabulated in Table 1.

Table 1: Drug induced liver diseases

| LIVER DISEASE | | AGENTS | |
|--------------------------------|--------|--|--|
| Acute hepatitis | viral | Acebutolol, Indomethacin, Phenylbutazone, Allopurinol, Isoniazid, Phenytoin, Atenolol, Ketoconazole, Piroxicam, Carbamazepine, Quinine, Diltiazem, Naproxen, Ranitidine, Enflurane, Para-aminosalicylic acid, Sulfonamides, Ethambutol, Penicillins, Sulindac, Labetalol, Probenecid, Cimetidine, Maprotiline, Pyrazinamide, Dantrolene, Metoprolol, Quinidine, Diclofenac, Mianserin, Ethionamide, Phenelzine, Tricyclic antidepressants, Halothane, Phenindione, Valproic acid, Ibuprofen, Phenobarbital, Verapamil | |
| | | Adrenocortical steroids, Phenothiazines, Sulfonamides, Antithyroid drugs, Phenytoin, Tetracyclines, Isoniazid, Salicylates, Valproic acid, Methotrexate | |
| Acute fatty liver infiltration | | Actinomycin D, Chlorpropamide, Erythromycin, Amoxicillin/Clavulanate, Cloxacillin, Flecaïnide, Azathioprine, Cyclophosphamide, Flurazepam, Captopril, Cyclosporine, Flutamide, Carbamazepine, Danazol, Glyburide, Carbimazole, Diazepam, Gold, Cephalosporins, Disopyramide, Griseofulvin, Chlordiazepoxide, Enalapril, Haloperidol, Ketoconazole, Norethandrolone, Sulfonamides, Mercaptopurine, Oral contraceptives, Tamoxifen, Methyltestosterone, Oxacillin, Thiabendazole, Nifedipine, Penicillamine, Tolbutamide, Nitrofurantoin, Phenothiazines, Tricyclic antidepressants, Nonsteroidal, Phenytoin, Troleandomycin, Anti-inflammatory drugs, Propoxyphene, Verapamil | |
| | | Gold, Phenytoin, Aspirin, Hydralazine, Procainamide, Carbamazepine, Isoniazid, Guinidine, Chlorpromazine, Quinidine, Nitrofurantoin, Sulfonamides, Diltiazem, Penicillin, Tolbutamide, Disopyramide, Phenylbutazone | |
| Chronic hepatitis | active | Acetaminophen, Dantrolene, Methyl dopa, Isoniazid, Nitrofurantoin | |
| | | Methotrexate, Terbinafine HCl, Nicotinic acid | |
| Liver cirrhosis or fibrosis | | Chlorpromazine/valproic acid (combination), Imipramine, Thiabendazole, Phenothiazines, Tolbutamide, Chlorpropamide/Erythromycin (combination), Phenytoin | |
| | | Anabolic steroids, Oral contraceptives, Thorotrast, Danazol, Testosterone | |
| Liver tumors | | | |

MODE OF ACTION OF LIVER TOXICANTS USED IN EXPERIMENTAL PHARMACOLOGY

Paracetamol

Paracetamol is chemically characterized as N-acetyl-para-aminophenol, which is also known as acetaminophen. Acetaminophen is a safe and effective analgesic in recommended dose. However, acetaminophen is also well

characterized, dose dependent hepatotoxin that can lead to life-threatening acute liver failure when excessive doses are ingested. [16] The hepatotoxic mechanism include the formation of a reactive metabolite, presumably N-acetyl-p-benzoquinamine (NAPQI) through cytochrome P₄₅₀ pathway [17], which is quickly conjugated with hepatic glutathione to yield a harmless product called mercapturic acid. However, after acetaminophen overdose, the capacity for glucuronidation and sulfation is exceeded with the formation of excess NAPQI via cytochrome P450 2E1. This then leads to depletion in glutathione, excess of NAPQI binds to hepatic cell protein and DNA resulting in mitochondrial dysfunction [18] and development of acute hepatic necrosis.

Several P₄₅₀ enzymes such as P450 2E1 play an important role in acetaminophen bioactivation to NAPQI. [19] Studies demonstrated that acetaminophen induced hepatotoxicity can be modulated by substances that influence cytochrome P₄₅₀ activity. [20] Paracetamol induced hepatotoxicity causes rise in SGOT, SGPT, ALP and bilirubin with extensive vascular degenerative changes and centrilobular necrosis in hepatocytes (Fig. 1).

Ethanol

Ethanol is an alcohol that is used commonly as a solvent in medications. Toxicity occurs when an excessive amount is ingested. Alcohol is mostly metabolized in the liver through a series of chemical reactions known as oxidation reactions. In the predominant biological pathway for alcohol metabolism, known as alcohol dehydrogenase pathway, the enzyme alcohol dehydrogenase converts alcohol to a toxic intermediate substance, acetaldehyde [21] by removing two atoms of hydrogen from each alcohol molecule. Then a second enzyme, aldehyde dehydrogenase, quickly converts acetaldehyde to acetate [22] by again removing hydrogen and adding oxygen. A secondary pathway of alcohol metabolism is microsomal ethanol-oxidizing system (MEOS). MEOS is activated by long-term heavy alcohol consumption. [23] The MEOS pathway involves the enzyme cytochrome P450 2E1 or CYP 2E1 that strips hydrogen away from alcohol to produce acetaldehyde. [24] In both of these pathways, more markedly in the MEOS pathway-oxidation reactions spawn highly unstable free oxygen radicals.

Normally the body deploys molecules called antioxidants to clear oxygen radicals from the liver. However heavy alcohol use not only heightens the production of oxygen radicals but also depletes the supply of antioxidants in the body, creating an imbalance between oxygen radicals and antioxidants. This imbalance is known as oxidative stress which damages liver cell membranes and mitochondria. When oxidative stress is chronic, it contributes to necrosis and liver fibrosis. In addition to its direct effects on the liver, oxidative stress appears to stimulate autoimmune reactions that further damage liver cells. [25] In ethanol induced hepatotoxicity there is significant rise in SGOT, SGPT, ALP and total bilirubin and reduction in total protein, albumin and total cholesterol. The rats treated with ethanol show hepatic cords, fatty infiltration, mesenchymal hyperplasia, fibrosis (Fig. 2a) and fatty infiltration of hepatocytes, hyperplasia of connective tissue, and early manifestation of cirrhosis (Fig. 2b).

Thioacetamide

Thioacetamide is a thiono-sulfur containing compound having the fungicidal property. Reif *et al.* 1999 reported that intraperitoneal injection of thioacetamide produces potent

hepatotoxicity and carcinogenesis. [26] Chronic administration of thioacetamide produces liver cirrhosis. Thioacetamide is metabolized by cytochrome P₄₅₀ enzymes of liver microsomes and is converted to a toxic intermediate called thioacetamide S-oxide due to oxidation process. [27] Thioacetamide S-oxide induced oxidative stress in the hepatic cells. [28] It is responsible for the changes in cell permeability, increase in intracellular concentration of Ca⁺⁺, increase in nuclear volume, enlargement of nucleoli and also inhibits mitochondrial activity which leads to cell death and severely affects those cells which are located in the perivenous acinus region. [29] Thioacetamide hepatotoxicity showed significant rise in SGOT, SGPT, ALP and total bilirubin and decrease in total protein. Thioacetamide causes pericellular hepatocyte necrosis, inflammation, infiltration of leukocytes with cytoplasmic vacuolation (Fig. 3).

Isoniazid

Isoniazid is chemically isonicotinylnhydrazine that is used to prevent and treat tuberculosis. Isoniazid can cause adverse effects on the liver, ranging from mild transient elevations in aminotransferases to overt hepatitis. N-acetyltransferase (NAT2) metabolizes isoniazid to acetyl isoniazid, which is then hydrolyzed to acetylhydrazine. [30] Acetylhydrazine is further metabolized by CYP 2E1 to produce hepatotoxic derivatives. Metabolic oxidation of acetylhydrazine leads to formation of a reactive acylating species that binds covalently to microsomal protein. Acetylhydrazine and hydrazine act as acylating agents by binding with the liver cell macromolecules, causing hepatocyte injury. [31] Isoniazid increases SGOT, SGPT, ALP and bilirubin, while decreasing total protein and albumin. The rats treated with isoniazid show hepatocellular disintegration and vacuolation in the centrilobular region (Fig. 4).

Carbon tetrachloride (CCl₄)

Liver toxicant CCl₄ causes lipid peroxidative degradation of biomembrane, which is one of the principal causes of hepatotoxicity. [32] In liver CCl₄ is biotransformed by cytochrome P₄₅₀ to produce its active metabolite trichloromethyl free radical (CCl₃*) [33], which binds to the macromolecule and induces peroxidative degradation of membrane lipids of endoplasmic reticulum rich in polyunsaturated fatty acids. This leads to the formation of lipid peroxide which then gives toxic aldehyde that causes damage to the liver. Secondary mechanisms link carbon tetrachloride metabolism that could promote the generation of toxic products arising directly from carbon tetrachloride metabolism or from peroxidative degeneration of membrane lipids. The possible involvement of toxic intermediates radical species are such as trichloromethyl (CCl₃*), trichloromethylperoxy (OCCl₃*) and chlorine (Cl*) free radicals as well as phosgene and aldehydic products of lipid peroxidation.

This radical can bind to cellular molecules (nucleic acid, protein and lipid) impairing crucial cellular processes such as lipid metabolism, with the potential outcome of fatty degeneration (steatosis). This radical can also react with oxygen to form a highly reactive species, trichloromethylperoxy radical (CCl₃OO*). CCl₃OO* initiates the chain reaction of lipid peroxidation which attacks and destroys polyunsaturated fatty acids, in particular those associated with phospholipids. This affects the permeability of mitochondrial, endoplasmic reticulum and plasma membranes resulting in the loss of cellular calcium

sequestration and homeostasis, which can contribute heavily to subsequent cell damage. Carbon tetrachloride significantly increases SGOT, SGPT, ALP and total bilirubin where as decreasing total protein, albumin and total cholesterol. Rat liver tissue treated with carbon tetrachloride shows hepatocellular necrosis, fatty vacuole and microvesicular fatty changes (Fig. 5).

Galactosamine

Galactosamine is a hexosamine derived from galactose. It causes liver injury via the generation of free radicals and depletion of UTP nucleotides. Galactosamine produces the hepatotoxic effect by selectively reducing the uridine pool in hepatocytes. This in turn inhibits mRNA and protein synthesis, alters the composition of cellular membranes and finally leads to cellular damage as a result of lipid peroxidation. [34] The hepatocyte death is represented as apoptosis and subsequently necrosis. [35] Other mechanism of galactosamine hepatotoxicity stated that galactosamine increases intestinal permeability and subsequently facilitates bacterial translocation to the liver. [36] Lipopolysaccharides activate kupffer cells to secrete tumor necrosis factor- α [37], which raises expression of intercellular adhesion molecule 1 in endothelial cells [38] and this promotes the adhesion of polymorphonuclear cells to vascular [39] and hepatic endothelial cells [40], leading to polymorphonuclear infiltration and hepatocyte damage. Galactosamine induces rise in SGOT, SGPT and total bilirubin where as decrease in total protein. Galactosamine shows pathological changes like moderate degeneration and necrosis of hepatocyte (Fig. 6).

Cadmium

Cadmium metals and metalloids affect almost every organ of the body, including the liver. One such metal is cadmium, which is of concern because of its increasing prevalence as an environmental contaminant. [41] Prolonged exposure to cadmium results in injury to the liver. A large bolus dose of cadmium causes injury to a number of tissues, including the liver. [42] Cadmium induces oxidative damage in different tissues by enhancing peroxidation of membrane lipids in tissues and altering the antioxidant systems of the cells. The peroxidative damage to the cell membrane may cause injury to cellular components due to the interaction of metal ions with the cell organelles. [43] Cadmium depletes glutathione and protein bound sulfhydryl groups resulting in enhanced production of reactive oxygen species such as superoxide ions, hydroxyl radicals, and hydrogen peroxides. These reactive oxygen species result in increased lipid peroxidation, [44] hepatic congestion, ischemia and hypoxia. The resultant ischemic hypoxia leads to neutrophil infiltration, kupffer cell activation, and inflammation which could potentially contribute to the widespread hepatocellular apoptosis and necrosis. [45] Cadmium causes increase in serum concentrations of urea, creatinine, glucose, AST, acid phosphatase, alkaline phosphatase, alanine transaminase, aspartate transaminase, and serum bilirubin where as reducing serum protein and tissue protein concentration. Cadmium treated rat shows the histopathological changes like periportal inflammation (Fig. 7a), microvesicular steatosis and balloon degeneration (Fig. 7b).

Aflatoxin B1 (AFB1)

AFB1 is a naturally occurring fungal toxin that causes both acute hepatotoxicity and liver carcinoma in humans and animals. AFB1 produces the hepatotoxicity through the formation of adducts with DNA, observed both in vitro and

in rat liver. [46] These adducts are derived from highly reactive exo-epoxide metabolites of AFB1, as a result of oxidation reactions within the liver. [47] Several cytochromes P₄₅₀ have been implicated in this activation and in human these were identified as CYP 1A2 and CYP 3A4. [48] The formation of adducts between the AFB1-exo-epoxide and the N7 of guanine bases in DNA leads to mutations and is strongly associated with the presence of preneoplastic lesions, characterized as GST-P-positive foci, providing a direct link between DNA damage, mutagenesis and carcinogenesis. AFB1 causes acute toxicity as well as carcinogenicity in rats and as observed in early studies. [49] Acute toxicity was initially attributed to mainly genotoxic effects of the epoxide; dependent on the formation of DNA adducts which at high levels lead to cell death. However, a dialdehyde metabolite of AFB1 that rapidly forms from the epoxide, can form adducts with proteins and these were proposed to contribute to the acute toxicity. [50] In addition, such cellular necrotic damage caused by AFB1 dialdehyde may lead to compensatory liver hyperplasia and by so doing may promote the incorporation of mutations into the DNA of dividing cells and contribute towards carcinogenicity initiated by the AFB1-exo-epoxide. [51] AFB1 increases serum concentrations of SGOT, SGPT, alkaline phosphatase and bilirubin, and decrease in serum cholesterol. The prominent gross pathologic and histopathologic changes in the liver are hemorrhage, necrosis, and massive accumulation of lipid (Fig. 8).

Allyl alcohol

The toxicity of allyl alcohol is considered to be mediated via acrolein, which is generated from allyl alcohol by the enzyme alcohol dehydrogenase. [52] Acrolein is a highly toxic member of a class of α - β unsaturated aldehydes, namely 2-alkenals. [53] Acrolein, is a powerful electrophile and reacts with nucleophiles such as sulphhydryl groups. [54] The reaction is accelerated by the activity of cytosolic glutathione S-transferase [55] to form an aldehyde-GSH adducts which is metabolized to acrylic acid. Glutathione is primarily involved in the reaction, which result in a depletion of cellular glutathione stores, followed by hepatocellular necrosis. [56] Allyl alcohol induces increase in SGOT, SGPT and total bilirubin where as decrease in total protein. The rats treated with allyl alcohol shows necrosis around branches of the central hepatic vein and presence of a large amount of nuclear debris (Fig. 9).

Halothane

Halothane is chemically 2-bromo-2-chloro-1,1,1-trifluoroethane. It has been used widely as an inhaled anaesthetic [57] and as liver toxicant in animal models. It is well established that halothane is metabolized in the liver as a lipophilic xenobiotic to hepatotoxic intermediates by monooxygenases through the cytochrome P450 2E1 system. [58] Thus, halothane anaesthesia causes hepatocellular necrosis, destruction of the lipid-protein interactions in human erythrocyte membranes, decrease in activities of membrane enzymes and alteration of cerebral glucose-6-phosphate dehydrogenase (E.C.1.1.1.49, G6PDH) activities. [59] Halothane treated rat liver shows extensive centrilobular necrosis and denaturation (Fig. 10).

BIOCHEMICAL ALTERATIONS IN HEPATIC DAMAGE

The biochemical changes in the blood reflect the histologic pattern of toxicity in hepatic injury. Biochemical alterations provide a static assessment of the degree of liver injury.

Serum aminotransferase enzymes

Serum concentrations of aspartate aminotransferase (AST) or glutamate oxaloacetate transaminase (SGOT) and alanine aminotransferase (ALT) or glutamate pyruvate transaminase (SGPT) are the most commonly used biochemical markers of hepatocellular necrosis. [60] These serum activities presumably increase as a result of cellular membrane damage and leakage. [61] Serum levels of SGOT and SGPT are increased on damage to the tissues producing them. Serum aminotransferase activities are increased in all types of hepatic injury. Thus serum estimation of SGPT which is fairly specific for liver tissue is of greater value in liver cell injury, whereas SGOT level may raise in acute necrosis or ischemia of other organs such as the myocardium, besides liver cell injury. The highest increases are observed with acute hepatocellular injuries, such as xenobiotic-induced necrosis or acute viral hepatitis. [62] Serum activities are generally within the reference interval or only slightly increased in alcoholic liver disease. [63]

Serum alkaline phosphatase

Serum alkaline phosphatase is produced by many tissues especially bone, liver, intestine, placenta and is excreted in the bile. Serum alkaline phosphatase increases to some extent in most types of liver injury. Bile acids induce alkaline phosphatase synthesis and exert a detergent effect on the canalicular membrane, allowing leakage into serum. [61, 64] The highest concentrations are observed with cholestatic injuries. [60]

Serum total protein and albumin

Proteins form the major portion of dissolved substances in the plasma. Liver cells synthesize albumin, fibrinogen, prothrombin, alpha-1-antitrypsin, haptoglobin, ceruloplasmin, transferrin, alpha fetoproteins and acute phase reactant proteins. The blood levels of these plasma proteins are decreased in extensive liver damage.

Serum albumin, the major plasma protein synthesized in the human liver, is a clinically useful marker of hepatic synthetic function. [60] Alcoholic cirrhosis with or without accompanying ascites generally lowers serum albumin concentrations. [61]

Serum total and direct bilirubin

The serum bilirubin level is one of the best tests of liver function. Bilirubin is the metabolic product of the break down of heme derived from senescent red blood corpuscles. An increase in urinary bilirubin is nearly always indicative of a corresponding increase in the serum, attributable to intrahepatic or extrahepatic cholestasis. The degree of increase in serum bilirubin values has prognostic significance in chronic liver injuries, but not in acute injuries. [65] If the direct or conjugated bilirubin is low, while the total bilirubin is high, this reflects liver cell damage or bile duct damage.

Serum bile acids

Measurement of bile acid concentrations is a good indicator of hepatobiliary function. An increase in serum bile acid concentrations in fasting is highly specific for liver injury and serves to exclude congenital or hemolytic causes of hyperbilirubinemia. [66] The greatest increases are observed in acute viral hepatitis or extrahepatic cholestasis. [60]

Serum lipid profile

Cholesterol is the main lipid found in the blood, bile and brain tissues. Liver metabolizes the cholesterol and it is transported to the blood stream by lipoproteins. Decreased levels are found in malabsorption, malnutrition, hyperthyroidism, anemia and liver diseases.

Triglycerides are simple lipids formed in the liver by glycerol and fatty acids. They are transported by very low density lipoprotein (VLDL) and low density lipoprotein (LDL). They constitute about 90 % of fat, stored as source of energy in the tissue and plasma.

Serum γ -glutamyl transferase

The measurement of serum γ -glutamyl transferase is a frequently used parameter of liver diseases. The serum enzyme originates from liver and is cleared from the circulation by the galactose receptor in liver. The rate of uptake will thus vary with the amount of terminal galactose residues on the enzymes carbohydrate moiety. The enzyme is inducible by chronic alcohol use liver abscess and by drugs such as phenytoin. [67]

Serum 5'-nucleotidase

5'-Nucleotidase (5NT) is an intrinsic membrane glycoprotein produced by the liver present as an ectoenzyme in a wide variety of mammalian cells, hydrolyzes 5'-nucleotides to their corresponding nucleosides. Despite its ubiquitous distribution, serum concentrations of 5NT appear to reflect hepatobiliary disease with considerable specificity. [68] The primary utility of serum 5'-nucleotidase activities is in the diagnosis of cholestatic liver injury in childhood or pregnancy.

Serum lactic dehydrogenase

Lactate dehydrogenase, also called lactic dehydrogenase (LDH), is an enzyme found in the cells of many body tissues, including the heart, liver, kidneys, skeletal muscle, brain, red blood cells, and lungs. It is responsible for converting muscle lactic acid into pyruvic acid, an essential step in producing cellular energy. Lactic dehydrogenase is present in almost all body tissues, so the LDH test is used to detect tissue alterations and as an aid in the diagnosis of heart attack, anemia, and liver disease.

Alpha-fetoprotein

Alpha-fetoprotein (AFP) is a protein normally produced by the fetal liver and is present in the fluid surrounding the fetus (amniotic fluid), and crosses the placenta into the mother's blood. At birth, infants have relatively high levels of AFP, which fall to normal adult levels by the first year of life. AFP probably has no normal function in adults. The most widely used biochemical blood test for liver cancer is AFP.

Mitochondrial antibodies

Anti-mitochondrial antibodies (AMA) represent a heterogeneous mixture of antibodies to at least 9 different antigens, which are designated M1-M9. Primary biliary sclerosis (PBC) is an autoimmune disease that causes destruction of intrahepatic bile ducts. PBC is often associated with other autoimmune disorders, particularly Sjogren syndrome. Anti-mitochondrial antibodies (AMA) are found in almost all patients with PBC and are considered the serological hallmark of the disease. AMA is useful diagnostically in distinguishing primary biliary cirrhosis from other types of chronic liver disease. [69]

HISTOPATHOLOGICAL ALTERATION IN HEPATIC DAMAGE

The liver reacts with eight different types of responses to injury towards variety of metabolic, toxic, microbial,

circulation and neoplastic insults. [70] Damage from toxic or immunologic insult may cause hepatocytes to take on a swollen and edematous appearance (ballooning degeneration) with irregularly clumped cytoplasm and large clear spaces. Alternatively, retained biliary material may impart a diffuse foamy swollen appearance to the hepatocyte (foamy degeneration). Substances may accumulate in viable hepatocytes including iron and copper. Accumulation of fat droplets within hepatocytes is known as 'steatosis' and appear in such conditions as alcoholic liver disease and acute fatty liver of pregnancy. A single large droplet that displaces the nucleus (macrovesicular steatosis) may be seen in the alcoholic liver or in the liver of obese or diabetic individuals. Necrosis is defined as focal death along with degradation of tissue by hydrolytic enzymes liberated by cells. Various agents such as hypoxia, chemical, physical agents, microbial agents and immunological injury can cause necrosis. Two essential changes bring about irreversible cell injury in necrosis is cell digestion by lytic enzymes and denaturation of proteins. These processes are morphologically identified by characteristic cytoplasmic and nuclear changes in necrotic cell. The cytoplasm appears homogeneous and intensely eosinophilic. Occasionally, it may show vacuolation or dystrophic calcification. The nuclear changes include condensation of nuclear chromatin which may either undergo dissolution or fragmentation into many granular clumps. Centrilobular necrosis frequently exhibits a zonal distribution. The most obvious necrosis of hepatocytes immediately around the terminal hepatic vein, an injury that is characteristic of ischemic injury occurred for a number of drug and toxic reactions. A variable mixture of hepatocellular death and inflammation is encountered. The hepatocyte necrosis may be limited to scattered cells within hepatic lobules. Bridging necrosis is more severe inflammatory injury; necrosis of continuous hepatocytes may span adjacent lobules in a portal to portal, portal to central and central to central fashion. Submassive necrosis of entire lobules or most of the liver is usually accompanied by hepatic failure. With disseminated candidal or bacterial infection, macroscopic abscesses may occur. In fat necrosis, the necrosed fat cells have cloudy appearance, surrounded by an inflammatory reaction. Formation of calcium soaps is identified in the tissue sections as amorphous, granular and basophilic material. Microscopically, fibrinoid necrosis is identified by brightly eosinophilic, hyaline-like deposition in the vessel wall or on the hepatocytes. All the histopathological changes characteristic of different types of liver diseases is tabulated in Table 2.

COMMON LIVER FUNCTION TESTS

Liver function tests routinely combine markers of function (albumin and bilirubin) with markers of liver damage (alanine transaminase, alkaline phosphatase, and γ -glutamyl transferase). Abnormalities in liver enzyme activities give useful information about the nature of the liver insult: a predominant rise in alanine transaminase activity (normally contained within the hepatocytes) suggests a hepatic process. These tests can also distinguish between acute and chronic liver disorders and between hepatitis and cholestasis. The most commonly performed blood tests include:

Serum glutamate oxaloacetate transaminase (SGOT) test

This enzyme is released from damaged liver, heart, muscle, kidney or brain cells. Its normal serum level is up to 46 IU/L at 37°C. [15] SGOT levels are 10 to 200 fold elevated in

patients with acute hepatic necrosis, viral hepatitis, CCl₄ and drug induced poisoning. SGOT levels are also elevated by 10 fold in patients of post hepatic jaundice, intra hepatic cholestasis and less than 10 fold in alcoholic and hepatic steatosis. [71]

Serum glutamate pyruvate transaminase (SGPT) test

This enzyme is released from damaged liver cells. Normal serum level of SGPT is up to 49 IU/L at 37°C and its levels are very high in patients of viral hepatitis and hepatic necrosis, 10 to 200 fold higher in patients of post hepatic jaundice, intrahepatic cholestasis and below 10 fold in patients of metastatic carcinoma, cirrhosis and alcoholic hepatitis. [72]

Serum alkaline phosphates test

Elevated levels of alkaline phosphatase, an enzyme found in the bile, usually indicate an obstruction of bile flow, liver injury, or certain cancers (Dial, 1995). Elevation in normal serum alkaline phosphatase (range 3-13 King Armstrong units/dl or 25-85 IU/dl) activity is found in diseases of bone, liver and in pregnancy. In the absence of bone disease or pregnancy, an elevated serum alkaline phosphatase level generally reflects hepatobiliary disease. The greatest elevation (3-10 times of normal) occurs in biliary tract obstruction. Slight to moderate increase is seen in parenchymal liver diseases such as hepatitis, cirrhosis and metastatic liver disease. [15]

Serum total protein and albumin test

Routinely estimated total proteins are in the normal range of 5.5 to 8 g/dl. The blood levels of plasma protein are decreased in extensive liver damage. Albumin (normal range 3.5 to 5.0 g/dl) synthesized in the liver constitutes a major part of the total proteins in the body and the other part being globulin. A low serum albumin concentration suggests chronic liver disease. [73] Hypoalbuminaemia may occur in liver diseases caused by significant destruction of hepatocytes. Hyperglobulinaemia may be present in chronic inflammatory disorders such as in cirrhosis and in chronic hepatitis.

Serum total and direct bilirubin test

Each day about 7.5 g of hemoglobin is catabolized with the corresponding production of 250 mg of bilirubin. Normally 0.25 mg/dl of conjugated bilirubin is present in the blood of an adult. Normal range for total bilirubin is from 0.2 to 1.2 mg/dl. Bilirubin level rises in diseases of hepatocytes, obstruction to biliary excretion into duodenum, hemolysis and in defects of hepatic uptake and conjugation of bilirubin such as in Gilberts disease. [15] Elevated levels of bilirubin often indicate an obstruction of bile flow or a defect in the processing of bile by the liver. If the direct or conjugated bilirubin is low, while the total bilirubin is high, this reflects liver cell damage or bile duct damage. [74]

Serum bile acids

Fasting bile acids concentrations in excess of 15 μ mol/L can be the result of hepatobiliary disease. The probability of hepatobiliary disease is high if fasting bile acid concentration is greater than 25-30 μ mol/L.

Serum lipid profile test

Liver toxicants cause disturbances in synthesis and metabolism of triglycerides, cholesterol and lipoproteins, thus damaging the basic resource for living cells. Cholesterol and bile salts are synthesized by liver cells thus liver intoxication decreases level. Normal range of cholesterol levels are up to 200 mg/dl or lower for a total count, but it is

Table 2: Characteristic histopathological changes in different types of liver diseases

| LIVER DISEASE | CAUSES | CHARACTERISTIC HISTOPATHOLOGICAL CHANGE |
|--------------------------------------|---|---|
| Hepatitis | | |
| Acute | Alcohol abuse and hepatotropic viral infection | Cell injury in centrilobular zone, ballooning degeneration, apoptosis, dropout necrosis and bridge necrosis, inflammation infiltrate in portal tracts and kupffer cell hyperplasia |
| Chronic active | Autoimmune chronic active hepatitis, hepatitis B and C virus, hepatitis B virus with delta infection, Wilson's disease, haemochromatosis, alcohol and alpha-1-antitrypsin deficiency | Intense portal inflammatory infiltrate spills over hepatic parenchyma, piecemeal necrosis, and bridging necrosis between adjacent triads, progressive fibrosis beginning in the portal triads and radiating into the liver parenchyma |
| Chronic persistent | Hepatitis B and C virus and combined hepatitis B and delta infection | Lymphocyte and macrophage infiltration is limited to the portal triads and no significant hepatocyte necrosis |
| Chronic liver disease | | |
| Liver cirrhosis | Alcohol abuse, hepatitis, viruses, certain drugs, chemicals, bile duct obstruction, autoimmune diseases, obstruction of blood outflow from liver, alpha-1-antitrypsin deficiency, high blood tyrosine levels at birth, diabetes, malnutrition and Wilson's Disease or hemochromatosis | Diffuse damage to hepatic parenchyma cells, with nodular regeneration, fibrosis, and disturbance of normal architecture |
| Liver fibrosis | Infection, inflammation and injury | Pericellular, perivenular fibrosis and web appearance |
| Autoimmune liver disorders | | |
| Hemochromatosis | Absorption of too much iron from food | No histopathological change |
| Wilson's disease | Retention of too much copper in the liver | No histopathological change |
| Alcohol-induced liver disease | | |
| Fatty liver | Excessive consumption of alcohol | Swelled hepatocyte, deposition of large fat droplets and formation of microvesicles, fatty changes in centrilobular and mid-zone and diffuse change Collagen laid down around the cells and hepatic veins - pericellular and perivenular fibrosis. |
| Alcoholic hepatitis | Excessive consumption of alcohol | Ballooning degeneration and necrosis of hepatocytes near the central veins, retention of secretory proteins and water, causing cell swelling and formation of Mallory bodies Hepatocytes may show slight increased cytoplasmic haemosiderin |
| Alcoholic cirrhosis | Excessive consumption of alcohol | |
| Congenital liver defects | | |
| Biliary atresia | Absence or abnormally developed bile ducts | No histopathological change |
| Choledochal cyst | Malformation of the hepatic duct | No histopathological change |
| Benign Liver tumors | | |
| Hemangioma | Birth defect | Tumor is found as hemorrhagic lesion composed of blood spaces of various sizes covered with single layer of endothelial cells separated by thin fibrous stroma |
| Focal nodular hyperplasia | Congenital vascular malformation or vascular injury | Central stellate scar with radiating fibrous septa, regions of nodular hepatocellular proliferation separated by radiating bands and surrounding myxomatous scar |
| Hepatocellular adenoma | Use of oral contraceptive pills with a high oestrogen content | Intratumoral hemorrhage, large intratumoral vessels, fatty change and peliosis |
| Malignant liver tumors | | |
| Hepatocellular carcinoma | Prolonged infection with Hepatitis B and C virus, alcohol, mycotoxin and chemical carcinogens | Large solid tumor mass separated by vascular space, inconspicuous sinusoids and central cystic space formed by degeneration |
| Haemangio-sarcoma | Vinyl chloride monomers, thorotrast and arsenic | Cystic degeneration, haemorrhage and necrosis |
| Hepatoblastoma | Beckwith-Wiedemann syndrome, familial adenomatous polyposis and hemihypertrophy. Inborn errors of metabolism such as tyrosinemia, glycogen storage disease type I, galactosemia and alpha-1-antitrypsin deficiency | Circumscribed tumor having areas of cystic degeneration, haemorrhage and necrosis |

important to check HDL and LDL levels for a better analysis. Fatty degeneration of the liver causes increased triglyceride (normal range >150 mg/dl) content in the blood.

γ -glutamyl transferase test

Gamma glutamyl transferase (γ -GT) also known as γ -glutamyl transpeptidase is a microsomal enzyme with wide tissue distribution. This enzyme is produced by the liver, pancreas, kidneys and released into the blood when these organs are injured. The normal γ -GT serum level is upto 26 U/L. In alcoholics with liver abscess, it is increased by 2-5 times.

5-Nucleotidase test

Normal range of 5-Nucleotidase is 2 to 17 U/L. The liver releases this enzyme when injured due to bile duct obstruction or impaired bile flow. Greater than normal values indicate liver cell destruction, liver ischemia, necrosis, hepatitis, cholestasis or liver tumor.

Lactic dehydrogenase test

Reference ranges for total LDH vary from laboratory to laboratory. Normal values are also higher in childhood. For adults, in most laboratories, the range can be up to approximately 200 U/L, but is usually found within 45-90 U/L. When disease or injury affects tissues containing LDH, the cells release it into the bloodstream, identified as higher than normal levels. The LDH is also elevated in heart attack, diseases of the liver, in certain types of anemia, and in cases of excessive destruction of cells, as in fractures, trauma, muscle damage and shock.

Alpha-fetoprotein test

In adults, high blood levels (over 500 ng/ml) of AFP are seen in only three situations like hepatocellular carcinoma, germ cell tumors and metastatic cancer in the liver. Also, pregnant women carrying babies with neural tube defects may have high levels of AFP.

Mitochondrial antibodies test

AMA are present in less than 1 % of normal people and in less than 5% of patients with systemic lupus erythematosus,

rheumatoid arthritis and other autoimmune diseases. Patients with extrahepatic biliary obstruction, Wilson's disease, hemochromatosis, and alcoholic cirrhosis rarely have elevated titers. The presence of mitochondrial antibodies remains a useful diagnostic tool in the differential diagnosis between primary biliary cirrhosis and extrahepatic biliary obstruction.

Prothrombin time (PTT) test

This test measures the time it takes for blood to clot. Blood clotting requires vitamin K and coagulation factors like II, V, VII, and IX synthesized in the liver. Liver cell damage and bile flow obstruction can both interfere with proper blood clotting.

EXPERIMENTAL PHARMACOLOGICAL STUDIES IN ANIMAL LIVER

To investigate and evaluate hepatoprotective substance, it is customary to subject animals to a range of toxic agents. These hepatotoxicants include carbon tetrachloride, galactosamine, thioacetamide, ethanol, aflatoxin B1, alpha amanitine, phalloidin, cadmium, paracetamol, hydrazine, halothane, isoniazid etc that causes damage of rat liver, resulting in biochemical and histopathological changes. Different toxicants used for experimental liver damage with dose range, route, vehicle and detailed schedule of treatment are presented in Table 3.

Table 3: Different toxicants used for experimental liver damage with dose, route, vehicle and detailed schedule of treatment

| TOXICANT | VEHICLE | ANIMAL | ROUTE | DOSE | SCHEDULE | REF |
|----------------------|------------------------------|--------------------|------------|---|---|--------|
| Paracetamol | 0.5 % CMC or 0.2% tragacanth | Wistar albino rat | I.P. | 600 mg/kg | Daily treatment for 14 days | 75 |
| Ethanol | Sterile water | Wistar albino rat | Oral | 500 mg/kg to 3 gm/kg | Single dose on 3 rd , 7 th or 8 th day | 76, 77 |
| Thioacetamide | Sterile water or saline | Wistar albino rat | S.C. | 1.5 gm/kg to 12 gm/kg or 15% v/v or 5 ml/100 gm | 30 days to 45 days | 78, 79 |
| | | Sprague-Dawley rat | I.P. | 100 mg/kg | Single dose on 21 st day of drug treatment | 80 |
| Isoniazid | Sterile water or saline | Rabbit | I.P. | 50 mg/kg | Single dose on every 24 th hr for 3 consecutive days | 81, 82 |
| Carbon tetrachloride | Olive oil | Wistar albino rat | S.C. | 1 ml/kg | Single dose on 7 th day or 4 consecutive days | 85, 86 |
| | | Wistar albino rat | S.C. | 0.15 ml/kg | There times a week for 10 weeks | 87 |
| Galactosamine | Saline | Swiss albino rat | I.P. | 650 mg/kg to 800 mg/kg | Single dose | 88, 89 |
| Cadmium chloride | Sterile water or saline | Swiss albino rat | S.C. | 3 mg/kg | Every alternate days for 2 weeks | 90 |
| | | Wistar albino rat | Oral | 6 mg/kg | One month to three months | 91 |
| | | Sprague-Dawley rat | I.P. | 3.5 mg/kg | Single dose on 7 th day of treatment | 92 |
| | | Sprague-Dawley rat | I.V. | 4 mg/kg | Single dose on 3 rd day of treatment | 93 |
| Aflatoxin B1 | 8% dimethyl-sulfoxide | Sprague-Dawley rat | I.P. | 1 mg/kg to 2 mg/kg | Single dose | 94, 95 |
| Allyl alcohol | Sterile water or saline | Wistar albino rat | Oral | 0.4 mg/kg | Single dose of toxicant administered on 3 rd day | 96 |
| Halothane | 21% with O ₂ | Swiss albino mice | I.P. | 75 µl/kg | Single dose | 97 |
| | | Guinea pig | Inhalation | 1% v/v | Inhalation for 4 hrs | 98, 99 |




Fig. 1




Fig. 2a




Fig. 2b




Fig. 3




Fig. 4




Fig. 5

Fig. 1: Extensive vascular degenerative changes and centrilobular necrosis induced by paracetamol. Fig. 2: Ethanol intoxication induces fatty infiltration, mesenchymal hyperplasia and fibrosis (Fig. 2a), and hyperplasia of connective tissue along with early manifestation of cirrhosis (Fig. 2b). Fig. 3: Thioacetamide causes perilobular hepatocyte necrosis, infiltration of leukocytes with cytoplasmic vacuolation. Fig. 4: Isoniazid shows hepatocellular disintegration and vacuolation in the centrilobular region. Fig. 5: Carbon tetrachloride treatment shows hepatocellular necrosis, fatty vacuole and microvesicular fatty changes.



Fig. 6




Fig. 7a




Fig. 7b




Fig. 8




Fig.9



Fig. 10

Fig. 6: Galactosamine shows pathological changes like moderate degeneration and necrosis of hepatocyte. Fig. 7: Cadmium treated rat shows the histopathological changes like periportal inflammation (Fig. 7a), microvesicular steatosis and ballooning degeneration (Fig. 7b). Fig. 8: The prominent gross pathologic and histopathologic changes like hemorrhage, necrosis, and massive accumulation of lipid induced by Aflatoxin B1. Fig. 9: The rats liver treated with Allyl alcohol shows necrosis around branches of the central hepatic vein and presence of a large amount of nuclear debris. Fig. 10: Halothane causes extensive centrilobular necrosis and denaturation.

Animals

Hepatotoxicity studies can be carried out in different strains of mice and rats of either sex maintained at uniform laboratory conditions. Animals should be housed at a temperature of $25\pm 2^{\circ}\text{C}$, relative humidity of $50\pm 15\%$ and 12:12, light: day. All animals are allowed to free access to water and fed with standard commercial pellet rat chaw. The animals are acclimatized for a period of 7 days before performing the experiment. Inhalation anaesthetic induced hepatotoxicity studies are usually done on guinea pigs. Rabbits are choice of species for drug induced chronic hepatotoxicity study.

Methodology

Before the commencement of experiment the animals are kept fasted overnight and then divide into several groups. One group will serve as a vehicle control group, receiving vehicle only. Hepatotoxicant (dissolve in the suitable solvent as depicted in Table 3) is administered to animals of the other groups through the suitable route as per the study protocol. The standard drug is administered to different group serves as positive control and only toxicant is administered to another group designating negative control. The test drug is administered in varying concentration or dose for a specified duration depending on the design of the experiment. At the end of experiment, all the animals are sacrificed after a predecided period (18-48 h after last dose) under light ether anesthesia or by direct cervical dislocation. Biochemical estimation is performed on serum by drawing blood either from carotid artery or through cardiac puncture in hyphenised tube. The blood samples are left to clot at room temperature for atleast 1 h. Serum is separated by centrifugation at 3000 rpm for 20 min at 4°C to carry out the different assay. In some cases the blood can also be collected from retino-bulbar venous plexus under light ether anesthesia for the biochemical estimation. For histopathology liver is removed,

washed with infusion of cold saline and weighed. Liver sections are taken from each lobe of liver, which are immediately frozen and stored at -70°C or fixed in 10 % normal formaline until analyzed.

Biochemical estimation

The activities of serum glutamate oxaloacetate transaminase (SGOT or AST) and serum glutamate pyruvate transaminase (SGPT or ALT) are measured in serum according to the methods described by Reitman and Frankel. [100] Lactase dehydrogenase (LDH) activity is measured by the method of Varley. Alkaline phosphatase (ALP) activity is determined according to Kind and King. [101] Total bilirubin and urea levels are estimated according to Malloy and Evlyn. [102] Plasma protein concentration was measured according to the method of Bradford. [103] Hepatic glucose-6-phosphatase is determined in the soluble fraction of liver homogenates according to the described by Harper. [104] Acid phosphatase (ACP) activity in liver was assayed using a Boehringer kit. The activity of albumin and total cholesterol are measured according to the method of Varley. [105] Hepatic lipid peroxidation is assayed by measuring the concentration of malondialdehyde (MDA) in 10% w/v liver homogenate according to the procedure of Okhawa. [106] Glutathione-S-transferase content in liver cytosolic fraction is determined by the method of Ellman. [107]

Histopathological observation

After fixing in 10 % neutral formalin solution, liver tissues are dehydrated with ethanol solution, embedded in paraffin, cut into 5 μm section, stained with haematoxylin-eosin dye and then observed under a photomicroscope. Different hepatotoxicants show the different morphological changes that may be disturbance of normal architecture of parenchyma cells, swelled hepatocyte, and formation of mallery bodies, microvesicles sinusoidal congestion, infiltration of lymphocytes, kupffer cells around the central

vein and loss of cell boundaries. Histopathological changes may also include massive fatty changes, ballooning degeneration, lymphocyte, macrophage infiltration, nodular regeneration, steosis, fibrosis, and tumor formation.

Assessment of liver function

Functional ability of liver is assessed on thiopentone induced sleeping time, bromosulphthalein clearance and viability by trypan blue exclusion test.

Barbiturate induced sleeping time

Hepatic damage can be measured by accessing activity of hepatic microsomal drug metabolizing enzymes. The hepatotoxicant attacks the membranes of smooth and rough endoplasmic reticulum thus reducing the quantity of microsomal enzymes. Intoxicated liver prolongs duration of sleeping time for hexobarbitone, thiopentone, zoxazolamine and pentobarbitone etc in animals. Thiopentone, hexobarbitone, or pentobarbitone induced sleeping time is increased in animals with liver intoxication as the enzyme responsible for metabolism of thiopentone is reduced or destroyed. ^[108]

Bromosulphthalein uptake test

The liver cells remove the dye, bromosulphthalein (BSP) from the plasma and excrete it into bile. It has been reported that BSP is secreted into bile as mercaptide conjugate of glutathione and cystein, which takes place in liver. Bromosulphthalein clearance test is the most sensitive and dependable method to assess the physiological status of liver function. The test indicates the excretory function of the liver. It is generally agreed that in the passage of BSP from the plasma to the bile, it undergoes storage, metabolism, and excretion by the liver. Concentration of BSP in plasma greater than 5.8 mg/ml is considered as indicative of liver damage after 15 min of 100 mg/kg (i.v.) BSP administration. ^[109]

BSP clearance rate can be estimated spectrophotometrically *in-vitro* in isolated liver slices following method of Rajan and Subrahmanyam. ^[110]

Trypan blue exclusion test

Loss of cell viability is most often measured as loss of membrane integrity. This event may be primarily due to necrosis or secondarily due to apoptosis. Trypan blue exclusion is a cell viability assay based on the ability of the liver cells to exclude the trypan blue and uptake of the dye by the dead cells due to alteration in the membrane permeability which can be measured following method of William *et al.*, 1971. ^[111]

While undertaking hepatoprotective potential assessment of a drug or chemical the study parameters should be well designed covering all aspects of liver functionality. Effectiveness of a particular drug against a specific type of liver disease can only be claimed when it is experimentally proved in preclinical animal studies. Preclinical study reports are also necessary for approval to carry out clinical studies. To study hepatoprotective potential of any herbal product, isolated phytochemical or synthetically developed moiety induction of experimental liver damage is prerequisite. Liver damage can be caused by exposure to a range of toxic chemicals. Toxic liver injury produced by drugs and chemicals may virtually mimic any particular form of naturally occurring liver disease. Dose and duration of exposure with toxicants are also crucial in concern to achievement of acute to chronic type of liver damage. The

selection of liver toxicant must depends on targeted nature of liver damage required.

The hepatoprotective potential of any promising drug can be evaluated by treating the experimental animal along with the toxicant and assessing functional parameters of liver. A range of liver function tests is employed for accurate diagnosis of disease prognosis and therapy evaluation. This review precisely compiles the details of different liver toxicants used in experimental pharmacology like, dose, route, and along with mechanism of damage. The literature compiles biochemical alterations and histological characteristic damage of liver cytology in detail. Different functional ability test and viability assessment studies can be referred from the review. This extensively cited and well documented review will definitely help the researcher in liver protective study protocol preparation and cross referencing the published methods.

REFERENCE

1. Dyce KM, Sack WO, Wensing CJ. Text Book of Veterinary Anatomy. W B Saunders Co, Philadelphia, 1987, 542-595.
2. Waugh A, Grant A. Ross and Wilson Anatomy and Physiology in Health and Illness. Edn 9, Churchill Livingstone, Spain, 2001, 307-311.
3. Lemke TL, William DA, Roche VF, Zito SW. Foye's Principles of Medicinal Chemistry. Edn 6, Lippincott Williams & Wilkins, USA, 2007, 253-325.
4. Park BK, Pirmohamed M, Kitteringham NR. The role of cytochrome P450 enzymes in hepatic and extrahepatic human drug toxicity. *Pharmacology and Therapeutics* 1995; 68: 385-424.
5. Smith G, Stubbins MJ, Harries LW, Wolf CR. Molecular genetics of the human cytochrome P450 monooxygenase superfamily. *Xenobiotica* 1998; 28: 1129-1165.
6. Werck-Reichhart D, Feyereisen R. Cytochromes P450: a success story. *Genome Biology* 2000; 1(6): 3003.1-3003.9.
7. Williams RT. Detoxification Mechanisms. Edn 2, Chapman & Hall, London, 1959, 114-126.
8. Wieland T, Wieland O. Chemistry and toxicology of the toxins of *Amanita phalloides*. *Pharmacological Reviews* 1959; 11: 87-107.
9. Zimmerman HJ, Lewis JH. Chemical and toxin induced hepatotoxicity. *Gastroenterology Clinics of North America* 1995; 24(4): 1027-1045.
10. Watanabe S, Phillips MJ. Acute phalloidin toxicity in living hepatocytes: evidence for a possible disturbance in membrane flow and for multiple functions for actin in the liver cell. *American Journal of Pathology* 1986; 122: 101-111.
11. Trauner M, Meier PJ, Boyer JL. Molecular pathogenesis of cholestasis. *New England Journal of Medicine* 1998; 339: 1217-1227.
12. Cullen JH. Mechanistic classification of liver injury. *Toxicologic Pathology* 2005; 33: 6-8.
13. Faubion WA, Guicciardi ME, Miyoshi H, Bronk SF, Roberts PJ, Svingen PA, Kaufmann SH, Gores GJ. Toxic bile salts induce rodent hepatocyte apoptosis via direct activation of Fas. *Journal of Clinical Investigation* 1999; 103: 137-145.
14. Pessayre D, Mansouri A, Haouzi, D, Fromenty B. Hepatotoxicity due to mitochondrial dysfunction. *Cell Biology and Toxicology* 1999; 15: 367-373.
15. Harshmohan. Textbook of Pathology. Edn 5, Jaypee Brothers: Medical Publishers (P) Ltd., New Delhi, 2005, 22-24 and 608-668.
16. Makin AJ, Wendon J, Williams R. A 7-year experience of severe acetaminophen-induced hepatotoxicity (1987-1993). *Gastroenterology* 1995; 109: 1907-1916.
17. Dahlin D, Miwa G, Lee A. N-acetyl-p-benzoquinonamine: a cytochrome P450 dependent oxidation product of acetaminophen. *Proceeding of the National Academy Science* 1984; 81: 327-331.
18. Parmar D, Kandakar M. Mitochondrial ATPase: A target for paracetamol induced hepatotoxicity. *European Journal of Pharmacology* 1995; 293: 225-229.
19. Raucy JL, Lasker JM, Lieber CS, Black M. Acetaminophen activation by human liver cytochromes P450 1E1 and P450 1A2. *Archives of Biochemistry and Biophysics* 1989; 271: 270-283.

20. Mitchell JR, Jollow DJ, Potter WZ, Davis DC, Gillette JR, Brodie BB. Acetaminophen-induced hepatic necrosis. I. Role of drug metabolism. *Journal of Pharmacology and Experimental Therapeutics* 1973; 187: 185-194.
21. Satoru K, Susumu I, Toyoka H, Yoshiyasu Y, Takeyuki M, Yoshihiko F. Acetaldehyde as well as ethanol are metabolized by human CYP2E1. *Journal of Pharmacology and Experimental Therapeutics*. 1997; 280: 527-532.
22. Koivula T, Koivusalo M. Different forms of rat liver aldehyde dehydrogenase and their subcellular distribution. *Biochimica et Biophysica Acta* 1975; 397: 9-23.
23. Lieber CS, Decarli LM. Hepatic microsomal ethanol oxidizing system. *Journal of Biological Chemistry* 1970; 254: 2505-2512.
24. Asai H, Imaoka S, Kuroki T, Monna T, Funae Y. Microsomal ethanol oxidizing system activity by human hepatic cytochrome P450. *Journal of Pharmacology and Experimental Therapeutics* 1996; 277: 1004-1009.
25. Albano E, French S, Ingelman SM. Cytochrome P450 2E1, hydroxyethyl radicals and immune reactions associated with alcoholic liver injury. *Alcoholism: Clinical and Experimental Research* 1988; 22: 740-742.
26. Reif S, Weis B, Aeed H. The Ras antagonist, farnesylthiosalicylic acid (FTS), inhibits experimentally-induced liver cirrhosis in rats. *Journal of Hepatology* 1999; 31: 1053-1061.
27. Kim KH, Bae JH, Cha SW, Han SS, Park KH, Jeong TC. Role of metabolic activation by cytochrome P450 in thioacetamide-induced suppression of antibody response in male BALB/c mice. *Toxicology Letters* 2000; 114: 225-235.
28. Zaragoza A, Andres D, Sarrion D, Cascales M. Potentiation of thioacetamide hepatotoxicity by phenobarbital pretreatment in rats, inducibility of FAD monooxygenase system and age effect. *Chemico-Biological Interactions* 2000; 124: 87-101.
29. Amad A, Pillai KK, Najmi AK, Pal SN. Evaluation of hepatoprotective potential of jigrine post-treatment against thioacetamide induced hepatic damage. *Journal of Ethnopharmacology* 2002; 79: 35-41.
30. Huang YS, Chern HD, Su WJ. Cytochrome P450 2E1 genotype and the susceptibility to antituberculosis drug-induced hepatitis. *Hepatology* 2003; 37: 924-930.
31. Noda A, Hsu KY, Noda H, Yamamoto Y, Kurozumi T. Is isoniazid hepatotoxicity induced by the metabolite hydrazine? *Journal of Uoeh* 1983; 5: 183-190.
32. Cotran RS, Kumar V, Robbins SL. Genetic Disorders. In: Cotran RS, Kumar V, Robbins SL. *Pathologic Basis of Disease*. Edn 5, W B Saunders Co, Philadelphia, 1994, 143.
33. Kaplowitz NTY, Simon FR, Stolz A. Drug induced hepatotoxicity. *Annals of Internal Medicine* 1986; 104: 826-839.
34. Shinozuka H, Farber JL, Konishi Y, Anukarhanonta T. D-galactosamine and acute liver cell injury. *Federation Proceeding* 1973; 32: 1516-1526.
35. Leist M, Gantner F, Bohlinger I, Germann PG, Wendel A. Murine hepatocyte apoptosis induced in vitro and in vivo by TNF- α requires transcription arrest. *Journal of Immunology* 1994; 153: 1778-1787.
36. Kasravi FB, Wang L, Wang D, Molin G, Bengmark S, Jeppsson B. Bacterial translocation in acute liver injury induced by D-galactosamine. *Hepatology* 1996; 23: 97-103.
37. Michael DJ, Bahjat FR, Fukuzuka K, Ksontini R, Solorzano CC, Edwards CK, Tannahill CL, Mackay SD, Copeland EM, Moldawer LL. Lipopolysaccharide and D-galactosamine-induced hepatic injury is mediated by TNF- α and not Fas ligand. *American Journal of Physiology* 2000; 278: R1196-R1201.
38. Komatsu Y, Shiratori Y, Kawase T, Hasimoto N, Han K, Shiina S, Matsumura M, Niwa Y, Kato N, Tada M, Ikeda Y, Tanaka M, Omata M. Role of polymorphonuclear leukocytes in galactosamine hepatitis: mechanism of adherence to hepatic endothelial cells. *Hepatology* 1994; 20: 1548-1556.
39. Klebanoff SJ, Vadas MA, Harlan JM, Sparks LH, Gamble JR, Agosti JM, Waltersdorff AM. Stimulation of neutrophils by tumor necrosis factor. *Journal of Immunology* 1987; 136: 4220-4225.
40. Komatsu Y, Shiratori Y, Hikiba Y, Hashimoto N, Han K, Kawase T, Yoshida H, Okano K, Omata M. Role of Platelet-activating factor in pathogenesis of galactosamine-lipopolysaccharide-induced liver injury. *Digestive Disease and Sciences* 1996; 41: 1030-1037.
41. Jarup L, Berglund M, Elinder CG, Nordberg G, Vahter M, Scand J. Health effects of cadmium exposure-a review of the literature and a risk estimate. *Journal of Work, Environment and Health* 1998; 24: 1-51.
42. Dudley RE, Svoboda DJ, Klaassen CD. Acute exposure to cadmium causes liver severe injury in rats. *Toxicology and Applied Pharmacology* 1982; 65: 302-313.
43. Sarkar S, Yadav P, Trivedi R, Bansal AK, Bhatnagar D. Cadmium induced lipid peroxidation and the status of antioxidant system in rat tissues. *Journal of Trace Elements in Medicine and Biology* 1995; 9: 144-149.
44. Stohs SJ, Bagchi D, Hassoun E, Bagchi M. Oxidative mechanisms in the toxicity of chromium and cadmium ions. *Journal of Environmental Pathology, Toxicology and Oncology* 2000; 19(3): 201-213.
45. Rikans LE, Yamano T. Mechanisms of cadmium-mediated acute hepatotoxicity-a review. *Journal of Biochemical and Molecular Toxicology* 2000; 14(2): 110-117.
46. Essigmann JM, Croy RG, Nadzan AM, Busby WF, Reinhold VN, Buchi G, Wogan GN. Structural identification of the major DNA adduct formed by aflatoxin B1 in vitro. *Proceedings of the National Academic Sciences* 1977; 74: 1870-1874.
47. Swenson DH, Miller EC, Miller JA. Aflatoxin B1-2, 3-oxide: evidence for its formation in rat liver in vivo and by human liver microsomes in vitro. *Biochemical and Biophysical Research Communication* 1974; 60: 1036-1043.
48. Gallagher EP, Wienkers LC, Stapleton PL, Kunze KL, Eaton DL. Role of human microsomal and human complementary DNA-expressed cytochromes P450 1A2 and P450 3A4 in the bioactivation of aflatoxin B1. *Cancer Research* 1994; 54: 101-108.
49. Wogan GN, Newberne PM. Dose-response characteristics of aflatoxin B1 carcinogenesis in the rat. *Cancer Research* 1967; 27: 2370-2376.
50. Neal GE, Judah DJ, Stirpe F, Patterson DSP. The formation of 2, 3-dihydroxy-2, 3-dihydroaflatoxin B1 by liver microsomes isolated from certain avian and mammalian species. *Toxicology and Applied Pharmacology* 1981; 58: 431-437.
51. Roebuck BD. Hyperplasia, partial hepatectomy, and the carcinogenicity of aflatoxin B1. *Journal of Cellular Biochemistry* 2004; 91: 243-249.
52. Serafini-Cessi F. Conversion of allyl alcohol into acrolein by rat liver. *Biochemical Journal* 1972; 128: 1103-1107.
53. Schauenstein E, Esterbauer H, Zollner H. Aldehydes in Biological Systems. In: Lagnado JR. *Aldehydes in Biological Systems, Their Natural Occurrence and Biological Activities*. Pion Limited, London, 1977, 1-7.
54. Esterbauer H, Zollner H, Scholz N. Reaction of glutathione with conjugated carbonyls. *Zeitschrift Fur Naturforschung* 1975; 30: 466-473.
55. Boyland E, Chasseaud LF. Enzyme-catalysed conjugation of glutathione with unsaturated compounds. *Biochemical Journal* 1967; 204: 95-102.
56. Jaeschke H, Kleinwaechter C, Wendel A. The role of acrolein in allyl alcohol-induced lipid peroxidation and liver cell damage in mice. *Biochemical Journal*. 1987; 335: 85-94.
57. McLain GE, Sipes IG, Brown BR. An animal model of halothane hepatotoxicity: roles of enzyme induction and hypoxia. *Anesthesiology*. 1979; 51: 321-323.
58. Spraklin DK, Hankins DC, Fisher JM, Thunmel KF, Kharash FD. Cytochrome P450 2E1 is the principal catalyst of human oxidative halothane metabolism in vitro. *Journal of Pharmacology and Experimental Therapeutics* 1997; 281: 400-411.
59. Trulsson E, Ullissey MJ. Halothane anesthesia alters cerebral enzymes: a histochemical study in the rat. *Acta Anatomica* 1997; 130: 163-167.
60. Friedman LS, Martin P, Munoz SJ. Liver Function Tests and the Objective Evaluation of the Patient with Liver disease. In: Zakin D, Boyer TD. *Hepatology: A Textbook of Liver Disease*. Edn 3, W B Saunders Co, Philadelphia, 1996, 791-833.
61. Kaplan MM. Laboratory tests. In: Schiff L, Schiff ER. *Diseases of the Liver*. Edn 7, J B Lippincott, Philadelphia, 1993, 108-144.
62. Clermont RJ, Chalmers TC. The transaminase tests in liver disease. *Medicine* 1967; 46: 197-207.
63. Matloff DS, Selinger MJ, Kaplan MM. Hepatic transaminase activity in alcoholic liver disease. *Gastroenterology* 1980; 78: 1389-1392.
64. Kaplan MM. Serum alkaline phosphatase-another piece is added to the puzzle. *Hepatology* 1986; 6: 526-528.
65. Dickson ER, Grambsch PM, Fleming TR, Fisher LD, Langworthy A. Prognosis in primary biliary cirrhosis: model for decision making. *Hepatology* 1989; 10: 1-7.

66. Skrede S, Solberg HE, Blomhoff JP, Gjone E. Bile acids measured in serum during fasting as a test for liver disease. *Clinical Chemistry* 1978; 24: 1095-1099.
67. Keefe EB, Sutherland MC, Gabourel JD. Serum gamma-glutamyl transpeptidase activity in patients receiving chronic phenytoin therapy. *Digestive Diseases and Sciences* 1986; 31: 1056-1061.
68. Hill PG, Sammons HG. An assessment of 59-nucleotidase as a liver function test. *The Quarterly of Journal of Medicine* 1967; 36: 457-468.
69. Richer G, Viallet A. Mitochondrial antibodies in extrahepatic biliary obstruction. *Digestive Diseases and Sciences* 1974; 19(8): 740-744.
70. Cotran RS, Collins T. The Liver and Biliary Tract. In: Robbins Pathologic Basis of Disease. Edn 6, W B Saunders Co, Philadelphia, 1999, 846-891.
71. Dial SM. Clinicopathologic evaluation of the liver. *Veterinary Clinics of North America: Small Animal Practice* 1995; 25(2): 257-273.
72. Hall RL. Laboratory evaluation of liver disease. *Veterinary Clinics of North America: Small Animal Practice* 1985; 1: 3-19.
73. Lehrer JK. Medical Encyclopedia: Albumin-Serum. Available from: www.nlm.nih.gov/medlineplus/ency/article/003480.htm. [Accessed on 29 July 2009].
74. Tilley L, Smith F. The 5-Minute Veterinary Consult: Canine and Feline. Edn 2, Lippincott Williams & Wilkins, Baltimore, 2000, 710-711.
75. Ita SO, Akpanyung EO, Umoh BI, Ben EE, Ulkafia. Acetaminophen induced hepatic toxicity: protective role of ageratum conyzoides. *Pakistan Journal of Nutrition* 2009; 8(7): 928-932.
76. Muruges KS, Yeligar VC, Maiti BC, Maity TK. Hepatoprotective and antioxidant role of *Berberis tinctoria* Lesch leaves on paracetamol induced hepatic damage in rats. *Iranian Journal of Pharmacology and Therapeutics* 2005; 4(1): 64-69.
77. Juraj G, Chovanova Z, Marekova M, Chavkova Z, Tomeckova V, Mojzisoava G, Kusnir J. Effect of quercetin on paracetamol-induced rat liver mitochondria dysfunction. *Biologica Bratislava* 2004; 59(3): 399-403.
78. Ghosh T, Maity TK, Das M, Bose A, Dash GK. Hepatoprotective activity of *Bacopa monnieri* L. against ethanol-induced hepatotoxicity in rats. *Pharmacognosy Magazine* 2007; 3(10): 95-100.
79. Mahendran P, Devi CSS. The modulating effect of *Garcinia Cambogia* extract on ethanol induced peroxidative damage in rats. *Indian Journal of Pharmacology* 2001; 33: 87-91.
80. Haldar PK, Gupta M, Mazumder UK, Kandar CC, Manikandan L. Hepatoprotective effect of *Wedelia Calendulaceae* against thioacetamide induced liver damage in rats. *Pharmacologyonline* 2007; 3: 414-421.
81. Huang HC, Wang SS, Chan CY, Chen YC, Lee FY, Chang FY, Chu CJ, Lin HC, Lu RH, Lee SD. Role of hepatic nitric oxide synthases in rats with thioacetamide-induced acute liver failure and encephalopathy. *Journal of the Chinese Medical Association* 2007; 70(1): 16-23.
82. Kim NH, Hyun SH, Jin CH, Lee SK, Lee DW, Jeon TW, Lee JS, Chun YJ, Lee ES, Jeong TC. Pretreatment with 1, 8-cineole potentiates thioacetamide-induced hepatotoxicity and immunosuppression. *Archives of Pharmacol Research* 2004; 27(7): 781-789.
83. Whitehouse LW, Tryphonas L, Paul CJ, Solomonraj G, Thomas BH, Wong LT. Isoniazid- induced hepatitis steatosis in rabbits: an explanation for susceptibility and its antagonism by pyridoxine hydrochloride. *Canadian Journal of Physiology and Pharmacology* 1983; 61: 478-487.
84. Karthikeyan S, Krishnamoorthy MS. Effect of subacute administration of isoniazid and pyridoxine on lipids in plasma, liver and adipose tissues in the rabbit. *Drug and Chemical Toxicology* 1991; 14: 293-303.
85. Pandit S, Sur TK, Jana U, Debnath PK, Sen S, Bhattacharya D. Prevention of carbon tetrachloride induced hepatotoxicity in rats by *Adhatoda vasika* leaves. *Indian Journal of Pharmacology* 2004; 36(5): 312-320.
86. Junnila M, Rakho A, Sukura, Lindberg LA. Reduction of carbon tetrachloride induced hepatotoxic effects by oral administration of betaine in male hans-wistar rats: a morphometric histological study. *Veterinary Pathology* 2000; 37: 231-238.
87. Lee HS, Keum KY, Ku SK. Effects of *Picrorrhiza* rhizome water extracts on the subacute liver damages induced by carbon tetrachloride. *Journal of Medicinal Food* 2007; 10(1): 110-117.
88. Imaoka M, Kato M, Tamanaka M, Hattori H, Manabe S. Aggravation of galactosamine hepatotoxicity by albumin in rats. *Journal of Toxicologic Pathology* 2008; 21: 175-183.
89. Zhou Y, Park CM, Cho CW, Song YS. Protective effect of pinitol against D-galactosamine-induced hepatotoxicity in rats fed on a high-fat diet. *Bioscience, Biotechnology, and Biochemistry* 2008; 72(7): 1657-1666.
90. Nair SP. Protective effect of tefroli-a polyherbal mixture (tonic) on cadmium chloride induced hepatotoxic rats. *Pharmacognosy Magazine* 2006; 2(6): 112-118.
91. Jeyaprakash K, Chinnaswamy P. Effect of spirulina and Liv.52 on cadmium induced toxicity in albino rats. *Indian Journal of Experimental Biology* 2005; 43: 773-781.
92. Amin A, Hamza AA, Daoud S, Hamza W. Spirulina protects against cadmium-induced hepatotoxicity in rats. *American Journal of Pharmacology and Toxicology* 2006; 1(2): 21-25.
93. Lee JR, Park SJ, Lee HS, Jee SY, Seo J, Kwon YK, Kwon TK, Kim SC. Hepatoprotective activity of licorice water extract against cadmium-induced toxicity in rats. *Evidence-Based Complementary and Alternative Medicine* 2009; 6(2): 195-201.
94. Luyendyk JP, Copple BL, Barton CC, Patricia E, Ganey PE, Roth RA. Augmentation of aflatoxin B1 hepatotoxicity by endotoxin: involvement of endothelial and the coagulation system. *Journal of Toxicological Sciences* 2003; 72: 171-181.
95. Rastogi R, Srivastava AK, Rastogi AK. Long term effect of aflatoxin B (1) on lipid peroxidation infracts liver and kidney: effect of picroliv and silimarin. *Phytotherapy Research* 2001; 6(3): 319-321.
96. Dhalwal K, Shinde V, Mahadik KR, Kadam SS. Hepatoprotective activity of *Sida rhombifolia ssp. retusa* against thioacetamide and allyl alcohol intoxication in rats. *Pharmacologyonline* 2006; 3: 259-266.
97. Alam K, Nagi MN, Al-Shabanah OA, Al-Bekairi AM. Beneficial effect of nitric oxide synthase inhibitor on hepatotoxicity induced by allyl alcohol. *Journal of Biochemistry and Molecular Toxicology* 2001; 15(6): 317-321.
98. Marit GB, Dodd DE, George ME, Vinegar A. Hepatotoxicity in guinea pigs following acute inhalation exposure to 1,1-dichloro-2,2,2-trifluoroethane. *Toxicologic Pathology* 1994; 22: 404-416.
99. Lind RC, Gandolfi AJ, Burnell R, Brown BR, Pauline de la M, Hall PDL. Halothane hepatotoxicity in guinea pigs. *Anesthesia and Analgesia* 1987; 66: 222-228.
100. Reitman S, Frankel A. Colorimetric method for determination of serum glutamate oxaloacetate and glutamic pyruvate transaminase. *American Journal of Clinical Pathology* 1957; 28: 56-58.
101. Kind PRN, King EJ, Varley H, Gowenlock AH, Bell M. Method of Practical Clinical Biochemistry. Heinman, London, 1980, 899-900.
102. Malloy HT, Evelyn KA. The determination of bilirubin with photometric colorimeter. *Journal of Biological Chemistry* 1937; 119: 481-490.
103. Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein dye binding. *Journal of Biological Chemistry* 1976; 72: 248-254.
104. Harper AE. Methods of Enzymatic Analysis. Academic Press, New York, 1965, 788-790.
105. Varley H. Practical Clinical Biochemistry. Edn 4, C B S Publishers and Distributors, New Delhi, 2003, 1-80.
106. Ohkawa H, Ohishi N, Yagi K. Assay for lipid peroxidation in animal tissues by thiobarbituric acid reaction. *Analytical Biochemistry* 1979; 95: 351-358.
107. Ellman GL. Tissue sulphhydryl groups. *Archives of Biochemistry and Biophysics* 1959; 82: 70-77.
108. Singh B, Saxena AK, Chandan BK, Agrawal SG, Anand KK. In vivo hepatoprotective activity of active fraction from ethanolic extracts of *Eclipta alba* leaves. *Indian Journal of Physiology and Pharmacology* 2001; 45(4): 435-441.
109. Turner RA. Screening Methods in Pharmacology, Edn 1, Academic Press, India, 2009, 299-300.
110. Rajan R, Subrahmanyam K. Uptake of sodium phenol tetrabromophthalein (bromsulphalein) by rat liver slices under different conditions. *Indian Journal of Experimental Biology* 1965; 24: 100-103.
111. William GM, Elizebeth K, Weisburger HJ. Isolation of long term cell culture of epithelial like cells from rat liver. *Experimental Cell Research* 1971; 69: 106-112.