

Opportunities for Oral Healthcare Professionals in Tobacco Cessation in Adolescents

Dr. Shilpa Khullar

Professor, Dept. of Prosthodontics,
Inderprastha Dental College &
Hospital, Sahibabad, Ghaziabad, UP.

Dr. S. Sood

Post-graduate Student,
Dept. of Public Health Dentistry
ITS-CDSR, Muradnagar,
Ghaziabad UP.

Dr. A. Mittal

Professor & H.O.D.
Dept. of Conservative
Dentistry

Dr. K. Bhargava

Professor & H.O.D.
Dept. of Oral Pathology
& Microbiology

Dr. Pankaj Datta

Principal & H.O.D.
Dept. of Prosthodontics

Dr. S. Sharma

Dept. of Pedodontics,

Inderprastha Dental College & Hospital, Sahibabad, Ghaziabad UP.

Abstract

Tobacco use is one of the leading preventable causes of morbidity and mortality. The most powerful predictor of adult tobacco use is its initiation during adolescence, the most susceptible time for onset of this habit. Initiation of tobacco use is associated with peer pressure, parental use, school factors, cultural norms, lower self esteem, accessibility, moderate pricing, desire for experimentation and aggressive marketing by tobacco companies.

While dentists have a positive attitude regarding their role in tobacco cessation, the same is not extrapolated into practice. Several barriers to counseling in the dental clinic have been identified. Dental professionals can render tobacco cessation services to the youth. Brief interventions, self-help materials, and nicotine replacement therapy for established nicotine dependence form the mainstay of therapy. The purpose of this paper is to identify the several factors leading to tobacco initiation in adolescent and discuss the role of dentist in tobacco cessation programs.

Key Words: Adolescents, tobacco, initiation, cessation, oral health, dentist

Introduction

Tobacco is considered to be the single most important cause of preventable deaths in the world. Preventing and treating diseases caused by tobacco usage is one of the major challenges of public health today. Despite the current knowledge of the harm caused by tobacco, its consumption continues to increase. The tobacco epidemic is shifting from industrialized to developing countries, due to steady population growth coupled with tobacco industry ensuring that millions of people become fatally addicted each year. Nearly 5 million people die due to tobacco use every year and this figure will increase to 10 million tobacco attributable deaths per year by 2020. Of these, 7 million deaths will occur in the developing countries, mainly China and India.¹

Tobacco And Youth

There is a need to identify high-risk segments of the population such as youth as an entry point to tobacco use as targeted by industry promotion. Most tobacco users start using it before the age of 18 years, while some start as young as 10 years², the time for discovery, challenge and experimentation, when they are far too young to understand or resist social expectations. The early age of initiation underscores the urgent need to intervene and protect this vulnerable group from falling prey to this addiction. The risks of tobacco use are highest among those who start early and continue its use for a long period. In order to reduce the long-term burden of tobacco related diseases, adoption of successful prevention and cessation

strategies is the only feasible solution in less-resourceful countries.

It is important for Oral health professionals to understand the factors which lead to tobacco abuse in youth, to counter this public health threat by their intervention.

Determinants of Tobacco Use in Youth

Tobacco use in adults is primarily due to nicotine dependence whereas in adolescents there are unique social and behavioral factors associated with tobacco use.³

Socio-environmental factors such as advertisements by tobacco companies and effect of role models in movies who are smokers and portrayed as smart, successful and courageous considerably increases the chances of smoking initiation among teens.⁴ Enhancement of athletic ability, concentration powers and being 'cool' have been wrongly associated with tobacco use. Moderate pricing and easy availability of tobacco products to even the small kids despite the ban by the governments has led to sharp rise in its use. Children may also start using tobacco for psycho-social reasons like peer influences, curiosity, desire for experimentation or as a remedy for stress.⁵

Personal characteristics of adolescent tobacco users include low self-esteem, low aspirations, depression/anxiety and risk taking. These personal traits are associated with certain behavior patterns such as poor school performance, violence, gang membership, and alcohol and drug abuse. Nicotine gives these users excitement and relief from the all pervasive gloom of life.⁵

Development of Dependence

For the past two decades, the onset of dependence has been conceptualized by the Stage theory as a slow and sequential process, with the daily use of tobacco over an extended period of time as a prerequisite for nicotine dependence.⁶ Contradicting this theory, preliminary results from the DANDY (Development and Assessment of Nicotine Dependence in Youth) study suggested that the first symptoms of nicotine dependence can appear within a matter of days or weeks of the onset of intermittent tobacco use.^{6,7}

Once adolescents have experimented with tobacco, approximately 50% continue its use and become addicted. Preventing this use requires intervention in the early adolescence prior to the time when these behaviors have already become ingrained.⁸

III. Effects of Tobacco on General Health

Tobacco use is a common risk factor to several systemic and oral diseases. Second hand smoke from parents puts the children at an increased risk of developing lung cancer, respiratory and cardiovascular diseases, middle-ear infections and delayed development of permanent teeth by as many as 4 months.⁹

Smoking during pregnancy increases the

chance of a low-birth-weight baby, premature birth, stillbirth, sudden infant death syndrome and six times greater chance of cleft palate formation.¹⁰

Although the most serious health outcomes associated with smoking typically emerge later in life, adolescent smokers show evidence of airway obstruction, slowed growth in lung function and other respiratory symptoms, compared with non-smokers. In addition, the earlier individuals begin to smoke, the higher their risk for heart disease, stroke, and chronic obstructive lung disease, risk of developing anxiety disorders/depression and nicotine addiction.¹¹

III Effects of Tobacco on Oral Health

Tobacco-induced oral diseases contribute significantly to the global oral disease burden^{12,13}. A clear association between tobacco use and the prevalence and severity of periodontal diseases exists.¹⁴ Tobacco use is responsible for up to half of all periodontitis cases among adults¹³. Moreover, tobacco use impedes the effectiveness of periodontal therapy and wound healing.¹⁵ Smoking has been shown to affect both taste and smell acuity. Tobacco, whether chewed or smoked, causes halitosis and staining of teeth¹⁶.

Smoking a pack of cigarettes a day or using smokeless tobacco quadruples the risk of developing oral cavity or oro-pharyngeal cancer, which kills about 50% of its victims within 5 years of diagnosis. Vast majority of oral cancers are preceded by precancerous lesions and conditions caused by the use of tobacco in some form.¹⁷ Such lesions can be easily seen due to their peculiar oral location, making oral cancers particularly amenable to prevention.

The clear link between oral diseases and tobacco use provides an ideal opportunity for oral health professionals to take part in tobacco control and cessation programs.

Role of Dentist in Tobacco Control Measures

Dentists can play a major role to control tobacco menace by Public Health Education at community level and counseling and use of Nicotine Replacement Therapy for tobacco cessation in the clinics.

Health Education and Information

Prevention against the diseases that come with tobacco usage is based primarily on public and individual education to drop the habit or preferably not to begin in the first place.

Pediatric dentist can take initiative to conduct school-based tobacco prevention programs to educate the adolescents about the health risk of tobacco consumption, risk of addiction and benefits of tobacco cessation. Such programs identify the social influences which promote tobacco use among the youth and teach skills to resist such influences which can produce a long term relative

improvement of 25-30% in quit rates,¹⁸ or delay adolescent tobacco use, especially if strengthened by booster session and community programs involving parents and community participation.

Public health Dentist can write articles about benefits of tobacco control, participate in talk shows, demonstrations, discussions and link with NGO's to involve youth in anti-tobacco advocacy. The focus should not be only on primary prevention, that is not only on discouraging young people from taking up the habit but also on providing help and support for those who wish to quit tobacco usage.

Tobacco Cessation for Youth in Dental Clinics

The scope of preventive dentistry is constantly expanding and can be as far reaching as a professional's imagination, sense of responsibility and efforts. Dentists have been recognized as "ideally positioned to counsel against the use of tobacco products." They can relay specific information concerning the oral ill effects of tobacco use. The dental encounter probably constitutes a "teachable moment" when the patient is receptive to counseling about life-style issues. Oral health professionals should integrate tobacco use, prevention and cessation services into their routine and daily practice¹⁹ for the following reasons.

1. They are especially concerned about the adverse effects of tobacco practices in the oro-pharyngeal region of the body.

2. They have easy access to children, youths and their caregivers, thus providing opportunities to influence individuals to avoid all together, postpone initiation or quit using tobacco before they become dependent.

3. They often have more time with patients than many other clinicians, providing opportunities to integrate education and intervention methods into practice.

4. They often treat women of childbearing age, thus are able to inform such patients about the potential harm to their babies from tobacco use.

5. They can build their patient's interest in discontinuing tobacco use by showing actual tobacco effects in the mouth.

Dentists in many parts of the world have a positive attitude about intervening for their adolescent patients. The majority of dentists consider smoking cessation and prevention for adolescents and children as part of their responsibility²⁰. Pediatric dentists should encourage, advice and assist tobacco users to quit²¹.

Most dentists and hygienists don't counsel children to prevent tobacco use due to lack of confidence and doubt about the effectiveness of their intervention efforts.^{20,22}

Barriers Mitigating Provision of Tobacco Cessation Counseling

Numerous barriers have been identified for the limited involvement of dental professionals in tobacco cessation programs for youth.^{20,21,22,23,24} Some of the consistently reported barriers are lack of time, lack of reimbursement or incentives, resistance from the patients or parents, lack of skills or formal training translating into lack of confidence, lack of readily accessible patient education materials, perception of poor effectiveness and fear that giving unwanted tobacco cessation counseling may upset the dentist-patient relationship. Children and adolescents also under-report tobacco use on health history forms that ask them to specify whether they use tobacco products²⁵ making it difficult to identify them in the first place.

Globally about 70% of 13-15 year olds who currently smoke have a desire to quit. Though a high number of tobacco users want to quit, few make a dedicated quit attempt (around 40%) and a negligible proportion of these (around 3%) are successful in achieving long term abstinence¹⁸. Parents are more likely to approve than disapprove of the dentist counseling the child and the parent.²⁶

The lack of training at the graduate level is also an important barrier that hinders large-scale involvement of dentists. The professional skills required by the dentists to provide tobacco cessation counseling to their patients ideally should be learnt during the dental curriculum and reinforced within continuing education. Dental colleges need to incorporate into their curricula not just didactic instruction on the oral health impact of tobacco use, but relevant counseling techniques and training in nicotine replacement therapy.

ideally should be learnt during the dental curriculum and reinforced within continuing education. Dental colleges need to incorporate into their curricula not just didactic instruction on the oral health impact of tobacco use, but relevant counseling techniques and training in nicotine replacement therapy.

Brief Interventions

Dentists can help youth by providing advice on when to quit, help the patient to identify problems and strategies to deal with problems.

Behavioral interventions for tobacco use

conducted by oral health professionals in the dental clinic and community setting may increase tobacco abstinence rates among smokeless tobacco users²⁷ Dental treatment often necessitates frequent contact with patients over an extended period of time, providing a mechanism for long-term contact and reinforcement, coupled with visible changes in the oral cavity in response to counseling²⁸.

Brief interventions typically involve an assessment of tobacco use, dependence, and motivation to quit; advice on the benefits and methods of quitting; and assistance with quitting, including referrals to other treatment modalities.

Motivating Factors for Tobacco Cessation


1. It is important to note that adolescents consistently rank physical attractiveness, dental concerns, and oral health as greatly important.²⁹

2. Relating tobacco to short-term adverse effects such as staining of teeth, halitosis, loss of taste may be more relevant and meaningful to an adolescent than long-term health effects such as cardiovascular or lung diseases³⁰

3. Highlighting role models abstaining from tobacco use and making the dental clinic adopt a "no tobacco policy" can also be used.

However, it is important to realize that tobacco cessation is a process and a number of stages are encountered in the process. Some of the attributes required for the clinician are to be persistent, supportive and not to give up.³¹

The 5 A's (ask, advice, access, assist and arrange) is a brief intervention method, used to guide the dentist in tobacco cessation counseling. It is important to include some sort of intervention to bring behavior change, in cases where the adolescent wishes to quit tobacco.³²



Ask every adolescent a simple question about current tobacco use and use record system in the clinic to document his/ her tobacco use status during every visit. Once a tobacco user is identified, assess willingness to make a quit attempt, the dentist should urge him or her, in a “clear, strong and personalized manner,” to quit. The dentist can assist tobacco users by helping them set a quit date; referring them to a telephone counseling service, cessation group or intensive cessation program; prescribing pharmacotherapy; and providing educational materials about tobacco cessation. Follow-up contact to support and guide a patient's quit attempt should be arranged otherwise, users may slip back to earlier stages of change.

Users unwilling to make a quit may respond to a motivational intervention that provides the clinician an opportunity to educate, reassure, and motivate. Motivational intervention is built around the '5 R's': Relevance, Risks, Rewards, Roadblocks and Repetition. Such counseling involves talking about tobacco and quitting and then reinforcing the points most likely to motivate adolescents to quit. Information should be of their relevance, such as health concerns, rewards or specific barriers to quitting. A discussion of the health effects of tobacco and the benefits of quitting (such as immediate improved oral health and financial savings) may allow the dentist to identify and highlight risks and rewards that seem most relevant to the user. Identifying the patient's perceptions of roadblocks to quitting, such as fear of withdrawal symptoms or weight gain and address those barriers. The motivational intervention should be repeated during every clinic visit by a user, who is unwilling to quit.

Nicotine replacement therapy (NRT)

The pharmacological means include NRT and antidepressants like bupropion. While intensive therapy is not in the realm of dentists providing brief interventions but NRT holds plenty of promise. Even though these methods may seem exclusive of each other, the existing data suggest that a combination of the two is often essential to achieve good success rates.³³ Normally these services will lie outside the dental practice although some trained dental teams will be able to provide these services.

Adolescent tobacco users are different from older tobacco users in that their motivation to stop smoking tends to be more unstable. It is sensible, therefore, to check that they are fully committed to trying to stop smoking permanently before offering them NRT and to attempt to establish that they are dependent³⁴

Self-help

The self-help, non-interactive approach includes minimal interventions that do not require responses from the adolescent and are delivered through written or audio-visual materials or on a computer; to motivate the patient to quit the habit.

Challenges to Youth Tobacco Cessation Programs

1. Tobacco cessation programs especially for youth are a resource-intensive intervention.
2. Negligible cessation clinics are available for adolescents or implemented in the

developing countries.

3. Cessation programs implemented in isolation with lack of supportive environment and government policies for tobacco control.
4. Low importance placed by oral health professional on tobacco cessation.

Conclusion

Though we do not fully understand all the factors that contribute to onset of tobacco use, which leads to addiction and eventual adverse health outcomes; we do need to understand better the patterns of use and how the determinants of initiation interact.

As oral health care providers, dentists must take the responsibility of providing tobacco cessation services and encourage non-users to be tobacco free, though few translate this into practice. Admittedly, there are several barriers in this process, both real and perceived, which should be addressed with further research. Screening for tobacco use, interventions, referring adolescents to additional resources for cessation, and establishing a follow-up system that will track each adolescent's progress should be made mandatory. Brief advice from a dentist for adolescents is cost-effective and has a potentially large reach.

Success in relation to cessation does not only mean that more number of patients has quit, but it also includes educating the masses, so that the number of individuals taking to this habit afresh will also reduce. Today, we the members of the oral health profession along with policy makers should help in achievement of a tobacco-free society so that we can protect the health of the coming generations.

References

1. WHO: The World Health Report: Reducing Risks, Promoting Healthy Life. Geneva: WHO, 2002.
2. Tobacco use among youth: a cross country comparison: The GYTS collaborative group. Tobacco control. September 2002;11(3):252-272
3. Sanchez del Mazo -- May 2005. Youth Smoking Cessation - What can we do? Available from: <http://www.fhi.se/Documents/Vart-uppdag/tobak/youth-smoking-cessation-0505.pdf>
4. Diane Logan, Beatriz Carlini-Marlatt. Smoking and Adolescence: some issues on prevention and cessation. available at http://www.mentorfoundation.org/pdfs/prevention_perspectives/5.pdf (accessed 12.11.10)
5. Chadda RK, Sengupta SN. Tobacco use by Indian adolescents. Tobacco Induced Diseases 2002;1(2): 11119.
6. DiFranza JR, J A Savageau, et al. Development of symptoms of tobacco dependence in youths: 30 month follow up data from the DANDY study. Tobacco Control 2002;11:228235.
7. DiFranza JR, Rigotti NA, McNeill AD, et al. Initial symptoms of nicotine dependence in adolescents. Tobacco Control 2000;9:31319.
8. Sinha DN, Reddy KS, et al. Linking GYTS data to the WHO FCTC: The case for India. Indian J of Public Health 2006; 50: 76-89.
9. International Consultation on Environmental Tobacco Smoke (ETS) and Child Health. Consultation Report, WHO, 1999. Available from <http://old.ash.org.uk/html/passive/html/who-ets.html>
10. Tobacco use and reproductive outcomes. In: Reddy KS, Gupta PC (eds). Report on tobacco control in India. New Delhi: Ministry of Health and Family Welfare, Government of India; 2004:108-10.
11. Pbert L, Moolchan ET, Muramoto M, Winickoff JP, Curry S, Lando H, et al. The State of Office-Based Interventions for Youth Tobacco Use. Pediatrics 2003;111:e650-60.
12. Reibel J. Tobacco and oral diseases: an update on the evidence, with recommendations. Med Princ Pract 2003; 12 (suppl. 1): 22-32.
13. Tomar SL, Asma S. Smoking attributable periodontitis in the United States: findings from

- NHANES III. J Periodontol 2000; 71: 743-51.
14. Bergstrom, J., Eliasson, S., Dock, J. 10-year prospective study of tobacco smoking and periodontal health. Journal of Periodontology 2000; 71, 1338-47.
15. Davis JM. Tobacco Cessation for the Dental Team: A Practical Guide Part: I background and overview. J Contemp Dent Pract 2005;3:158-66.
16. Allard, R., Johnson, N., Sardella A et al. Tobacco and Oral Diseases: Report of EU Working Group. Journal of Irish Dental Association 1999; 46, 12-23.
17. WHO: Tobacco-related Oral Mucosal Lesions and Dental Diseases 4.7. Available from: <http://www.whoindia.org/SCN/Tobacco/Report/03-Chapter-04.7.pdf>
18. Global youth meet on tobacco control (GYM 2009). Youth advocacy for global tobacco control: insight, issues and incentives. HRIDAY
19. Petersen PE. The World Oral Health Report 2003. Continuous improvement of oral health in the 21st century and the approach of the World Health Organization Global Oral Health Programme. Geneva, World Health Organization, 2003
20. Wyne AH, Chohan AN, Al-Moneef MM, Al-Saad AS. Attitudes of General Dentists about Smoking Cessation and Prevention in Child and Adolescent Patients in Riyadh, Saudi Arabia. J Contemp Dent Pract 2006;1:35-43.
21. Gansky SA, Ryan JL, Ellison JA, Isong U, Miller AJ, Walsh MM. Patterns and correlates of tobacco control behavior among American Association of Pediatric Dentistry members: a cross-sectional national study. BMC Oral Health 2007;7:13.
22. Kast KR, Berg R, Deas A, Lezotte D, Crane LA. Colorado Dental Practitioners' Attitudes and Practices Regarding Tobacco-Use Prevention Activities for 8- Through 12-Year-Old Patients. J Am Dent Assoc 2008;139:467-75.
23. Goodman H S., Vargas C M. et al. Maryland. General and pediatric dentists' attitudes regarding tobacco use prevention and cessation in their child and adolescent patients. Public health and the environment. APHA 2004. p. 6-10.
24. Beaglehole, R.H. The role of oral health professionals in tobacco control in OECD countries: policies and initiatives. Master's Thesis. University College London, 2003.
25. Hennrikus D, Rindal DB, Boyle RG, Stafne E, Lazovich D, Lando H. How well does the health history form identify adolescent smokers? J Am Dent Assoc 2005;136:1113-20.
26. Vergotine R J, Koerber A, Shires D. Pediatric Dentist's role in Tobacco Counseling for Children: Parental Attitudes. IADR 86th General session and Exhibition.
27. Carr A, Ebbert J. Interventions for tobacco cessation in the dental setting. Cochrane Database of Systematic Reviews 2006, Issue 1. Art. No.: CD005084. DOI: 10.1002/14651858. CD005084. pub2.
28. Albert DA, Severson HH, Andrews JA. Tobacco use by adolescents: the role of the oral health professional in evidence-based cessation programs. Pediatr Dent 2006;28:177-87.
29. Johnson JG, Cohen P, Pine DS, Klein DF, Kasen S, Brook JS. Association between cigarette smoking and anxiety disorders during adolescence and early adulthood. JAMA 2000;284:2348-51.
30. Gansky SA, Ryan JL, Ellison JA, Isong U, Miller AJ, Walsh MM. Patterns and correlates of tobacco control behavior among American Association of Pediatric Dentistry members: a cross-sectional national study. BMC Oral Health 2007;7:13.
31. Albert DA, Severson HH, Andrews JA. Tobacco use by adolescents: the role of the oral health professional in evidence-based cessation programs. Pediatr Dent 2006;28:177-87.
32. Fiore MC, Bailey WC, Cohen SJ, Dorfman SF, Goldstein MG, Gritz ER, et al. Treating tobacco use and dependence. Quick reference guide for clinicians. Rockville, Maryland: US Department for Health and Human Services, Public Health Service, October 2000.
33. Mallin R. Smoking Cessation: Integration of Behavioral and Drug Therapies. Am Fam Physician 2002;65:1107-14.
34. Nicotine replacement therapy: guidance for health professionals on changes in the licensing arrangements for nicotine replacement therapy Ash, London, December 2005. Available from: www.nrt_guidance_uk_1205.