## **The Editorial**

## **Stem Cells of Epidermis: A Critical Introduction**

cells are cells Stem the having some distinguishing characteristics like longevity, high capacity of self-renewal and differentiation, quiescence and highly error-free proliferation. Almost all stem cells have the potential of lineage reprogramming, i.e. inter-conversion of cell lineages. They also have the potential to differentiate into almost all kinds of cells. These cells have been found in almost every organs of human body. Pool of stem cells found in epidermis is termed as Epidermal Stem Cells (Blanpain and Fuchs, 2006).

Many researchers around the world have reported different kinds of stem cells in skin, based on their cell surface makers, while they have not categorized these cells chronically (De Rosa and De Luca, 2012). It is worth mentioning here that all kinds of stem cells reported in skin, i.e. keratinocyte stem cells, limbal stem cells, hair follicle and bulge stem cells, SG (sebaceous gland) stem cells, and spinous keratinocytes express specific types of cytokeratin protein (e.g. K1, K3, K5, K10, K12, K14, K15, K19 etc.) on their surfaces (Bose et al., 2013; Forni et al., 2012; Ghadially, 2012).

Biologists have defined that almost all of these stem cells share single origin, i.e. Basal Layer of Embryonic Skin. As the embryonic skin passes the developmental stages, the basal layer produces two mother stem cells of skin, keratinocyte stem cells, and so called Limbal Stem Cells (in cornea) (Chee et al., 2006; Lavker and Sun, 2000).

Keratinocyte stem cells give rise to the cells expressing Cytokeratin proteins on their surfaces. So, all the cells expressing cytokeratin are tracked back to these stem cells in origin , while undergone natural lineage reprogramming or differentiation (Potten and Booth, 2002). The stem cells in the basal-layer give rise to keratinocyte stem cells which can be found in the basal layer of the adult skin (Kaur et al., 2004). During the developmental stages, this basal layer, containing keratinocyte stem cells, gives rise to limbal invagination of

\*Corresponding author E-mail: Muhammad Irfan-Maqsood Managing Editor, JCMR <u>muhammadirfanmaqsood@gmail.com</u> corneal region, a linage conversion mechanism happens, and the keratinocyte stem cells are naturally reprogramed into the limbal stem cells as shown in figure 1 (Dua and Azuara-Blanco, 2000; Pellegrini et al., 2001).



Figure 1: Proposed hierarchy of epidermal stem cells

In basal layer of the epidermis, the keratinocyte stem cells give rise to bulge, hair follicle, and SG stem cells when placode formation takes place.

In future, we need studies to find out which kind of cytokeratin protein is expressed early in these cells. In another word it would be helpful to define the order of cytokeratin expression regarding these lineage developmental processes from embryonic to mature skin and from embryonic basal layer to the formation of cornified epithelial cells.

## **References:**

1- Blanpain C. and Fuchs E. (2006) Epidermal stem cells of the skin. Annual review of cell and developmental biology 22:339-373.

2- Bose A., Teh M. T., Mackenzie I. C. and Waseem A. (2013) Keratin k15 as a biomarker of epidermal stem cells. International journal of molecular sciences 14:19385-19398.

3- Chee K. Y., Kicic A. and Wiffen S. J. (2006) Limbal stem cells: the search for a marker. Clinical & experimental ophthalmology 34:64-73.

4- De Rosa L. and De Luca M. (2012) Cell biology: Dormant and restless skin stem cells. Nature 489:215-217.

5- Dua H. S. and Azuara-Blanco A. (2000) Limbal stem cells of the corneal epithelium. Survey of ophthalmology 44:415-425.

6- Forni M. F., Trombetta-Lima M. and Sogayar M. C. (2012) Stem cells in embryonic skin development. Biological research 45:215-222.

7- Ghadially R. (2012) 25 years of epidermal stem cell research. The Journal of investigative dermatology 132:797-810.

8- Kaur P., Li A., Redvers R. and Bertoncello I. (2004) Keratinocyte stem cell assays: an evolving science. The journal of investigative dermatology. Symposium proceedings / the Society for Investigative Dermatology, Inc. [and] European Society for Dermatological Research 9:238-247. 9- Lavker R. M. and Sun T. T. (2000) Epidermal stem cells: properties, markers, and location. Proc Natl Acad Sci U S A 97:13473-13475.

10- Pellegrini G., Dellambra E., Golisano O., Martinelli E., Fantozzi I., Bondanza S., Ponzin D., McKeon F. and De Luca M. (2001) p63 identifies keratinocyte stem cells. Proc Natl Acad Sci U S A 98:3156-3161.

11- Potten C. S. and Booth C. (2002) Keratinocyte stem cells: a commentary. The Journal of investigative dermatology 119:888-899.