

Received: 27 December 2013 • Accepted: 10 February 2014

Research

Prevalence of *Enterobius Vermicularis* (pinworm) in Kermanshah city nurseries, using Graham: 2014

Zahra Sha-Mohammadi¹, Fariba Ghahramani², Mohammad Mahboubi^{3*}, Farzad Jalilian⁴, Mina Neiakane-Shahri⁵, Mohsen Mohammadi⁶

¹ Student Research Committee, Kermanshah University of Medical Sciences, Kermanshah, Iran

² MSc in Epidemiology, Shiraz University of Medical Sciences, Shiraz, Iran

³ PhD in Health Services Administration, Kermanshah University of Medical Sciences, Kermanshah, Iran

⁴ Ph.D Student of Health Education and Promotion; Department of Public Health, School Public Health, Kermanshah University of Medical Sciences, Kermanshah, Iran

⁵ Bs. Kermanshah University of Medical Sciences, Kermanshah, Iran

⁶ MSc Student in Health Services Administration, Kermanshah University of Medical Sciences, Kermanshah, Iran

*correspondence should be addressed to Mohammad Mahboubi, PhD in Health Services Administration, Kermanshah University of Medical Sciences, Kermanshah, Iran; Tell: +98; Fax: +98; Email: mmahboubi@kums.ac.ir.

ABSTRACT

Oxyuriasis is the most prevalent parasitic diseases in children and over 200 million people on Earth are infected. Aim of this study was investigation of prevalence of *Enterobius Vermicularis* in Kermanshah city nurseries. This was cross sectional (descriptive - analysis) study. Cluster Random Sampling was donning. Data were collected by questionnaire. According to the prevalence of this disease, Sample size was determined 92 subjects in Kermanshah city nurseries. Samples were examined in the Kermanshah reference laboratory of parasitology. Chi-square test was used to determine the relationship between infection and different variables. Data were analyzed by using SPSS version 18. In this study, 95 children were observed from kindergartens in Kermanshah. In 14.7% of the children oxyuriasis (pinworm) was diagnosed. In total, 13.7% were infected for the first time. Significant relationship between sex and oxyuriasis where observed. Among these children 7% had sucking habits, 14.3% had itching symptoms and 57% had sign of fatigue. The relationship between father job and mother job with Oxyuriasis were statistically significant. The prevalence of oxyure among children in Kermanshah city kindergartens was 14.7%. Attention to personal hygiene by parents can help in reducing the disease. In all societies, whether poor or rich, if they do not observe hygiene and care towards children, prevalence of this disease is increased. Girls more attention to health and therefore they are less Oxyuriasis.

Key words: *EnterobiusVermicularis*, Oxyure, Pinworm, nursery, Children, Kermanshah

Copyright © 2014 Zahra Sha-Mohammadi et al. This is an open access article distributed under the Creative Commons Attribution License.

1. INTRODUCTION

Oxyuriasis is the most prevalent parasitic diseases in children that caused by *Enterobiusvermicolaris* infections. It is estimated that over 200 million people on Earth are infected. Female worm exits the anus at night and lays their eggs on the skin around the anus that they will contain larvae. Anal itching is a major symptom of the disease. For this reason, the eggs were transferred to the hands,

especially the fingernails, and thereby will be able to infect others and himself (Autoinfection). The Eosinophilia is uncommon and low in this disease (1). Although the worm is often considered as symbiotic (commensally), but in many cases, especially in children, causing symptoms that include anal itching, abdominal pain, nausea and vomiting, irritability, and teeth grinding (2). Advances in medical sciences have decreased to morbidity and mortality in recent years. Neverthe-

less, parasitic infection is one the most health problems in developing countries and so developed countries. It appears to prevalence of infectious diseases is specific to underserved and poor areas, But the interesting thing is that privileged people have this disease too (3-6). Given the above, the rate of infection varies in different parts of the world. In North America and Western Europe, the incidence of infection is about 30% and among children is probably between 60-50% (7). In Iran, the spread of intestinal parasites and oxyuriasis is Unavoidable, because of type of work, life style and health status, unsanitary drinking water and the use of animal and human manure, weakness sewage system, waste spill in water in broken pipe in urban and rural areas, and the weakness in crisis management (8-10). In one study, an increased incidence of urinary tract infection in oxyuriasis was reviewed and it was shown that these cases in girls are more than boys. It has been observed that 57% of children with urinary tract infection were infected by oxyure, while only 10% of children had no urinary tract infection were infected by oxyure (11). Due to lack of time in research related to oxyuriasis recent years, this research was done in Kermanshah. The researchers decided to do this study to determine of prevalence of *VermicularisEnterobius* (pinworm) in Kermanshah city nurseries because of geographical distribution and cultural diversity in rural an urban residents, residents of different levels of social welfare, possibility of neglect of the disease by health officials and the importance of the disease.

2. MATERIALS AND METHODS

This was cross sectional (descriptive - analysis) study. According to the prevalence of this disease, Sample size was determined 92 subjects in Kermanshah city nurseries. Nurseries that have the lowest levels of health were selected according to report of the Welfare Organization of Kermanshah. Nurseries that had been examined and had been trained by experts in personal health and well-being of prevention were excluded. Other kindergartens were entered in the statistical community. In order to ensure of the accuracy of data collection, 100 patients were selected. Cluster Random Sampling was donnig. Data were collected by questionnaire. The questionnaire included questions on age, sex, child care, parental education level, mother's occupation, family size, health status, and healthy habits of children. The city was divided to three levels; high, middle and low. Then from each region 2 or 3 nurseries

and kindergartens were selected. Depending upon the number of children in each kindergarten samples was prepared. Welfare and parental consent was obtained for each nursery. Parasite-specific method of Graham and scotch tape was used for the detection of (3, 12). Teaching points for test and packed slides were given to parents. Two slides with attached tape were delivered for each child. Mothers were advised, two days, once in the morning before defecation, 3-4 cm from the tape attached to the baby anal area for one minute. Then it is removed and attach on the slide, and the slides packed in same day delivery to the nursery. Samples were examined in the reference laboratory parasitology of Kermanshah University of Medical Sciences for searching parasites by light microscopy using objective lenses 10 and 40 (1). Five samples were excluded due to sampling errors by parents. To determine the relationship between infection and different variables chi-square test was used. Data were analyzed by using SPSS version 18.

3. RESULTS AND DISCUSSION

In this study, 95 children were observed from kindergartens in Kermanshah. 49.5% were female and 50.5% were male. In 14.7% of the childrenoxyuriasis (pinworm) was diagnosed. 19% of children who had a positive test had experienced one or more times this disease before. In total, 13.7% were infected for the first time. Among boys and girls 18.8% and 10.6% had positive tests respectively. Significant relationship between sex and oxyuriasis where observed ($P = 0.02$). 32.6% of children who observed good personal hygiene. 54.7% of children who observed moderate personal hygiene and 12.7% were in the poor level of personal hygiene. Personal hygiene in girls was more than boys. But the difference was not statistically significant. 5.3% of all of these children were already sometimes afflicted with this disease. Among these children 7% had sucking habits, 14.3% had itching symptoms and 57% had sign of fatigue. 24.2% of the studied children were from families with high income, 51.6% of them were from middle income and 24.2% had low income family. There was no relationship between income and Oxyuriasis. Most positive tests (60%) were observed in children whose fathers were employed. The relationship between father job and Oxyuriasis were statistically significant ($p = 0.03$). Most positive tests (57.1%) were observed in children whose mothers were employed. 42.9% of mothers were housewives. This difference was statistically

significant ($p = 0.008$). The highest percentages of positive cases (73.6%) were observed in children whose fathers were undergraduate education. Father's education was no significantly associated with oxyuriasis. This study showed that parents' jobs affect oxyuriasis disease. Descriptive variables are shown in a table (Table 1).

Table 1. Prevalence of oxyuriasis based on some variables in Kermanshah city nurseries

Variables	Options	Prevalance			
		Total (95)	Positive cases (14)	Percent	Negative cases (81)
Fathers Job	Self-Employed	38	3	40	40
	Employee	57	11	60	41
Economic Status	Good	23	4	24.2	21
	Middle	49	4	51.6	20
	Bad	23	6	24.2	40
Fathers Ejuication	Diploma and Lower	16	9	16.8	15
	Technician and Bachelor	70	3	72.7	29
	MSc and Higher	9	2	9.5	36
Mothers Ejuication	Diploma and Lower	21	5	22.2	17
	Technician and Bachelor	65	7	68.4	39
	MSc and Higher	9	2	9.4	25
Children Age Group	One year	4	3	4.2	18
	Two years	10	5	10.5	8
	Three years	18	2	18.9	16
	Four years	20	2	21.1	14
	Five years	18	1	18.9	25
	Six years	25	1	26.4	0
Living Area	High	30	3	31.6	28
	Middle	53	4	55.8	32
	Low	12	7	12.6	21

The study findings showed that the patients had a family history status is not indicator, The result of individual and family health status as a major factor in causing the disease has been introduced. Living in determining disease status can now be oxyuriasis (Table 2).

Table 2. Family history Oxyuriasis

Family history of pollution	Negative cases (81)	Positive cases(14)
Year 2013-14	0	2
Before 3013	11	8
Total	11	10

Oxyuriasis is one of parasitic disease in childhood. It has been reported, 18 million people are infected to Oxyure in Canada and America. The infection rates are from 3% to 80%. In the present study all areas were examined, whereas previous studies emphasized the poor location and high population so that the results of these studies are consistent with our findings (1). Based on the results obtained in the present study, the prevalence of oxyure among children in Kermanshah city kindergartens was 14.7%. In studies conducted in Yasooj and Oruma have been reported that prevalence of oxyure was 37.64% and 9%, respectively (9, 13). A study conducted in Korea shown that Oxyuriasis in nursery children was 9.2% (14). High infected levels, because of parents are busy and less attentions to children's health and also lack commitment of kindergarten teachers to this issue. It can be conclude that if children live in impoverished areas, but parents and educators of kindergarten attend to hygiene, catching Oxyuriasis simply not possible (15). In the present study, the oxyuriasis prevalence in boys more than girls, in a study conducted in Orumienursuries had shown similar results (16). Girls more attention to health and therefore they are less Oxyuriasis. The study found that mothers of infected children experienced Oxyuriasis at least twice. The results are consistent with studies outside Iran (17). So the fear of drug use during pregnancy and expressed shame about oxyuriasis causes not go to the doctor and health centers and can increase oxyuriasis. Based on the findings, significant relationship between parental occupation and oxyuriasis prevalence so that the prevalence of oxyuriasis in children who have parents working is higher than others. This result has been achieved in the study of Valizadeh (9). Attention to personal hygiene by parents can help in reducing the disease.

4. CONCLUSION

Oxyuriasis just does not have children, but is most common in children. Living in affluent places and privileged feel to be there to believe that the probability of disease is low oxyuriasis. Only short-term training at an early age cannot prevent diseases. In all societies, whether poor or rich, if they do not observe hygiene and care towards children, prevalence of this disease is increased. Periodic check in public places especially kindergarten could be decreases spread of the oxyuriasis disease.

ACKNOWLEDGMENT

This study is the result of plan (92218) numbering sponsored by the Department of Research and Technology, Kermanshah University of Medical Sciences has been done.

AUTHORS CONTRIBUTION

This work was carried out in collaboration between all authors.

CONFLICT OF INTEREST

Authors have declared that no conflict interests exist.

REFERENCES

1. Garcia LS, Bruckner DA. Diagnostic medical parasitology: American Society for Microbiology (ASM); 1997.
2. Nourozian MB, Youssefi MR. Prevalence of Enterobius vermicularis in Babol Medical School, 2011. World Applied Sciences Journal. 2012;19(5):634-6.
3. Peterson LR, Bailey WR, Scott EG, Finegold SM, Baron EJ. Diagnostic microbiology: Mosby; 1994.
4. Gilman RH, Marquis GS, Miranda E. Prevalence and symptoms of Enterobius vermicularis infections in a Peruvian shanty town. Transactions of the Royal Society of Tropical Medicine and Hygiene. 1991;85(6):761-4.
5. Makhlof S, Sarwat M, Mahmoud D, Mohamad A, Richards Jr F, Flores Z, et al. Parasitic infection among children living in two orphanages in Cairo. Journal of the Egyptian Society of Parasitology. 1994;24(1):137-45.
6. Savioli L, Bundy D, Tomkins A. Intestinal parasitic infections: a soluble public health problem. Transactions of the Royal Society of Tropical Medicine and Hygiene. 1992;86(4):353-4.
7. Marquardt W, Demaree R, Grieve R. Parasitology and Vector Biology, 2000. Academic Press, San Diego, CA.
8. Organization WH. Basic laboratory methods in medical parasitology. 1991.
9. Bagher Nourozian M, Reza Youssefi M. Investigation of Oxyuris (Enterobius vermicularis) prevalence in kindergarten and primary school children of Babol city, Mazandaran, Iran 2009. Annals of Tropical Medicine & Public Health. 2013;6(1).
10. Maraghi S. Prevalence of Enterobius infection in day-care center children in Ahvaz. J Med Sci Ahvaz University. 1997;21:14-8.
11. Sh G, Mobedi E, Ziaee H, Sharif M. Intestinal helminth parasites in dog and jackal in inferent areas Sari in the year 1992-1993. J Mazand Univ Med Sci. 1999;9:22-3.
12. Neva FA, Brown HW. Basic clinical parasitology: Appleton & Lange; 1994.
13. MOGHIMI M, SHARIFI A. PREVALENCE OF INTESTINAL PARASITES IN PRESCHOOL CHILDREN OF YASUJ (1380-1381). ARMAGHAN DANESH. 2002.
14. Yoon H-J, Choi Y-J, Lee S-U, Park H-Y, Huh S, Yang Y-S. Enterobius vermicularis egg positive rate of pre-school children in Chunchon, Korea (1999). The Korean journal of parasitology. 2000;38(4):279-81.
15. Cateau E, Yacoub M, Tavilien C, Becq-Giraudon B, Rodier M-H. Enterobius vermicularis in the kidney: an unusual location. Journal of medical microbiology. 2010;59(7):860-1.
16. HAZRATI TK, MOHAMMADZADEH H, KHASHAVEH S. PREVALENCE OF INTESTINAL PARASITIC INFECTIONS AMONG PRIMARY SCHOOL STUDENTS IN BARANDOOZ-CHAY RURAL REGION OF URMIA, 2007. THE JOURNAL OF URMIA UNIVERSITY OF MEDICAL SCIENCES. 2010.
17. Young C, Tataryn I, Kowalewska-Grochowska KT, Balachandra B. Enterobius vermicularis infection of the fallopian tube in an infertile female. Pathology-Research and Practice. 2010;206(6):405-7.