

Protective Effect of Royal Jelly on *In Vitro* Fertilization (IVF) in Male Mice Treated with Oxymetholone

Ensieh Zahmatkesh, M.Sc.^{1*}, Gholamreza Najafi, Ph.D.², Vahid Nejati, Ph.D.¹

1. Department of Histology and Embryology, Faculty of Science, Urmia University, Urmia, Iran
2. Department of Anatomy and Embryology, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran

*Corresponding Address: P.O.Box: 5756151818, Department of Histology and Embryology, Faculty of Science, Urmia University, Urmia, Iran
Email: ensieh_zahmatkesh@yahoo.com

Received: 14/Jul/2014, Accepted: 23/Oct/2014

Abstract

Objective: This study aimed to investigate the effects of royal jelly (RJ) on catalase, total antioxidant capacity and embryo development in adult mice treated with oxymetholone (OXM).

Materials and Methods: In this experimental study, 32 male and 96 female adult Naval Medical Research Institute (NMRI) mice (7-9 weeks of age) with a ratio of 1:3 for fertilization purposes were randomly divided into 4 groups as follows: i. Control group (n=8) receiving 0.1 ml/mice saline daily by gavage for 30 day, ii. RJ group (n=8) treated with RJ at a dose of 100 mg/kg daily by gavage for 30 days, iii. OXM group (n=8) receiving OXM at the dose of 5 mg/kg daily by gavage for 30 days and iv. RJ+OXM group (n=8) receiving RJ at the dose of 100 mg/kg daily by gavage concomitant with 100 mg/kg OXM administration for 30 days.

Results: Analysis revealed a significant reduction in catalase, total antioxidant, as well as embryo development in OXM group ($P<0.05$). However, RJ group showed a salient recovery in the all of the above mentioned parameters and embryo toxicity.

Conclusion: The results of this study indicated a partially protective effect of RJ against OXM-induced embryo toxicity.

Keywords: Catalase, Embryo, Fertilization, Oxymetholone, Royal Jelly

Cell Journal(yakhteh), Vol 17, No 3, Autumn 2015, Pages: 569-575

Citation: Zahmatkesh E, Najafi Gh, Nejati V. Protective effect of royal Jelly on in vitro fertilization (IVF) in male mice treated with oxymetholone. Cell J. 2015; 17(3): 569-575.

Introduction

Infertility is a fairly common problem and defined by World Health Organization (WHO) failure to conceive after 12 months of unprotected intercourse (1, 2). Infertility has different aspects including psychological and interpersonal stress which severely affects infertile couples. Globally, ranging between 8 and 12% among women of childbearing age, infertility affects 50-80 million people. Some of its aspects important roles in Eastern countries and, become a psychosocial burden (3, 4). A substantial share of the total infertility falls on males (5, 6). Oxymetholone (OXM) is considered as a synthetic derivative of testosterone, made by methylation of 17-alpha carbon and saturation of 5-alpha carbon, and synthesized for

the first time by Ringold in 1959 (7). Anabolic-androgenic steroids (AAS), particularly OXM, are used at small doses (1-5 mg/Kg) for treating anemia, failure to thrive (FTT) in children, and heart failure (8). This type of drug increases erythropoietin production that is followed by affecting bone marrow directly to increase hemoglobin level and red blood cell count (9). Additionally, OXM can promote synthesis of protein, nitrogen retention and deposition of calcium in bones (10). In order to assume anabolic properties, these steroids must be used at 10-100 times their normal (therapeutic) doses in which their adverse reactions occur, (11) having profound effects on male endocrine and reproductive systems. It has been found that AASs-induced low male infertility can be re-

versible. Also, researches have indicated that persistent hypogonadotrophic hypogonadism caused by steroid abuse could be resolved after the AAS withdrawal (12). Royal jelly (RJ) is a milky white highly viscous substance secreted from the salivary gland of the honeybee worker, *Apis mellifera* (Apidae), which is essential in the development of queen bees (13). This substance possesses many physical and chemical properties, including anti-inflammatory, antioxidant, anti-tumor, and immunomodulatory functions in experimental animals (14), which are beneficial to human health, leading to its wide use in commercial and medical products, health food, and cosmetics (15). *In vitro* fertilization (IVF), first successfully applied in 1978, is the last option for infertile couples who failed to conceive. In this international method, different standard therapies like surgery, fertility drugs and artificial insemination are applied to treat infertility cases like male and immunological infertility, infertility caused by endometriosis, and other unexplained conditions (16, 17). This study aims to investigate the effects of RJ on catalase, total antioxidant capacity and embryo development in adult mice treated with OXM.

Materials and Methods

Animal model

In this experimental study, two groups of 7-9-week old adult Naval Medical Research Institute (NMRI) mice, 32 male and 96 female, weighing 30 ± 2 g and 27 ± 2 g, respectively, were purchased from the Animal House of Science Faculty, Urmia University, Urmia, Iran. All animals were allowed free access to water and food ad libitum in controlled conditions of temperature ($22 \pm 2^\circ\text{C}$), relative humidity ($55 \pm 5\%$) and normal photoperiod.

Drugs

OXM was used at a dose of 5 mg/kg (pilot). Drug was dissolved in saline before oral administration. RJ was used at the dose of 100 mg/kg (18).

Experimental protocol

In this investigate, 32 male and 96 female with a ratio of 1:3 for fertilization purposes were divided

into following 4 groups: i. control group (n=8) receiving 0.1 ml/mice saline daily by gavage for 30 days, ii. OXM group (n=8) treated with 5 mg/kg OXM daily by gavage for 30 days, iii. RJ group (n=8) treated with 100 mg/kg RJ daily by gavage for 30 days and iv. RJ+OXM (n=8) group treated with 5 mg/kg OXM by gavage along with 100 mg/kg RJ daily for 30 days. All experimental procedures were approved by the Ethics Committee for Animal Experimentation of Urmia University, Urmia, Iran.

Mouse preparation for *in vitro* fertilization (Harvesting of oocytes)

Each female mouse was injected subcutaneously with 10 IU pregnant mare's serum gonadotropin (PMSG, Boxmeer, Netherlands) 48 hours prior to receiving an intraperitoneal injection of 10 IU human chorionic gonadotropin (hCG, Folligon, Netherlands). The animals were euthanized by dislocation of cervical vertebrae, 12-14 hours after hCG administration. Their ampulla of oviducts were removed and transferred to a petri dish that contained 1 ml human tubular fluid (HTF, Sigma, St. Louis, USA) medium+4 mg/ml bovine serum albumin (BSA, Sigma, St. Louis, USA). Using a stereo microscope (Model TL2, Olympus Co., Tokyo, Japan), the ampullary portion of fallopian tube was found and oocytes were dissected out.

Preparation of culture media for *in vitro* fertilization

One day before conception, required media were prepared for fertilization and incubated 5% CO₂ and 37°C for 12 hours. For each group, separate conception dishes were considered containing 1 ml HTF medium combined with 4 mg/ml BSA. A drop of 500 ml and a few drops of 100 ml per dish were used for washing the IVF dishes and they were then covered with mineral oil (Sigma, St. Louis, USA).

Sperm preparation

Male mice were euthanized by dislocation of cervical vertebrae to prepare needed sperm. Then, the skin of the abdomen was sterilized with 70% ethanol (Merck, USA) and after making an incision in the abdomen and removing the surrounding connective tissue, caudal

epididymis was isolated from testes and placed in a petri dish containing 1 ml HTF medium combined with 4 mg/ml BSA which had reached equilibrium before. After making several cuts in the tail of the epididymis and using pressure in vas deferens, sperm output was placed in an incubator at 5% CO₂ and 37°C for 30 minutes. Spermatozoa were then spread out in the medium.

Ovulation and fertilization in the *in vitro* fertilization laboratory

Between 12 and 14 hours after injection of hCG (in the morning after), female mice were euthanized by dislocation of cervical vertebrae, and the ampule of fallopian tubes were then removed and put in the HTF medium at 37°C. Using dissection techniques, oocytes were removed, washed with the HTF, and transferred to the fertilization droplets under mineral oil containing HTF+BSA. Following capacitation step, sperms (1×10⁶/1 ml HTF) were added to the medium, one million per ml of culture. Fertilization is determined about 4 to 6 hours after releasing the sperm by observing two pre-nucleus. After granulosa cells were denuded and washed, these zygotes were transferred into the fresh pre-equilibrated medium and cultured for five days.

Assessment of two-cell embryos growth

Evaluation of two-cell embryos was done 24 hours after fertilization. The percentage of blastocyst-stage embryos was performed on days 4 and 5 of fertilization.

Measurement of catalase activity

Catalase activity was determined based on its ability to decompose hydrogen peroxide (H₂O₂) in homogenized testicular tissue using the method of Aebi (19). Decomposition of H₂O₂ is considerable by reducing the absorption at 240 nm in an absorption spectrum. For this purpose, 30 mM H₂O₂ (Merck, USA) as substrate and 50 mM phosphate buffer solution (PBS, PH=7, Merck, USA) were used as an alternative substrate in the blank solution. The testis tissue pieces were homogenized in PBS and then centrifuged (Eppendorf AG 5810R, Germany)

at 3400 rpm for 15 minutes. Subsequently 1 ml H₂O₂ was added to 2 ml supernatant and absorbance was measured at 240 nm in a spectrophotometer (pharmacia novaspec, and biochrom, England). The values were expressed in terms of U/g tissue.

Measurement of total antioxidant ferric reducing antioxidant power test

For automatic measurement and evaluation of antioxidants, the ferric reducing antioxidant power (FRAP) was applied. Reduction of ferric to ferrous ions in low PH forms a colored complex of ferrous-tripyridyltriazine (Fe^{II}-TPTZ). Reactive oxygen species (ROS) that are potentially harmful are produced as byproducts of normal aerobic metabolism and usually removed or disabled by antioxidants groups in *in vivo*. In low PH, ferric tripyridyltriazine (Fe^{III}-TPTZ) complex is reduced to form the Fe^{II} and to produces an intense blue color with a high absorption at 593 nm. FRAP reagent (with a ratio of 10:1:1) including 25 ml of (300 mM PBS, 10 mM TPTZ in 40 mM HCL, Merck, USA) and 2.5 ml of 0.0540 g FeCl₃ (Merck, USA) were poured into a flask and brought to volume of 10 ml with distilled water sample was weighted dissolved into 1.5 M of 10% (W/V) KCL buffer (Merck, USA) and grounded in a mortar. The solution was then poured in a micro-tube and centrifuged at 1000 rpm for 5 minutes. One hundred ml of supernatant was poured into a test tube, mixed with 3 mL of FRAP reagent and incubated in water bath at 37°C for 7-10 minutes. The absorbance of blue complex was then determined at 593 nm. Data are expressed in mmol tissue weight (FRAP value) (20, 21).

Statistical analysis

All data were analyzed using the one-way ANOVA by Tukey-Kramer test. The level of significance was considered at P<0.05. All analysis for each sample were done individually.

Results

Catalase concentration

Results show that OXM caused a significant decrease in catalase enzyme compared to the control

group ($P<0.05$). However RJ+OXM group partially compensated this lack attributed to the OXM group (Table 1).

Total antioxidant concentration (FRAP test)

In the OXM group, total antioxidant concentration dropped as compared with the control group, while in the RJ group, there was a significant increase in total antioxidant concentration in comparison with the control group ($P<0.05$). The OXM+RJ showed a significant increase in this regard as compared with the control and OXM group

($P<0.05$, Table 1).

Fertilization and embryonic development

In this study, OXM caused a significant decrease in fertilization rate and percentage of blastocysts ($P<0.05$) and a significant increase in the percentage of arrested embryos ($P<0.05$), but did not affect percentage of the two-cell embryos. RJ caused an increase in fertilization rate and percentage of blastocyst and a decrease in percentage of the arrested embryos ($P<0.05$) and two-cell embryo (Table 2, Figs.1, 2).

Table 1: Effects of OXM, RJ as well as OXM and RJ on catalase activity and FRAP level in adult male mice

Groups	Catalase (U/g tissue)	FRAP (mMol/g tissue)
Control	0.516 ± 0.060	126.66 ± 0.927
OXM	0.230 ± 0.026 ^a	103.33 ± 4.371
RJ	0.530 ± 0.060 ^b	135.43 ± 7.361 ^b
OXM+RJ	0.320 ± 0.020	214.33 ± 8.685

Values are expressed as mean ± standard deviation (n=8).

^a; Significant differences as compared with the control group ($P<0.05$), ^b; Significant differences as compared with the OXM group ($P<0.05$), OXM; Oxymetholone, RJ; Royal jelly and FRAP; Ferric reducing antioxidant power.

Table 2: Effect of OXM , RJ as well as OXM and RJ on fertilization and embryonic development rates

Groups	Oocyte (n)	Zygote (n/%)	Two cell (%)	Blastocyst (%)	Arrest (%)
Control	38	(33) 86.72 ± 1.92	78.48 ± 2.89	73.04 ± 0.71	14.65 ± 1.00
OXM	47	(32) 67.81 ± 1.55 ^a	77.18 ± 4.34	40.64 ± 2.02 ^a	27.17 ± 3.58 ^a
RJ	85	(76) 89.32 ± 0.41 ^b	88.71 ± 2.64	82.40 ± 2.10 ^b	6.92 ± 2.51 ^b
OXM+RJ	58	(44) 75.76 ± 2.47	63.43 ± 1.92	60.51 ± 1.20 ^b	15.24 ± 1.27 ^b

Values are expressed as mean ± standard deviation (n=8).

^a; Significant differences as compared with the control group ($P<0.05$), ^b; Significant differences as compared with the OXM group ($P<0.05$), OXM; Oxymetholone and RJ; Royal jelly.




Fig.1: Fertilized and infertilized oocytes in OXM group ($\times 200$). OXM; Oxymetholone.




Fig.2: Blastocyst embryos in control group ($\times 200$).

Discussion

AAS alter the function of the hypothalamic- pituitary-gonadal (HPG) axis that results in affecting the target reproductive tissues. Acting on the

hypothalamus, circulating testosterone regulates the gonadotropin-releasing hormone (GnRH) release. Introducing exogenous AAS to the body increases androgen levels that cause a decrease in luteinizing hormone (LH) release from the pituitary, resulting in the suppression of endogenous testosterone production (22, 23), which is found to be reversible (12). ROS induced damage is major factors in male infertility (24, 25). In the last decade, numerous evidences have been gathered about etiology of oxidative stress (OS) in male infertility (26, 27). Our findings indicated that OXM causes a significant decrease in catalase activity and also total antioxidant concentration. The total antioxidant potential is a measure of the individual's ability to cope with or prevent a state of OS. It is noted that a state of disequilibrium is the rate of generation of ROS that overwhelms an individual's ability to remove them (28). Our findings also found that OXM reduces the fertilization rate and percentage of blastocysts, but increases percentage of arrested embryos which is probably due to the effects of OS system. MacLeod (29), observed that human spermatozoa produces ROS, i.e. highly reactive derivatives of oxygen such as H_2O_2 , and have harmful effects on different sperm functions. ROS involvement in etiology of male infertility is now well-established (30, 31). Since sperm functions are extremely sensitive to ROS -because of high content of polyunsaturated fatty acids (PUFA) and limited ability of DNA repair- the role of ROS and OS in human sperm function and pathophysiology of male infertility have been intensively investigated (32, 33). Very recently, scientists revealed that ROS not only affects the fertilization process negatively, but also raises serious questions about the health and well-being of the progeny since the male germ cell is exposed to high oxidants concentrations (34, 35). Furthermore the concept of OS causing damage to spermatozoa by impairing various functions of sperm such as motility, acrosome reaction or DNA integrity is widely accepted nowadays (26, 36, 37). Several studies reported positive effects of RJ which is a homogeneous substance secreted by worker honey bees for feeding young larvae and the adult queen bee on animal reproduction (38). The physical and chemical properties of RJ have been described in detail elsewhere (39, 40). Intramuscular or oral administration of RJ has proven to be effective in improving estrus responses and

pregnancy rate (41). The results of this study indicate that RJ increases catalase activity and total antioxidant capacity. Antioxidants including the enzymes superoxide dismutase, catalase and glutathione peroxidase that are constantly produced during normal metabolic and physiological processes (42) counteract with the deleterious effects of ROS mammalian spermatozoa with rich polyunsaturated fatty acids are very susceptible to ROS attack which results in decreased sperm motility, axonemal damage, decreased sperm viability, and increased midpiece morphology defects with deleterious effects on sperm capacitation and acrosome reaction that is presumably due to rapid loss of intracellular ATP (43). The key mechanism of this ROS-induced sperm damage is considered to be lipid peroxidation of sperm membrane that leads to infertility (44). Several studies using various antioxidants alone or in combination with others have shown a significant reduction in seminal ROS levels (45, 46) and improvement in sperm count and motility (47, 48), while other studies have found contrary results (49, 50). We also found that RJ increases fertility-related parameters such as fertilization rate and blastocysts percentage and also reduces percentage of arrested embryos which is probably due to its effect on the total antioxidant capacity. Antioxidants levels in seminal plasma of infertile men are significantly low (51), Renard et al. (52) reported that intravaginal administration of Egyptian bee honey and RJ might be a reasonable and effective method to treat infertility caused by asthenozoospermia. Amino acid content of both honey and RJ may play a main role in fertility either by enhancing acrosome reaction and sperm motility or by improving fertilization. Also sperm motility may be further enhanced by the short chain fatty acids specific to RJ, especially 10-hydroxy-2-decenoic acid (45).

Conclusion

It can be concluded that with the decline in total antioxidant capacity, OXM reduces fertility. Due to beneficial biological properties of its components, RJ showed to increase antioxidant capacity; therefore, it prevents embryo toxicity manifested by OXM when given after OXM administration. As the result, RJ with its antioxidant properties increases fertility.

Acknowledgments

This study was financially supported by Urmia University. The authors thank the personnel of Science Faculty of Urmia University for their assistant. There is no conflict of interest in this article.

References

- Cooper TG, Noonan E, von Eckardstein S, Auger J, Baker HW, Behre HM, et al. World Health Organization reference values for human semen characteristics. *Hum Reprod Update*. 2010; 16(3): 231-245.
- de Melo-Martin I. On cloning human beings. *Bioethics*. 2002; 16(3): 246-265.
- Schmidt L. Psychosocial burden of infertility and assisted reproduction. *Lancet*. 2006; 367(9508): 379-380.
- Whiteford LM, Gonzalez L. Stigma: the hidden burden of infertility. *Soc Sci Med*. 1995; 40(1): 27-36.
- Tekatli H, Schouten N, van Dalen T, Burgmans I, Smakman N. Mechanism, assessment, and incidence of male infertility after inguinal hernia surgery: a review of the pre-clinical and clinical literature. *Am J Surg*. 2012; 204(4): 503-509.
- Mbizvo MT, Danso AP. Male reproductive function: a review on sperm fertilising ability and infertility. *Cent Afr J Med*. 1991; 37(2): 40-47.
- Bi H, Du P, Masse R. Studies on anabolic steroids--8. GC/MS characterization of unusual seco acidic metabolites of oxymetholone in human urine. *J Steroid Biochem Mol Biol*. 1992; 42(2): 222-242.
- Shahidi NT, Calantoff DV. The role of puberty in red-cell production in hereditary haemolytic anemias. *Br J Haematol*. 1969; 17(4): 335-342.
- Barcelo AC, Martinez MP, Conti MI, Champin GM, Alippi RM, Bozzini CE. Inability of oxymetholone to elicit an erythropoietin hyper secretory state in orchidectomised male mice. *Comp Clin Pathol*. 2005; 13(4): 153-157.
- Karimi Jashni H, Bandak Sh, Mahjoor AA. Effects of oxymetholone on kidney tissue of one-day old offspring of pregnant rats. *J Jahrom Univ Med Sci*. 2011; 9(3): 8-13.
- Christiansen K. Behavioural effects of androgen in men and women. *J Endocrinol*. 2001; 170(1): 39-48.
- Alen M, Reinila M, Vihko R. Response of serum hormones to androgen administration in power athletes. *Med Sci Sports Exerc*. 1985; 17(3): 354-359.
- Lombardi C, Senna GE, Gatti B, Feligioni M, Riva G, Bonadonna P, et al. Allergic reactions to honey and royal jelly and their relationship with sensitization to compositae. *Allergol Immunopathol (Madr)*. 1998; 26(6): 288-290.
- Hattori N, Nomoto H, Fukumitsu H, Mishima S, Furukawa S. Royal jelly and its unique fatty acid, 10-hydroxy-trans-2-decenoic acid, promote neurogenesis by neural stem/progenitor cell in vitro. *Biomed Res*. 2007; 28(5): 261-266.
- Guo H, Kouzuma Y, Yonekura M. Structures and properties of antioxidative peptides derived from royal jelly protein. *Food Chem*. 2009; 113(1): 238-245.
- Stephoe PC, Edwards RG. Birth after the reimplantation of a human embryo. *Lancet*. 1978; 2(8085): 366.
- Pagidas K, Hemmings R, Falcone T, Miron P. The effect of antisperm autoantibodies in male or female partners undergoing in vitro fertilization-embryo transfer. *Fertil Steril*. 1994; 62(2): 363-369.
- Silici S, Ekmekcioglu O, Eraslan G, Demirtas A. Antioxidative effect of royal jelly in cisplatin-induced testes damage. *Urology*. 2009; 74(3): 545-551.

19. Aebi H. Catalase in vitro. *Methods Enzymol.* 1984; 105: 121-126.
20. Hosseinzadeh H, sadeghnia HR. Safranal, a constituent of *corcus sativus* (saffron), attenuated cerebral ischemia induced oxidative damage in rat hippocampus. *J Pharm Pharm Sci.* 2005; 8(3): 394-399.
21. Benzie IF, Strain JJ. The ferric reducing ability of plasma (FRAP) as measure of "antioxidant power": the FRAP assay. *Anal Biochem.* 1996; 239(1): 70-76.
22. Johnson LC, Fisher G, Silvester LJ, Ho fheins CC. Anabolic steroid: effects on strength, body weight, oxygen uptake, and spermatogenesis upon mature males. *Med Sci Sports.* 1972; 4(1): 43-45.
23. Klaiber EL, Henzl MR, Lloyd CW, Segre EJ. Corpus luteum-inhibiting action of oxymetholone. *J Clin Endocrinol Metab.* 1973; 36(1): 142-147.
24. Pasqualotto FF, Sharma RK, Nelson DR, Thomas AJ, Agarwal A. Relationship between oxidative stress, semen characteristics, and clinical diagnosis in men undergoing infertility investigation. *Fertil Steril.* 2000; 73(3): 459-464.
25. Ollero M, Gil-Guzman E, Lopez MC, Sharma RK, Agarwal A, Larson K, et al. Characterization of subsets of human spermatozoa at different stages of maturation: implicatin in the diagnosis and treatment of male infertility. *Hum Reprod.* 2001; 16(9): 1912-1921.
26. Aitken RJ, Clarkson JS. Cellular basis of defective sperm function and its association with the genesis of reactive oxygen species by human spermatozoa. *J Reprod Fertil.* 1987; 81(2): 459-69.
27. Aitken RJ. Free radicals, lipid peroxidation and sperm function. *Reprod Fertil Dev.* 1995; 7(4): 659-668.
28. Cerutti PA, Trump BF. Inflammation and oxidative stress in carcinogenesis. *Cancer Cells.* 1991; 3(1): 1-7.
29. MacLeod J. The role of oxygen in the metabolism and motility of human spermatozoa. *Am J Physiol.* 1943; 138(85): 512-518.
30. Aitken RJ. A free radical theory of male infertility. *Reprod Fertil Dev.* 1994; 6(1): 19-23.
31. Sharma RK, Agarwal A. Role of reactive oxygen species in male infertility. *Urology.* 1996; 48(6): 835-850.
32. Griveau JF, Le Lannou D. Reactive oxygen species and human spermatozoa: physiology and pathology. *Int J Androl.* 1997; 20(2): 61-69.
33. Shen H, Ong C. Detection of oxidative DNA damage in human sperm and its association with sperm function and male infertility. *Free Radic Biol Med.* 2000; 28(4): 529-536.
34. Lewis SE, Aitken RJ. DNA damage to spermatozoa has impacts on fertilization and pregnancy. *Cell Tissue Res.* 2005; 322(1): 33-41.
35. Zini A, Boman JM, Belzile E, Ciampi A. Sperm DNA damage is associated with an increased risk of pregnancy loss after IVF and ICSI: systematic review and meta-analysis. *Hum Reprod.* 2008; 23(12): 2663-2668.
36. Iwasaki A, Gagnon C. Formation of reactive oxygen species in spermatozoa of infertile patients. *Fertil Steril.* 1992; 57(2): 409-416.
37. Williams AC, Ford WC. Relationship between reactive oxygen species production and lipid peroxidation in human sperm suspensions and their association with sperm function. *Fertil Steril.* 2005; 83(4): 929-936.
38. Husein MQ, Kridli RT, Humphrey WD. Effect of royal jelly on estrus synchronization and pregnancy rate of ewes using flourogestone acetate sponges. *J Anim Sci.* 1999; 77 Suppl 1: 431.
39. Nation JL, Robinson FA. Concentration of some major and trace elements in honey bees, royal jelly and pollens, determined by atomic absorption spectrophotometry. *J Apic Res.* 1971; 10(1): 35-43.
40. Hanes J, Simuth J. Identification and partial characterization of the major royal jelly protein of the honey bee (*Apis mellifera* L.). *J Apic Res.* 1992; 31(1): 22-26.
41. Husein MQ, Kridli RT. Reproductive responses following royal jelly treatment administered orally or intramuscularly into progesterone-treated Awassi ewes. *Anim Reprod Sci.* 2002; 74(1-2): 45-53.
42. Kensler TW, Egner PA, Wang JB, Zhu YR, Zhang BC, Qian GS, et al. Strategies for chemoprevention of liver cancer. *Eur J Cancer Prev.* 2002; 11 Suppl 2: S58-64.
43. Lenzi A, Cualosso F, Gandini L, Lombardo F, Dondero F. Placebo-controlled, double-blind, cross-over trial of glutathione therapy, in male infertility. *Hum Reprod.* 1993; 8(10): 1657-1662.
44. Agarwal A, Ikemoto I, Loughlin KR. Relationship of sperm parameters with levels of reactive oxygen species in semen specimens. *J Urol.* 1994; 152(1): 107-110.
45. Comhaire FH, Christophe AS, Zalata AA, Dhooge WS, Mahmoud AM, Depudt CE. The effects of combined conventional treatment, oral antioxidants and essential fatty acids on sperm biology in subfertile men. *Prostaglandins Leukot Essent Fatty Acids.* 2000; 63(3): 159-165.
46. Vicari E, Calogero AE. Effects of treatment with carnitines in infertile patients with prostatic-vesiculo-epididymitis. *Hum Reprod.* 2001; 16(11): 2338-2342.
47. Lenzi A, Sgro P, Salacone P, Paoli D, Gilio B, Lombardo F, et al. A placebo-controlled double-blind randomized trial of the use of combined L-carnitine and L-acetyl-carnitine treatment in men with asthenozoospermia. *Fertil Steril.* 2004; 81(6): 1578-1584.
48. Akmal M, Qadri JQ, Al-Waili NS, Thangal S, Haq A, Saaloom KY. Improvement in human semen quality after oral supplementation of vitamin C. *J Med Food.* 2006; 9(3): 440-442.
49. Greco E, Iacobelli M, Rienzi L, Ubaldi F, Ferrero S, Tesarik J. Reduction of the incidence of sperm DNA fragmentation by oral antioxidant treatment. *J Androl.* 2005; 26(3): 349-353.
50. Menezes YJ, Hazout A, Panteix G, Robert F, Rollet J, Cohen-Bacrie P, et al. Antioxidants to reduce sperm DNA fragmentation: an unexpected adverse effect. *Reprod Biomed Online.* 2007; 14(4): 418-421.
51. Halliwell B. How to characterize a biological antioxidant. *Free Radic Res Commun.* 1990; 9(1): 1-32.
52. Renard P, Grizard G, Griveau JF, Sion B, Boucher D, Le Lannou D. Improvement of motility and fertilization potential of postthaw human sperm using glutamine. *Cryobiology.* 1996; 33(3): 311-319.