

## ON WEAK CONVERGENCE THEOREM FOR NONSELF I-QUASI-NONEXPANSIVE MAPPINGS IN BANACH SPACES

Pankaj Kumar Jhade and and A. S. Saluja

ABSTRACT. In this paper, we construct Ishikawa iteration scheme with error for nonself I-quasi nonexpansive maps and establish the weak convergence of a sequence of Ishikawa iteration of nonself I-quasi nonexpansive maps in a Banach space which satisfies Opial's condition.

### 1. Introduction and Preliminaries

Let  $K$  be a nonempty convex subset of a real Banach space  $E$ . The map  $T : K \rightarrow K$  is nonexpansive if  $\|Tx - Ty\| \leq \|x - y\|$  for all  $x, y \in K$ . Nonexpansive selfmaps ever since their introduction, remained a popular area of research in various fields. Iterative construction of fixed points of these maps is a fascinating field of research. In 1967, Browder [3] studied the iterative construction of fixed points of nonexpansive self maps on closed and convex subset of a Hilbert space.

Two most popular iteration procedure for obtaining fixed points of  $T$ , if they exists, are : Mann iteration [12], defined by

$$(1.1) \quad \begin{aligned} x_1 &\in K, \\ x_{n+1} &= (1 - \alpha_n) x_n + \alpha_n T x_n \quad n \geq 1 \end{aligned}$$

and, Ishikawa Iteration [8], defined by

$$(1.2) \quad \begin{aligned} x_1 &\in K, \\ x_{n+1} &= (1 - \alpha_n) x_n + \alpha_n T y_n \\ y_n &= (1 - \beta_n) x_n + \beta_n T x_n, \quad n \geq 1 \end{aligned}$$

---

2000 *Mathematics Subject Classification*. Primary 47H10; Secondary 47H09.

*Key words and phrases*. Fixed points, quasi-nonexpansive nonself mappings, Opial's condition, Ishikawa iteration scheme with error.

for certain choices of  $\{\alpha_n\}, \{\beta_n\} \subset [0, 1]$ . If we take  $\beta_n = 0$  in (1.2) then we obtain iteration (1.1). In sequel, let  $F(T) = \{x \in K : Tx = x\}$  be the set of fixed points of a mapping  $T$ .

The first nonlinear ergodic theorem was proved by Baillon [5] for general nonexpansive mappings in Hilbert space  $H$ : If  $K$  is a closed and convex subset of  $H$  and  $T$  has a fixed point, then for all  $x \in K$ ,  $\{T^n x\}$  is weakly almost convergent, as  $n \rightarrow \infty$ , to a fixed point of  $T$ . It was also shown by Pazy [1] that if  $H$  is a real Hilbert space and  $(\frac{1}{n}) \sum_{i=0}^{n-1} T^i x$  converges weakly, as  $n \rightarrow \infty$ , to  $y \in K$ , then  $y \in F(T)$ .

The concept of quasi-nonexpansive mapping was initiated by Tricomi in 1941 for real functions. Diaz and Metcalf ([6]) and Dotson ([11]) studied quasi-nonexpansive mappings in Banach spaces. Kirk ([10]) gave this concept in metric spaces which we adopt to a normed space as follows:  $T$  is called a quasi-nonexpansive mapping provided  $\|Tx - p\| \leq \|x - p\|$  for all  $x \in K$  and  $p \in F(T)$ .

Recall that a Banach space  $E$  is said to be uniformly convex if for each  $r$  with  $0 \leq r \leq 2$ , the modulus of convexity of  $E$  given by

$$\delta(r) = \inf \left\{ 1 - \frac{1}{2} \|x + y\| : \|x\| \leq 1, \|y\| \leq 1, \|x - y\| \geq r \right\}$$

satisfies the inequality  $\delta(r) > 0$ .

The space  $E$  is said to satisfy Opial's condition ([14]) if, for each sequence  $\{x_n\}$  in  $E$ , the condition  $x_n \rightarrow x$  implies that  $\overline{\lim}_{n \rightarrow \infty} \|x_n - x\| < \overline{\lim}_{n \rightarrow \infty} \|x_n - y\|$  for all  $y \in E$  with  $y \neq x$ .

The following definitions and Lemma will be needed for the proof of our result.

Let  $K$  be a subset of a normed space  $E = (E, \|\cdot\|)$  and  $T$  and  $I$  are self mappings of  $K$ . Then  $T$  is called  $I$ -nonexpansive on  $K$  if  $\|Tx - Ty\| \leq \|Ix - Iy\|$ .

$T$  is called  $I$ -quasi-nonexpansive on  $K$  if  $\|Tx - p\| \leq \|Ix - p\|$  for all  $x, y \in K$  and  $p \in F(T) \cap F(I)$ .

Let  $E$  be a real Banach space and  $K$  be a closed convex subset of  $E$ . A mapping  $T : K \rightarrow K$  is said to be demi-closed at the origin if, for any sequence  $\{x_n\}$  in  $K$ , the condition  $x_n \rightarrow x_0$  weakly  $Tx_n \rightarrow 0$  strongly imply  $Tx_0 = 0$ .

REMARK 1.1. If  $I$  is an identity map then  $I$ -nonexpansive maps and  $I$ -quasi-nonexpansive mappings reduces to nonexpansive and quasi nonexpansive mappings.

A subset  $K$  of  $E$  is said to be a retract of  $E$  if there exists a continuous map  $P : E \rightarrow K$  such that  $Px = x$  for all  $x \in K$ . A map  $P : E \rightarrow E$  is a retraction if  $P^2 = P$ . It easily follows that if a map  $P$  is a retraction, then  $Py = y$  for all  $y$  in the range of  $P$ . A set  $K$  is optimal if each point outside  $K$  can be moved to be closer to all points of  $K$ . Note that every nonexpansive retract is optimal. In strictly convex Banach spaces, optimal sets are closed and convex. However, every closed convex subset of a Hilbert space is optimal and also a nonexpansive retract.

LEMMA 1.1. ([15]) Let  $\{s_n\}$  and  $\{t_n\}$  be two nonnegative real sequences satisfying  $s_{n+1} \leq s_n + t_n$  for all  $n \geq 1$ . If  $\sum_{n=1}^{\infty} t_n < \infty$ , then  $\lim_{n \rightarrow \infty} s_n$  exists.

LEMMA 1.2. ([3]) *Let  $K$  be a nonempty closed convex subset of a uniformly convex Banach space and let  $T : K \rightarrow E$  be a nonexpansive map. Then  $I - T$  is demi-closed at 0.*

LEMMA 1.3. ([16]) *Suppose that  $E$  is a uniformly convex Banach space and  $0 < p \leq t_n \leq q < 1$  for all  $n \in N$ . Suppose further that  $\{x_n\}$  and  $\{y_n\}$  are sequences in  $E$  such that  $\limsup_{n \rightarrow \infty} \|x_n\| \leq r$ ,  $\limsup_{n \rightarrow \infty} \|y_n\| \leq r$  and  $\lim_{n \rightarrow \infty} \|t_n x_n + (1 - t_n) y_n\| = r$  hold for some  $r \geq 0$ . Then  $\lim_{n \rightarrow \infty} \|x_n - y_n\| = 0$*

There are many results on fixed points on nonexpansive and quasi-nonexpansive mappings in Banach spaces and metric spaces. For example Petryshyn and Williamson ([13]) studied the weak and strong convergence to a fixed points of quasi-nonexpansive maps. Their analysis was related to the convergence of Mann iterates studied by Dotson ([11]). Subsequently, Ghosh and Debnath ([7]) discussed the convergence of Ishikawa iterates of quasi-nonexpansive mappings in Banach spaces. In [9], the weak convergence theorem for  $I$ -asymptotically quasi-nonexpansive mapping defined in Hilbert space was proved.

In [2], Rhoades and Temir considered  $T$  and  $I$  self mappings of  $K$ , where  $T$  is an  $I$ -nonexpansive mapping. They established the weak convergence of sequence of Mann iterates to a common fixed point of  $T$  and  $I$ . Subsequently, Kiziltunc and Ozdemir [4] considered  $T$  and  $I$  be nonself mappings of  $K$  with  $T$  is  $I$ -nonexpansive mapping and establish the weak convergence theorem of the sequence of Ishikawa iterates to a common fixed point of  $T$  and  $I$ .

In this paper, we consider  $T$  and  $I$  nonself mappings of  $K$ , where  $T$  is an  $I$ -quasi nonexpansive mapping and establish the weak convergence of the sequence of Ishikawa iterates with error to a common fixed point of  $T$  and  $I$ .

**Iteration Scheme 1.4** [Ishikawa Iteration with error]: Let  $E$  be a uniformly convex Banach space, let  $K$  be a nonempty convex subset of  $E$  with  $P$  as a nonexpansive retraction. Let  $T : K \rightarrow E$  be a given nonself mapping. The Ishikawa iterative scheme with error is defined as follows:

$$(1.3) \quad \begin{cases} x_1 \in K \\ x_{n+1} = P(\alpha_n x_n + \beta_n T y_n + \gamma_n u_n) \\ y_n = P(\alpha'_n x_n + \beta'_n T x_n + \gamma'_n v_n), \quad n \geq 1 \end{cases}$$

Where  $\{\alpha_n\}, \{\beta_n\}, \{\gamma_n\}, \{\alpha'_n\}, \{\beta'_n\}$  and  $\{\gamma'_n\}$  are real sequences in  $[0, 1]$  such that  $\alpha_n + \beta_n + \gamma_n = 1 = \alpha'_n + \beta'_n + \gamma'_n$ ; and  $\{u_n\}, \{v_n\}$  are bounded sequences in  $K$ .

## 2. Main Results

Before proving our main result we begin with the following lemmas.

LEMMA 2.1. *Let  $K$  be a closed convex bounded subset of a uniformly convex Banach space  $E$  and let  $T, I$  be two nonself mappings with  $T$  be  $I$ -quasi-nonexpansive mapping,  $I$  a nonexpansive mapping on  $K$ . If  $\{x_n\}$  is defined as in (1.3) where  $\{\alpha_n\}, \{\beta_n\}, \{\gamma_n\}, \{\alpha'_n\}, \{\beta'_n\}$  and  $\{\gamma'_n\}$  are real sequences in  $[0, 1]$  such that  $\alpha_n +$*

$\beta_n + \gamma_n = 1 = \alpha'_n + \beta'_n + \gamma'_n$  ;  $\sum_{n=1}^{\infty} \gamma_n < \infty$ ,  $\sum_{n=1}^{\infty} \gamma'_n < \infty$  ;  $\{u_n\}$  and  $\{v_n\}$  are bounded sequences in  $K$ , then  $\lim_{n \rightarrow \infty} \|x_n - p\|$  exists.

PROOF. For  $p \in F(T) \cap F(I)$ , we have

$$\begin{aligned}
(2.1) \quad \|x_{n+1} - p\| &= \|P(\alpha_n x_n + \beta_n T y_n + \gamma_n u_n) - p\| \\
&\leq \alpha_n \|x_n - p\| + \beta_n \|T y_n - p\| + \gamma_n \|u_n - p\| \\
&\leq \alpha_n \|x_n - p\| + \beta_n \|I y_n - p\| + \gamma_n \|u_n - p\| \\
&\leq \alpha_n \|x_n - p\| + \beta_n \|y_n - p\| + \gamma_n \|u_n - p\|
\end{aligned}$$

where

$$\begin{aligned}
(2.2) \quad \|y_n - p\| &= \|P(\alpha'_n x_n + \beta'_n T x_n + \gamma'_n v_n) - p\| \\
&\leq \alpha'_n \|x_n - p\| + \beta'_n \|T x_n - p\| + \gamma'_n \|v_n - p\| \\
&\leq \alpha'_n \|x_n - p\| + \beta'_n \|I x_n - p\| + \gamma'_n \|v_n - p\| \\
&\leq \alpha'_n \|x_n - p\| + \beta'_n \|x_n - p\| + \gamma'_n \|v_n - p\|
\end{aligned}$$

Substituting the value of (2.2) into (2.1) we obtain,

$$\begin{aligned}
\|x_{n+1} - p\| &\leq (\alpha_n + \alpha'_n \beta_n + \beta_n \beta'_n) \|x_n - p\| + \gamma_n \|u_n - p\| + \beta_n \gamma'_n \|v_n - p\| \\
&\leq ((1 - \beta_n) + (1 - \beta'_n) \beta_n + \beta_n \beta'_n) \|x_n - p\| + \gamma_n \|u_n - p\| + \beta_n \gamma'_n \|v_n - p\| \\
&\leq \|x_n - p\| + d_n
\end{aligned}$$

where  $d_n = \gamma_n \|u_n - p\| + \beta_n \gamma'_n \|v_n - p\|$

Since  $\sum_{n=1}^{\infty} \gamma_n < \infty$ , and  $\sum_{n=1}^{\infty} \gamma'_n < \infty$  implies that  $\sum_{n=1}^{\infty} d_n < \infty$  and by Lemma (1.1)  $\lim_{n \rightarrow \infty} \|x_n - p\|$  exists. This completes the proof of the lemma.  $\square$

LEMMA 2.2. Let  $E$  be a uniformly convex Banach space and let  $K$  be a nonempty closed convex subset of  $E$ . Let  $T : K \rightarrow E$  be a  $I$ -quasi-nonexpansive mapping with  $F(T) \cap F(I) \neq \phi$  and  $I$  a nonexpansive mapping. Let  $\{\alpha_n\}, \{\beta_n\}, \{\gamma_n\}, \{\alpha'_n\}, \{\beta'_n\}$  and  $\{\gamma'_n\}$  are real sequences in  $[0, 1]$  such that  $\alpha_n + \beta_n + \gamma_n = 1 = \alpha'_n + \beta'_n + \gamma'_n$  and  $\varepsilon \leq \beta_n, \beta'_n \leq 1 - \varepsilon$  for all  $n \in N$  and some  $\varepsilon > 0$  ;  $\{u_n\}$  and  $\{v_n\}$  are bounded sequences in  $K$ . Then for the sequence  $\{x_n\}$  given by (1.3), we have  $\lim_{n \rightarrow \infty} \|x_n - T x_n\| = 0$ .

PROOF. For any  $p \in F(T) \cap F(I)$ , set

$$\begin{aligned}
r_1 &= \sup \{\|u_n - p\| : n \geq 1\}, \\
r_2 &= \sup \{\|v_n - p\| : n \geq 1\}, \\
r_3 &= \sup \{\|x_n - p\| : n \geq 1\}, \\
r &= \max \{r_i : 1 \leq i \leq 3\}
\end{aligned}$$

Now consider

$$\begin{aligned}
\|y_n - p\| &= \|P(\alpha'_n x_n + \beta'_n T x_n + \gamma'_n v_n) - p\| \\
&\leq \alpha'_n \|x_n - p\| + \beta'_n \|T x_n - p\| + \gamma'_n \|v_n - p\| \\
&\leq \alpha'_n \|x_n - p\| + \beta'_n \|I x_n - p\| + \gamma'_n \|v_n - p\| \\
&\leq \alpha'_n \|x_n - p\| + \beta'_n \|x_n - p\| + \gamma'_n \|v_n - p\| \\
&\leq (\alpha'_n + \beta'_n) \|x_n - p\| + \gamma'_n \|v_n - p\| \\
(2.3) \quad &\leq \|x_n - p\| + \gamma'_n r
\end{aligned}$$

Since by Lemma (2.1)  $\lim_{n \rightarrow \infty} \|x_n - p\|$  exists. Let  $\lim_{n \rightarrow \infty} \|x_n - p\| = c$ , then by the continuity of  $T$  the conclusion follows.

Now, let  $c > 0$ . We claim that  $\lim_{n \rightarrow \infty} \|x_n - T x_n\| = 0$ .

Since  $\{u_n\}$  and  $\{v_n\}$  are bounded, it follows that  $\{u_n - x_n\}$  and  $\{v_n - x_n\}$  are bounded.

Taking limit sup on both sides in the inequality (2.3), we have

$$(2.4) \quad \limsup_{n \rightarrow \infty} \|y_n - p\| < c$$

Next consider,

$$\begin{aligned}
\|T y_n - p + \gamma_n (u_n - x_n)\| &\leq \|T y_n - p\| + \gamma_n \|u_n - x_n\| \\
&\leq \|I y_n - p\| + \gamma_n r \\
&\leq \|y_n - p\| + \gamma_n r
\end{aligned}$$

Taking limit sup on both sides in the above inequality and using (2.4), we get

$$\limsup_{n \rightarrow \infty} \|T y_n - p + \gamma_n (u_n - x_n)\| \leq c$$

Then  $\|x_n - p + \gamma_n (u_n - x_n)\| \leq \|x_n - p\| + \gamma_n \|u_n - x_n\| \leq \|x_n - p\| + \gamma_n r$  yields

$$\limsup_{n \rightarrow \infty} \|x_n - p + \gamma_n (u_n - x_n)\| \leq c$$

Again  $\lim_{n \rightarrow \infty} \|x_{n+1} - p\| = c$  means that

$$(2.5) \quad \liminf_{n \rightarrow \infty} \|\beta_n (T y_n - p + \gamma_n (u_n - x_n)) + (1 - \beta_n) (x_n - p + \gamma_n (u_n - x_n))\| \geq c$$

On the other hand we have

$$\begin{aligned}
&\|\beta_n (T y_n - p + \gamma_n (u_n - x_n)) + (1 - \beta_n) (x_n - p + \gamma_n (u_n - x_n))\| \\
&\leq \beta_n \|T y_n - p\| + (1 - \beta_n) \|x_n - p\| + \gamma_n \|u_n - x_n\| \\
&\leq \beta_n \|I y_n - p\| + (1 - \beta_n) \|x_n - p\| + \gamma_n \|u_n - x_n\| \\
&\leq \beta_n \|y_n - p\| + (1 - \beta_n) \|x_n - p\| + \gamma_n \|u_n - x_n\| \\
&\leq \beta_n (\|x_n - p\| + \gamma'_n r) + (1 - \beta_n) \|x_n - p\| + \gamma_n \|u_n - x_n\| \\
&\leq \|x_n - p\| + \gamma'_n r + \gamma_n r
\end{aligned}$$

Therefore we obtain

$$(2.6) \quad \limsup_{n \rightarrow \infty} \|\beta_n(Ty_n - p + \gamma_n(u_n - x_n)) + (1 - \beta_n)(x_n - p + \gamma_n(u_n - x_n))\| \leq c$$

From (2.5) and (2.6) we get

$$\lim_{n \rightarrow \infty} \|\beta_n(Ty_n - p + \gamma_n(u_n - x_n)) + (1 - \beta_n)(x_n - p + \gamma_n(u_n - x_n))\| = c$$

Hence applying Lemma (1.3) we have  $\lim_{n \rightarrow \infty} \|Ty_n - x_n\| = 0$ .

Since  $P$  is a nonexpansive retraction we have

$$\begin{aligned} \|x_n - Tx_n\| &\leq \|x_n - Ty_n\| + \|Tx_n - Ty_n\| \\ &\leq \|x_n - Ty_n\| + \|Ix_n - Iy_n\| \\ &\leq \|x_n - Ty_n\| + \|x_n - y_n\| \\ &\leq \|x_n - Ty_n\| + \|Px_n - P(\alpha'_n x_n + \beta'_n Tx_n + \gamma'_n v_n)\| \\ &\leq \|x_n - Ty_n\| + \|x_n - (\alpha'_n x_n + \beta'_n Tx_n + \gamma'_n v_n)\| \\ &\leq \|x_n - Ty_n\| + \beta'_n \|x_n - Tx_n\| + \gamma'_n \|x_n - v_n\| \\ &\leq \|x_n - Ty_n\| + \beta'_n \|x_n - Tx_n\| + \gamma'_n r \end{aligned}$$

That is  $(1 - \beta'_n) \|x_n - Tx_n\| \leq \|x_n - Ty_n\| + \gamma'_n r$

On taking limit as  $n \rightarrow \infty$  both sides we get  $\lim_{n \rightarrow \infty} \|x_n - Tx_n\| = 0$ . This completes the proof of the lemma.  $\square$

Now we prove our main result.

**THEOREM 2.1.** *Let  $E$  be a uniformly convex Banach space satisfying the Opial's property and let  $K$ ,  $T$  and  $\{x_n\}$  be as in Lemma (2.2). If  $F(T) \cap F(I) \neq \phi$ , then  $\{x_n\}$  converges weakly to a fixed point of  $F(T) \cap F(I)$ .*

**PROOF.** For any  $p \in F(T) \cap F(I)$ , it follows from Lemma (2.1) that

$$\lim_{n \rightarrow \infty} \|x_n - p\|$$

exists. We now prove that  $\{x_n\}$  has a unique weak sub sequential limit in  $F(T)$ . By Lemmas (1.2) and (2.2), we know that  $p \in F(T)$ .

Let  $\{x_{n_k}\}$  and  $\{x_{m_k}\}$  be two sub sequences of  $\{x_n\}$  which converges weakly to  $p$  and  $q$ , respectively. We will show that  $p = q$ .

Suppose that  $E$  satisfies Opial's property and that  $p \neq q$  is in weak limit set of the sequence  $\{x_n\}$ . Then  $\{x_{n_k}\} \rightarrow p$  and  $\{x_{m_k}\} \rightarrow q$ , respectively. Since  $\lim_{n \rightarrow \infty} \|x_n - p\|$  exists for any  $p \in F(T) \cap F(I)$ , then by Opial's property we conclude that

$$\begin{aligned} \lim_{n \rightarrow \infty} \|x_n - p\| &= \\ \lim_{k \rightarrow \infty} \|x_{n_k} - p\| &< \lim_{k \rightarrow \infty} \|x_{n_k} - q\| < \lim_{j \rightarrow \infty} \|x_{m_j} - p\| = \\ \lim_{n \rightarrow \infty} \|x_n - p\| & \end{aligned}$$

a contradiction. This proves that  $\{x_n\}$  converges weakly to a fixed point of  $F(T) \cap F(I)$ . This completes the proof of the theorem.  $\square$

## References

- [1] A. Pazy, *On the asymptotic behaviour of iterates of nonexpansive mappings in Hilbert space*, Israel Journal of Mathematics, **26**(1977), no. 2, 197-204.
- [2] B. E. Rhoades and S. Temir, *Convergence theorems for I-nonexpansive mapping*, to appear in International Journal of Mathematics and Mathematical Sciences.
- [3] F. E. Browder, *Convergence of approximates to fixed points of nonexpansive nonlinear mappings in Banach spaces*, Arch. Rational Mech. Anal. **24**, pp.82-90, 1967.
- [4] H. Kiziltunc and M. Ozdemir, *On convergence theorem for nonself I-nonexpansive mappings in Banach spaces*, Applied Mathematical Sciences, **Vol. 1**, 2007, no.48, 2379-2383.
- [5] J. B. Baillon, *Un theoreme de type ergodique pour les contractions non lineaires dans un espace de Hilbert*, Comptes Rendus de l'Academie des Sciences de Paris, Serie A **280**(1975), no. 22, 1511-1514.
- [6] J. B. Diaz and F. T. Metcalf, *On the set of subsequential limit points of successive approximations*, Transactions of American Mathematical Society **135**(1969), 459-485.
- [7] M. K. Ghosh and L. Debnath, *Convergence of Ishikawa iterates of quasi-nonexpansive mappings*, J. Math. Anal. Appl. **207**(1997), no. 1, 96-103.
- [8] S. Ishikawa, *Generalized I-nonexpansive maps and best approximations in Banach spaces*, Demonstratio Mathematica **37**(2004), no. 3, 597-600.
- [9] S. Temir and O. Gul, *Convergence theorem for I-asymptotically quasi-nonexpansive mapping in Hilbert space*, Journal of Mathematical Analysis and Applications **329**(2007), 759-765.
- [10] W. A. Kirk, *Remarks on approximation and approximate fixed points in metric fixed point theory*, Annales Universitatis Mariae Curie-Sklodowska. Sectio A **51**(1997), no. 2, 167-178.
- [11] W. G. Dotson Jr., *On the Mann iterative process*, Transactions of the American Mathematical Society **149**(1970), no. 1, 65-73.
- [12] W. R. Mann, *Mean value methods in iteration*, Proc. Amer. Math. Soc. **4**(1953), 506-510.
- [13] W. V. Petryshyn and T. E. Williamson Jr., *Strong and weak convergence of the sequence of successive approximations for quasi-nonexpansive mappings*, Journal of the Mathematical Analysis and Applications **43**(1973), 459-497.
- [14] Z. Opial, *Weak convergence of the sequence of successive approximations for nonexpansive mappings*, Bull. Amer. Math. Soc. **73**(1967), 591-597.
- [15] K. K. Tan and H. K. Xu, *Approximating fixed points of nonexpansive mappings by the Ishikawa iteration process*, J. Math. Anal. Appl., **178**(1993), 301-308.
- [16] J. Schu, *Weak and strong convergence to fixed points of asymptotically nonexpansive mappings*, Bull. Austral. Math. Soc. **43**(1991), 153-159.

(Received by editors 20.11.2011, in revised form 15.02.2012; available online 17.02.2012)

DEPARTMENT OF MATHEMATICS, NRI INSTITUTE OF INFORMATION SCIENCE AND TECHNOLOGY, BHOPAL, INDIA-462021

*E-mail address:* pmathsjhade@gmail.com

DEPARTMENT OF MATHEMATICS, J H GOVERNMENT POST GRADUATE COLLEGE, BETUL, INDIA-460001

*E-mail address:* dssaluja@rediffmail.com