

# On the Error Term for the Number of Integral Ideals in Galois Extensions

Sanying Shi\*

School of Mathematics, Hefei University of Technology, Hefei, China  
 \*Corresponding author: vera123\_99@hotmail.com

**Abstract** Suppose that  $E$  is an algebraic number field over the rational field  $\mathbb{Q}$ . Let  $a(n)$  be the number of integral ideals in  $E$  with norm  $n$  and  $\Delta(x)$  denote the remainder term in the asymptotic formula of the  $l$ -th integral power sum of  $a(n)$ . In this paper the bound of the average behavior of  $\Delta(x)$  is given. This result constitutes an improvement upon that of Lü and Wang for the error terms in mean value.

**Keywords:** dedekind zeta-function, dirichlet series, mean value

**Cite This Article:** Sanying Shi, "On the Error Term for the Number of Integral Ideals in Galois Extensions." Turkish Journal of Analysis and Number Theory, vol. 2, no. 6 (2014): 230-232. doi: 10.12691/tjant-2-6-8.

## 1. Introduction and the Result

Let  $E$  be an algebraic number field of degree  $d$  over the rational field  $\mathbb{Q}$ , and  $\zeta(s, E)$  be its Dedekind zeta-function. Thus

$$\zeta(s, E) = \sum_{\mathfrak{a}} \frac{1}{(N\mathfrak{a})^s}, \quad (\Re(s) > 1),$$

where  $\mathfrak{a}$  runs over all integral ideals of the field  $E$ , and  $N\mathfrak{a}$  is the norm of  $\mathfrak{a}$ . If  $a(n)$  denotes the number of integral ideals in  $E$  with norm  $n$ , then we have

$$\zeta(s, E) = \sum_{n=1}^{\infty} \frac{a(n)}{n^s}, \quad s = \sigma + it, \quad \sigma > 1.$$

It is known that  $a(n)$  is a multiplicative function and satisfies

$$a(n) \ll \tau(n)^d, \quad (1)$$

where  $\tau(n)$  is the divisor function.

It is an important problem to study the function  $a(n)$ . In 1927, Landau [7] first proved that

$$\sum_{n \leq x} a(n) = \alpha x + O\left(x^{1-\frac{2}{d+1}+\epsilon}\right),$$

for any arbitrary algebraic number field of degree  $d \geq 2$ , where  $\alpha$  is the residue of  $\zeta(s, E)$  at its simple pole  $s = 1$ .

It is hard to refine Landau's result. Later, Huxley and Watt [3] and Müller [9] improved the results for the quadratic and cubic fields, respectively.

Until 1993, Nowak [10] obtained the best result

$$\sum_{n \leq x} a(n) = \alpha x + \begin{cases} O\left(x^{1-\frac{2}{d}+\frac{8}{d(5d+2)}} (\log x)^{\frac{10}{5d+2}}\right), & \text{for } 3 \leq d \leq 6 \\ O\left(x^{1-\frac{2}{d}+\frac{3}{2d^2}} (\log x)^{\frac{2}{d}}\right), & \text{for } d \geq 7 \end{cases}$$

for any arbitrary algebraic number field of degree  $d \geq 3$ .

In [1], Chandrasekharan and Good studied the  $l$ -th integral power sum of  $a(n)$  in some Galois fields, and they showed that

**Theorem 1.0.** If  $E$  is a Galois extension of  $\mathbb{Q}$  of degree  $d$ , then for any  $\epsilon > 0$  and any integer  $d \geq 2$ , we have

$$\sum_{n \leq x} a(n)^l = x Q_m(\log x) + O\left(x^{1-\frac{2}{md}+\epsilon}\right),$$

where  $m = d^{l-1}$ , and  $Q_l(t)$  is a suitable polynomial in  $t$  of degree  $m-1$ .

Recently, Lü and Wang [8] improved the classical result of [1] by replacing  $\frac{2}{md}$  with  $\frac{3}{md+6}$ .

Motivated by [2,4,5], the purpose of this paper is to study the remainder term in mean square, and we shall prove the following result.

**Theorem 1.1** Subject to assumptions in Theorem 1.0, and define

$$\Delta(x) := \sum_{n \leq x} a(n)^l - x Q_m(\log x). \quad (2)$$

Then we have

$$\int_1^X \Delta^2(x) dx \ll_{\epsilon} X^{3-\frac{6}{md+3}+\epsilon}$$

for any given  $\epsilon > 0$ .

**Notations.** As usual, the Vinogradov symbol  $A \ll B$  means that  $B$  is positive and the ratio  $A/B$  is bounded. The letter  $\epsilon$  denotes an arbitrary small positive number, not the same at each occurrence.

## 2. Proof of Theorem 1.1

To prove our Theorem, we need the following lemmas.

**Lemma 2.1** Let  $E/\mathbb{Q}$  be a Galois extension of degree  $d$ , and  $a(n)$  be defined in (1). Define

$$N_l(s) = \sum_{n=1}^{\infty} \frac{a(n)^l}{n^s}, (\Re s > 1). \tag{3}$$

Then we have

$$N_l(s) = \zeta^m(s, E) \cdot A_1(s),$$

for any integer  $l \geq 1$ , where  $m = d^{l-1}$ , and  $A_1(s)$  denotes a Dirichlet series, which is absolutely and uniformly convergent for  $\Re(s) > 1/2$ .

**Proof.** This is Lemma 2.1 in [8].

**Lemma 2.2.** Let  $E$  be an algebraic number field of degree  $d$ , then

$$\zeta(\sigma + it, E) \ll (1 + |t|)^{\frac{d}{3}(1-\sigma)+\epsilon},$$

for  $\frac{1}{2} \leq \sigma \leq 1 + \epsilon$  and any fixed  $\epsilon > 0$ .

**Proof.** By Lemma 2.2 in [8] and the Phragmen-Lindelöf principle for a strip (see, e.g. Theorem 5.53 in [6]), Lemma 2.2 follows immediately.

Now we begin to prove our theorem.

Let  $E$  be a Galois extension of  $\mathbb{Q}$  of degree  $d$ .

Recall  $a(n)$  denotes the number of integral ideals in  $E$  with norm  $n$ , and

$$\zeta(s, E) = \sum_{n=1}^{\infty} \frac{a(n)}{n^s}, \quad s = \sigma + it, \quad \sigma > 1.$$

Let

$$T = X^{\frac{3}{d^l+3}}.$$

From (1), (3) and Perron's formula (see Proposition 5.54 in [6], we get

$$\sum_{n \leq x} a(n)^l = \frac{1}{2\pi i} \int_{1-\epsilon-iT}^{1+\epsilon+iT} N_l(s) \frac{x^s}{s} ds + O\left(\frac{x^{1+\epsilon}}{T}\right).$$

By the property  $N_l(s)$  only has a simple pole at  $s = 1$  for  $\Re(s) > \frac{1}{2}$  and Cauchy's residue theorem, we have

$$\begin{aligned} \sum_{n \leq x} a(n^2)^l &= \frac{1}{2\pi i} \left\{ \int_{\frac{1}{2}+\epsilon-iT}^{\frac{1}{2}+\epsilon+iT} + \int_{\frac{1}{2}+\epsilon+iT}^{1+\epsilon+iT} + \int_{1+\epsilon+iT}^{1+\epsilon-iT} \right\} N_l(s) \frac{x^s}{s} ds \\ &\quad + \text{Res}_{s=1} N_l(s)x + O\left(\frac{x^{1+\epsilon}}{T}\right) \\ &= xQ_m(\log x) + J_1(x) + J_2(x) + J_3(x) + O\left(x^{1+\epsilon}T^{-1}\right). \end{aligned}$$

where  $m = d^{l-1}$ , and  $Q_m(t)$  is a suitable polynomial in  $t$  of degree  $m-1$ .

From the definition of  $\Delta(x)$  in (2), we have

$$\Delta(x) = J_1(x) + J_2(x) + J_3(x) + O\left(x^{1+\epsilon}T^{-1}\right).$$

Therefore to prove Theorem 1.1, we shall prove the following results.

$$\int_1^X J_i^2(x) dx \ll_{\epsilon} X^{3-\frac{6}{md+3}+\epsilon}, \quad i = 1, 2, 3 \tag{4}$$

and

$$\int_1^X \left( O\left(\frac{x^{1+\epsilon}}{T}\right) \right)^2 dx \ll_{\epsilon} X^{3-\frac{6}{md+3}+\epsilon}. \tag{5}$$

It is easy to get

$$\int_1^X \left( O\left(\frac{x^{1+\epsilon}}{T}\right) \right)^2 dx \ll \frac{X^{3+\epsilon}}{T^2} \ll X^{3-\frac{6}{md+3}+\epsilon}. \tag{6}$$

Now we consider the integral  $J_1(x)$ . We have

$$J_1(x) = \frac{1}{2\pi} \int_{-T}^T N_l\left(\frac{1}{2} + \epsilon + it\right) \frac{x^{\frac{1}{2}+\epsilon+it}}{\frac{1}{2} + \epsilon + it} dt.$$

Then

$$\begin{aligned} \int_1^X J_1^2(x) dx &= \frac{1}{4\pi^2} \int_1^X \left( \int_{-T}^T N_l\left(\frac{1}{2} + \epsilon + it_1\right) \frac{x^{\frac{1}{2}+\epsilon+it_1}}{\frac{1}{2} + \epsilon + it_1} dt_1 \right. \\ &\quad \left. \times \int_{-T}^T \overline{N_l\left(\frac{1}{2} + \epsilon + it_2\right)} \frac{x^{\frac{1}{2}+\epsilon-it_2}}{\frac{1}{2} + \epsilon - it_2} dt_2 \right) dx \\ &= \frac{1}{4\pi^2} \int_{-T}^T \int_{-T}^T \frac{N_l\left(\frac{1}{2} + \epsilon + it_1\right) \overline{N_l\left(\frac{1}{2} + \epsilon + it_2\right)}}{\left(\frac{1}{2} + \epsilon + it_1\right)\left(\frac{1}{2} + \epsilon - it_2\right)} \\ &\quad \times \left( \int_1^X x^{1+2\epsilon+i(t_1-t_2)} dx \right) dt_1 dt_2 \\ &\ll X^{2+2\epsilon} \int_{-T}^T dt_1 \int_{-T}^T \frac{|N_l\left(\frac{1}{2} + \epsilon + it_1\right)| |N_l\left(\frac{1}{2} + \epsilon + it_2\right)|}{(1+|t_1|)(1+|t_2|)(1+|t_1-t_2|)} dt_2 \end{aligned}$$

$$\begin{aligned} &\ll X^{2+2\epsilon} \int_{-T}^T dt_1 \int_{-T}^T \left( \frac{|N_l\left(\frac{1}{2} + \epsilon + it_1\right)|^2}{(1+|t_1|)^2} \right. \\ &\quad \left. + \frac{|N_l\left(\frac{1}{2} + \epsilon + it_2\right)|^2}{(1+|t_2|)^2} \right) \frac{dt_2}{1+|t_1-t_2|} \quad (7) \\ &\ll X^{2+2\epsilon} \int_{-T}^T \frac{|N_l\left(\frac{1}{2} + \epsilon + it_1\right)|^2}{(1+|t_1|)^2} dt_1 \int_{-T}^T \frac{dt_2}{1+|t_1-t_2|}. \end{aligned}$$

To go further, we get

$$\begin{aligned} \int_{-T}^T \frac{dt_2}{1+|t_1-t_2|} &\ll \int_{t_1-1}^{t_1+1} dt_2 + \left( \int_{t_1+1}^T + \int_{-T}^{t_1-1} \right) \frac{dt_2}{|t_1-t_2|} \\ &\ll 1 + \int_{t_1+1}^T \frac{dt_2}{t_1-t_2} \quad (8) \\ &\ll \int_1^{T+|t_1|} \frac{dt}{t} \ll \log 2T. \end{aligned}$$

By (7) and (8)

$$\int_1^X J_1^2(x) dx \ll X^{2+3\epsilon} \int_{-T}^T \frac{|N_l\left(\frac{1}{2} + \epsilon + it_1\right)|^2}{(1+|t_1|)^2} dt_1. \quad (9)$$

From (9), Lemma 2.1 and 2.3, we have (for  $d \geq 3$ )

$$\begin{aligned} \int_1^X J_1^2(x) dx &\ll X^{2+3\epsilon} + X^{2+3\epsilon} \int_1^T \left| \zeta^{d^{l-1}} \left( \frac{1}{2} + \epsilon + it, E \right) \right. \\ &\quad \left. \times A_1 \left( \frac{1}{2} + \epsilon + it_1 \right) \right|^2 t^{-2} dt \\ &\ll X^{2+3\epsilon} + X^{2+3\epsilon} \int_1^T \left| \zeta^m \left( \frac{1}{2} + \epsilon + it, E \right) \right|^2 t^{-2} dt \quad (10) \\ &\ll X^{2+3\epsilon} + X^{2+3\epsilon} \int_1^T \left( \frac{md}{t^6} + \epsilon \right)^2 t^{-2} dt \\ &\ll X^{2+3\epsilon} + X^{2+4\epsilon} T^{\frac{md}{3}-1} \\ &\ll X^{3-\frac{6}{md+3}+\epsilon}. \end{aligned}$$

Finally we estimate trivial bounds of the integrals  $J_2(x), J_3(x)$ . By Lemma 2.2, we get

$$\begin{aligned} J_2(x) + J_3(x) &\ll \int_{\frac{1}{2+\epsilon}}^{1+\epsilon} x^\sigma |\zeta^m(\sigma + iT, E)| T^{-1} d\sigma \\ &\ll \max_{1/2+\epsilon \leq \sigma \leq 1+\epsilon} x^\sigma T^{\frac{md}{3}(1-\sigma)+\epsilon} T^{-1} \\ &= \max_{\frac{1}{2+\epsilon} \leq \sigma \leq 1+\epsilon} \left( \frac{x}{T^{md/3}} \right)^\sigma T^{\frac{md}{3}-1+\epsilon} \\ &\ll \frac{x^{1+\epsilon}}{T} + x^{\frac{1}{2+\epsilon}} T^{\frac{d^l}{6}-1+\epsilon}, \end{aligned}$$

which yields

$$\begin{aligned} \int_1^X (J_2(x) + J_3(x))^2 dx &\ll \int_1^X \left( \frac{x^{1+\epsilon}}{T} + x^{\frac{1}{2+\epsilon}} T^{\frac{md}{6}-1+\epsilon} \right)^2 dx \\ &\ll \int_1^X \left( \frac{x^{1+\epsilon}}{T} \right)^2 dx \quad (11) \\ &\ll \frac{X^{3+\epsilon}}{T^2} + X^{2+2\epsilon} T^{\frac{md}{3}-2+2\epsilon} \\ &\ll X^{3-\frac{6}{md+3}+\epsilon}. \end{aligned}$$

The inequalities (4), (5) immediately follow from (6), (10) and (11). That is,

$$\int_1^X \Delta^2(x) dx \ll_\epsilon X^{3-\frac{6}{md+3}+\epsilon}.$$

Then this completes the proof of Theorem 1.1.

### Acknowledgement

This work is supported by The National Science Foundation of China (grant no. 11201107 and 11071186) and by Natural Science Foundation of Anhui province (Grant No. 1208085QA01).

### References

- [1] Chandrasekharan K., Good A., On the number of integral ideals in Galois extensions, *Monatsh. Math.*, 95, 99-109. 1983.
- [2] Heath-Brown D. R., *The number of Abelian groups of order at most x*, Journtes Arithmttiques, Luminy 1989.
- [3] Huxley M. N., Watt N., The number of ideals in a quadratic field II, *Israel J. Math.* Part A, 120, 125-153. 2000.
- [4] Ivic A., The number of finite non-isomorphic Abelian groups in mean square, *Hardy-Ramanujan J.*, 9, 17-23. 1986.
- [5] Ivic A., On the Error Term for the Counting Functions of Finite Abelian Groups, *Monatsh. Math.* 114, 115-124. 1992.
- [6] Iwaniec H., Kowalski E., *Analytic Number Theory*, Amer. Math. Soc. Colloq. Publ., 53, 204-216. 2004.
- [7] Landau E., *Einführung in die elementare und analytische Theorie der algebraischen Zahlen und der Ideale*, Teubner, 1927.
- [8] Lü G., Wang Y., Note on the number of integral ideals in Galois extension, *Sci. China Ser. A*, 53, 2417-2424. 2010.
- [9] Müller W., On the distribution of ideals in cubic number fields, *Monatsh. Math.*, 106, 211-219. 1988.
- [10] Nowak W.G., On the distribution of integral ideals in algebraic number theory fields, *Math. Nachr.*, 161, 59-74. 1993.